

J. C. Türp · F. Komine · A. Hugger

Efficacy of stabilization splints for the management of patients with masticatory muscle pain: a qualitative systematic review

Received: 22 December 2003 / Accepted: 5 April 2004 / Published online: 4 June 2004
© Springer-Verlag 2004

Abstract This study aimed at providing an answer to two clinical questions related to patients with masticatory muscle pain: 1) Does the use of a full-coverage hard acrylic occlusal appliance (stabilization splint) lead to a significant decrease of symptoms? and 2) Is the treatment success achieved with a stabilization splint more pronounced than the success attained with other forms of treatment (including placebo treatment) or no treatment? A systematic search was carried out in different electronic databases, supplemented by handsearch in four selected dental journals and by examination of the bibliographies of the retrieved articles. Thirteen publications, representing nine controlled clinical studies, could be identified. Reporting quality of most studies as assessed with the Jadad score ranged from 1 to 5. Based on the currently best available evidence it appears that most patients with masticatory muscle pain are helped by the incorporation of a stabilization splint. Nevertheless, evidence is equivocal if improvement of pain symptoms after incorporation of the intraoral appliance is caused by a specific effect of the appliance. A stabilization splint does not appear to yield a better clinical outcome than a soft splint, a non-occluding palatal splint, physical therapy, or body acupuncture. The scarcity of current external evidence emphasizes the need for more and better clinical research.

Keywords Evidence-based medicine · Myofascial pain · Occlusal splints · Systematic review · Temporomandibular joint disorders

Introduction

Muscle pain has been known for some time to be the leading cause of discomfort in the head and neck area [5], and masticatory muscle pain (myofascial pain) is the most common diagnosis among the various conditions encompassed by the term *temporomandibular disorders* (TMDs) [58, 81]. For about 40 years [76], stabilization splints have been one of the preferred modalities in the management of TMDs [72], although a great variety of other treatments are currently in use among clinicians [3, 96].

In the early 1990s, David Sackett and his colleagues from McMaster University, Hamilton (Ontario, Canada), introduced the concept of evidence-based medicine (EBM), “the conscientious, explicit and judicious use of current best evidence in making decisions about the care of individual patients” [82]. The practice of EBM relies on three equally important elements: the best research evidence from a systematic search of the literature (mostly from patient-centered clinical research), individual clinical expertise (clinical skills and past experience), and patient values (preferences, concerns, expectations) [83]. The application of EBM comprises several steps: formulation of an important and answerable clinical question derived from a clinical problem; selection of the most appropriate information source(s) and the most appropriate search strategy; and appraisal and application of the evidence found. In many editorials and articles, the introduction of EBM into dentistry has been encouraged. However, the incorporation of EBM into clinical decision-making depends heavily on the availability of external clinical evidence.

Over the past two decades, a few reviews about the effect of occlusal appliances for the management of TMDs have been published (e.g., [2, 10, 11, 30, 31, 52]). In these

J. C. Türp (✉)
Clinic for Reconstructive Dentistry
and Temporomandibular Disorders,
Dental School, University of Basel,
Hebelstr. 3, CH-4056 Basel, Switzerland
e-mail: jens.tuerp@unibas.ch
Tel.: +41-61-2672631
Fax: +41-61-2672660

F. Komine
Nihon University School of Dentistry,
Department of Crown and Bridge Prosthodontics,
1-8-13, Kanda-Surugadai, Chiyoda-Ku, 101-8310 Tokyo, Japan

A. Hugger
Department of Prosthodontics,
Dental School, University of Düsseldorf,
Moorenstr. 5, D-40225 Düsseldorf, Germany

publications, however, no distinction among specific TMD diagnoses such as masticatory muscle pain was made. The present paper, therefore, focuses on the most common clinical scenario in the management of TMDs by dentists: the management of a patient, who suffers from pain in the masticatory muscles, with a bite splint. Specifically, by systematically searching the literature, we want to provide an answer to the following two clinical questions. In patients with masticatory muscle pain:

1. Does the use of a full-coverage hard acrylic occlusal appliance (stabilization splint, Michigan splint) lead to a significant decrease of symptoms?
2. Is symptom improvement achieved with a stabilization splint more pronounced than the success attained with other forms of management (including placebo treatment) or no treatment?

Considering the great number of patients suffering from myofascial face pain, an answer to our questions would be of considerable clinical relevance.

Methods

To identify all relevant articles and dental congress abstracts of randomized controlled clinical trials (RCTs), in which stabilization splint therapy was used and compared to no therapy or concurrent treatments, the following information sources and search strategies were used:

Search in electronic databases

- *Ovid Medline* (online database). The subject terms included in the search as well as the search strategy are listed in Table 1.
- *Cochrane Library* (online database). The search term was “splint.”
- *ISI Web of Science* (Science Citation Index Expanded) (online database). The option “Full Search” was chosen. The keywords included in the search were “bite splint,” “occlusal appliance,” “occlusal splint” and “splint.”
- *Japana Centra Revuo Medicina* (CD-ROM). The search in this database was carried out in a similar way as in Medline. The following Japanese terms were used:
 - Randomized controlled trials: *Musakui taisyo shiken*, *Musakui hikaku shiken*
 - Clinical trials: *Rinsyo shiken*
 - Random allocation: *Randamu waritsuke*
 - Double-blind method: *Nijuu mouken shiken*

- Controlled clinical trials: *Hikaku rinsho shiken*, *Hikaku taisho rinsho shiken*
- Myofascial pain dysfunction (MPD) syndrome: *Kinmaku totsui kinoushugai shoukougoun*, *MPD shoukougoun*
- Temporomandibular joint (TMJ) dysfunction syndrome: *Gaku-kansetsusho*, *Gaku-kinouijo*, *Gaku-kinoushugai*, *Tougai kakushugai*
- Masticatory muscle: *Soshaku-kin*
- Myofascial pain syndrome: *Kinmaku-totsui shoukougoun*, *Kinmakutsui shoukougoun*
- Pain: *Totsui*
- Bite splint: *Baito supurinto*, *Baito purein*, *Kougou kyojo*
- Occlusal appliance: *Okuruuzaru apuriansu*
- Occlusal splint: *Okuruuzaru supurinto*
- Splint: *Supurinto*

Handsearch in selected journals

The most important, peer-reviewed journals of Austria, France, and Germany, which are currently not included in Medline, namely *Actualités odonto-stomatologiques* (Medline listing discontinued in 1991), *Stomatologie* (formerly—until 1996—*Zeitschrift für Stomatologie*; discontinued in Medline in 1990), and *Deutsche Zahnärztliche Zeitschrift* (listing in Medline discontinued in 1992), were handsearched through December 2003.

In addition, the abstracts published in the special issues of the *Journal of Dental Research*, which relate to the annual General Session and Exhibition of the International Association for Dental Research (IADR), were reviewed for the years 1990–2003.

Bibliography search of the identified publications and reviews

The references listed in the relevant articles were perused to identify additional publications pertinent to our clinical question. In addition, the reference lists of relevant review articles were checked.

Inclusion/exclusion criteria

Only trials in which patients had explicitly been diagnosed with masticatory muscle (myofascial) pain were considered. Studies in which additional diagnoses were allowed (e.g., TMJ arthralgia/osteoarthritis; disk interference disorders), were excluded from further analysis. Similarly, articles in which unspecific terminology was used to characterize the investigated patient samples (e.g., “mandibular dysfunction”, “[TMJ] pain dysfunction syndrome” or “temporomandibular disorders”) were not considered because these terms may also include TMJ-related conditions.

There were neither age restrictions of study participants nor language restrictions for inclusion. The last update of the search was made on 11 December 2003.

Table 1 Search strategy and results in Ovid Medline (date of the search: 11 December 2003)

	Search terms	Hits	Relevant
#1	trial\$.mp. or Clinical Trials/ or Randomized Controlled Trials/	328835	–
#2	Temporomandibular Joint Disorders/ or Masticatory Muscles/ or Craniomandibular Disorders/ or Temporomandibular Joint Dysfunction Syndrome/ or myofascial pain dysfunction syndrome.mp. or Myofascial Pain Syndromes/	15178	–
#3	Occlusal Splints/ or bite splint\$.mp. or Splints/	6140	
#1 AND #3		151	[17, 28, 98]
#2 AND #3		964	[7, 17, 28, 47, 80, 86, 98]

Assessment of reporting quality

The reporting quality of the identified articles of RCTs was assessed independently by two reviewers (J.C.T., A.H.). For this purpose, the quality score developed by Jadad et al. [46] was used. The Jadad scale consists of five items which focus on three dimensions of internal validity (randomization; double blinding; description of withdrawals and drop-outs). Uncertainties on data interpretation and discrepancies in scoring the reporting quality were resolved by discussion between the two reviewers.

Results

Altogether, 13 relevant publications, representing nine clinical studies, were identified by the two reviewers. The articles of six studies were published in English, the rest in Dutch [98], German [86], and Japanese [84]. Two studies were carried out in Sweden, two in the USA, one in Canada, one in Germany, one in Italy, one in Japan, and one in the Netherlands. Only two trials [91, 98] were conducted over an observation period of at least 1 year.

The results of the search in Ovid Medline are displayed in Table 1. Seven articles listed in this database are relevant to our question [7, 17, 28, 47, 80, 86, 98].

The search in the Japanese database yielded one hit [84]. The searches in the Cochrane Library and in ISI Web of Science identified no additional publications.

The handsearch in the special issues of the *Journal of Dental Research* yielded five meeting abstracts [42, 43, 44, 90, 91]. All five abstracts referred to the same prospective trial. In the following, only the two most recently published abstracts, which complement each other, are considered [44, 91].

The bibliography search of the identified publications identified no additional publication. (In one article [78], a paper from an Argentinean dental journal was cited [79] in which it was allegedly reported "that splint therapy associated with diazepam, in this order, produced more effective TMD pain relief when these therapies were applied exclusively." In the cited article by Roldan et al. [79], however, bite splints were not mentioned.)

The major characteristics (patients, inclusion and exclusion criteria, treatment groups, number of participants, study duration, outcome variables for treatment success) and findings (improvement of signs and symptoms, authors' conclusions, reviewers' comments) of the publications are summarized in Tables 2 and 3. In all but one study [86], stabilization splints were fabricated in the maxilla. Masticatory muscle pain was an outcome variable in every study. In all but two trials [7, 17], functional parameters such as mandibular mobility and TMJ sounds were also considered. In three articles [28, 47, 86], Helkimo's Clinical Dysfunction Index (CDI) [38] was used.

The methodology of the clinical investigations differed in several important aspects:

1. Recruitment of study participants: in all but two trials, study participants were restricted to patients seeking

care at or referred to clinical centers specialized for the diagnosis and management of orofacial pain. In the remaining two studies, participants were exclusively [80] or partly [17] recruited by notices in local newspapers or journals.

2. Number of participants: the number of participants included in most studies was small. The total number of participants lay between 26 [86] and 168 individuals [91] (Table 2).
3. Description of randomization/blinding: the method of randomization was described only in two publications [28, 86]. Five studies [17, 28, 47, 80, 98] used a blinded design: the examiner who evaluated the treatment was blind to the type of treatment the patient received. Lack of (double-)blinding leads to an overestimation of the treatment effect [85]. (It is controversial whether or not double-blinding is possible in trials comparing active and "placebo" appliances.) The publication by Ekberg et al. [28] is the only one with an appropriate description of both randomization and blinding, and with a Jadad score of 5 (Table 4). (It has to be considered that the reported information about the randomized studies of Huggins/Truelove et al. [44, 91] as well as Sakuma et al. [84] was limited in the available meeting abstracts.)
4. Appliance use: in three trials [17, 80, 86], the appliance was worn (nearly) 24 h/day. In four other studies, the splint was worn only at night [7, 28, 47, 98]. In the published meeting abstracts [44, 84, 91], no pertinent information was given.
5. Treatment provided in the control group(s): Michigan splints were compared with the following alternative approaches:
 - Non-occluding palatal appliance [17, 28, 80, 84]
 - Anteriorly occluding maxillary splint [86]
 - Full-covering maxillary soft appliance [44, 91]
 - Occlusal adjustment [98]
 - Physical therapy [98]
 - Body acupuncture [47]
 - Different treatments [44, 91]
 - No treatment [7, 47]

In one study [80], physical therapy and verbal support were given to each patient included in the trial.

Based on the results described in the identified publications, our two clinical questions can be answered as follows:

1. Management of myofascial face pain with a stabilization splint worn at night is likely to lead to a statistically significant short-term improvement when compared with no treatment [47].
2. Current evidence is inconclusive about the question of whether the observed improvement during and after stabilization therapy is greater than the one achieved by a non-occluding palatal appliance (i.e., a "placebo" splint). In two recently conducted trials [28, 84], there was a statistically more significant decrease of pain and functional impairment in the group that received a

Table 2 Characteristics of RCTs in which stabilization appliance therapy in patients with masticatory muscle pain was investigated

Study	Patients	Inclusion criteria	Exclusion criteria	Treatment groups	n	Study duration
Rubioff <i>et al.</i> [80]	Patients responding to a notice in a newspaper (Buffalo, New York, USA)	<ul style="list-style-type: none"> Diagnosis of MPD: "a complaint of facial pain and one or more of the following: limited jaw opening, joint sounds, deviation on opening, and tenderness on muscle palpation." 	<ul style="list-style-type: none"> Clinical or radiological evidence of TMJ pathology Complete upper or lower denture Recent major occlusal changes Third molar problems (<i>e.g.</i>, pericoronitis) Previously worn a appliance Alternative diagnosis 	(1) Maxillary stabilization appliance (24h/day) (2) Non-occluding palatal appliance (24h/day) <ul style="list-style-type: none"> Additional treatment in both groups: <ul style="list-style-type: none"> support by the dental therapist physical therapy (moist heat; daily home exercise) 	(1) 15 (2) 11/15	6 weeks
Siebert & Gundlach [86]	From the clinic population of the Dental School, University of Hamburg (Germany)	<ul style="list-style-type: none"> Myofascial pain Possibly: No satisfactory pain reduction by previous patient education and physical therapy (n=26/72) 	<ul style="list-style-type: none"> Evidence of TMJ pathology 	(1) Mandibular (I) stabilization appliance (use: as often as possible) (2) Maxillary anterior appliance (Sved appliance; "relaxing appliance") (use: as often as possible)	(1) 12 (2) 12/14	2.2 months (average)
Johansson <i>et al.</i> [47]	Consecutive series of patients referred to the Department of Stomatognathic Physiology, Faculty of Odontology, University of Gothenburg (Sweden), for TMD treatment	<ul style="list-style-type: none"> History including signs and symptoms of TMD Complaints of headache and/or facial pain Clinical examination demonstrating tenderness to palpation in the masticatory muscles Complete natural dentition (single crown permitted) 	<ul style="list-style-type: none"> Individuals with psychologic / psychogenic factors, trauma, surgery, or systemic joint, muscle, or skin diseases influencing the symptoms Radiologic evidence of TMJ, facial skeleton, or tooth pathology Previous acupuncture or stomatognathic therapy for the treatment of the disorder in the individuals selected 	(1) Maxillary stabilization appliance (at night) (2) Body acupuncture (3) No treatment	(1) 15 (2) 15 (3) 15	(1) 3 months (2) 3 months (3) 2 months

Table 2 (continued)

Study	Patients	Inclusion criteria	Exclusion criteria	Groups	n	Study duration
Dao <i>et al.</i> [17]	(a) Recruited through announcements published in local journals; or (b) referred by dentists to the research clinic, University of Montreal of Montreal (Canada)	<ul style="list-style-type: none"> Men and women between 16 and 45 years seeking treatment Chief complaint of frequent pain (at least four times/week) in the masticatory muscles of at least 12 weeks duration Positive report of tenderness to palpation of at least three sites in the masticatory muscles 	<ul style="list-style-type: none"> Clinical or radiological evidence of TMJ pathology Previous occlusal treatment with or without appliances Complete dentures or removable partial dentures with distal extensions Metabolic disease Neurologic disorders Vascular disease Neoplasia History of psychiatric disorders History of drug abuse Recent facial or cervical trauma Currently receiving medication or other treatments 	(1) Stabilization appliance (24h/day) (2) Stabilization appliance (restricted use: 30 minutes at each appointment = passive control) (3) Non-occluding palatal appliance (= active control) (24h/day)	(1) 22 (2) 19/20 (3) 20/21	2 weeks baseline + 8 weeks treatment
Cane <i>et al.</i> [7]	Patients referred to the Physical Medicine and Rehabilitation Service, S. Giovanni-Molinette Hospital, Turin (Italy)	<ul style="list-style-type: none"> Cervical pain Contractures of the neck muscles Reduction of cervical mobility Masticatory muscle pain (n=33) 	<ul style="list-style-type: none"> Clinical or radiological evidence of arthrosis Previous treatment for pain in the neck region History of trauma to the cervical region Cervical pain related to tumor Congenital malformation of the cervical-occipital junction 	(1) Maxillary stabilization appliance (night only) (2) No treatment	(1) 18 (2) 15	2 months

Table 2 (continued)

Study	Patients	Inclusion criteria	Exclusion criteria	Groups	n	Study duration
Truelove <i>et al.</i> [91]	Clinic patients, Department of Oral Medicine, University of Washington, Seattle (USA)	<ul style="list-style-type: none"> RDC/TMD diagnosis of myofascial pain 		(1) Stabilization appliance (use: not reported) (2) Soft appliance (use: not reported) (3) Conservative treatment without a appliance	(1) 65 (2) 55 (3) 48	12 months
Huggins <i>et al.</i> [44]	“	“		“	(1) 47 (2) 38 (3) 33	“
van der Glas <i>et al.</i> [98]	Clinic patients, University Medical Center Utrecht (Netherlands)	<ul style="list-style-type: none"> At least two months of myogenous TMD complaints Between 18 and 65 years old 		<ul style="list-style-type: none"> Patients without occlusal interferences: <ol style="list-style-type: none"> Maxillary appliance (use: evening and night) Physical therapy Patients with pronounced occlusal interferences: <ol style="list-style-type: none"> Maxillary appliance (use: evening and night), followed by (not earlier than 6 weeks later) occlusal adjustment (combination therapy) Occlusal adjustment 	(1) 35 (2) 36 (1) 25	(average) 1 year 24 weeks 1 year 16 weeks 1 year 27 weeks
Ekberg <i>et al.</i> [28]	Patients referred for (and requesting) TMD treatment to the Department of Stomatognathic Physiology, Faculty of Odontology, Malmö University (Sweden)	<ul style="list-style-type: none"> Diagnosis of myofascial pain with or without limited opening, according to the RDC/TMD Self-assessed myofascial pain of at least 40 mm on a 100-mm VAS 	<ul style="list-style-type: none"> TMJ pain Previous TMD treatment Complete dentures History of psychiatric disorders Symptoms related to disease in other components of the stomatognathic system 	(1) Maxillary stabilization appliance (use: not reported) (2) Non-occluding palatal appliance (= active control)	(2) 22 (1) 30 (2) 30	1 year 18 weeks 10 weeks
Sakuma <i>et al.</i> [84]	Clinic patients, Atchigakuin University Dental Hospital (Japan)	<ul style="list-style-type: none"> Masticatory muscle pain 		(1) Stabilization appliance (use: not reported) (2) Non-occluding palatal appliance	(1) 20 (2) 20	12 weeks

TMJ temporomandibular joint, MPD myofascial pain dysfunction, TMD temporomandibular disorders, CMD craniomandibular disorders, RDC/TMD Research Diagnostic Criteria for Temporomandibular Disorders [22], CDI Clinical Dysfunction Index [38], VAS visual analog scale, (?) not explicitly reported in the paper

Table 3 Characteristics of studies in which stabilization appliance therapy in patients with masticatory muscle pain was investigated

Study	Outcome variables for treatment success	Improvement of signs	Improvement of symptoms	Authors' conclusion	Reviewers' comments
Rubinoﬀ <i>et al.</i> [80]	<p><i>Signs</i></p> <ul style="list-style-type: none"> - Maximum interincisal distance - Mandibular deviation - Joint sounds - Tenderness on muscle palpation - Tenderness on TMJ palpation <p><i>Symptoms</i></p> <ul style="list-style-type: none"> - Pain intensity (pain diary) - Subjective symptom improvement (treatment success) 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant for stabilization group only: joint sounds, palpation <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • No statistically significant differences between the groups 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant for both groups: pain intensity <p><i>Specific symptoms</i></p> <p><u>Symptom improvement</u> (= moderate, great, or complete)</p> <p><i>Treatment groups</i></p> <p>(1) 87% (13/15)</p> <p>(2) 77% (10/13)</p> <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • No statistically significant differences between the groups 	<p>Average pain intensity on a scale from 0 to 5 (0 "no pain at all"; 5 "most intense pain ever experienced") after treatment: stabilization appliance group 0,73; non-occluding appliance group 0,98.</p> <p>"In summary, none of the measures showed significant differences in treatment effectiveness between the groups that received the occluding splint and the group that received the nonoccluding splint."</p> <p>"...both conventional and non-occluding appliances seem to relieve symptoms equally."</p>	<ul style="list-style-type: none"> • Randomization of patients. • Blinded evaluation: clinical examiner unaware of patient group affiliation. • Small sample size. • Significantly lower pretreatment pain levels in the stabilization appliance group at the beginning of the study: average pain intensity on a scale from 0 to 5: stabilization appliance group 1,27 (n=15); non-occluding appliance group 2,07 (n=11). [cf. 61]
Siebert & Gundlach [86]	<p><i>Signs</i></p> <ul style="list-style-type: none"> - Muscle palpation - CDI <p><i>Symptoms</i></p> <ul style="list-style-type: none"> - Masticatory muscle pain 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Symptom improvement ("treatment success") in both treatment groups <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • Statistically greater symptom improvement in the stabilization appliance group (1) than in the anterior appliance group (2) <p><i>Treatment groups</i></p> <p>(1) 92% (11/12)</p> <p>(2) 42% (5/12)</p>	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Symptom improvement ("treatment success") in both treatment groups <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • Statistically greater symptom improvement in the stabilization appliance group (1) than in the anterior appliance group (2) <p><i>Treatment groups</i></p> <p>(1) 92% (11/12)</p> <p>(2) 42% (5/12)</p>	<p>"In spite of the small number of patients the statistical evaluation of the results showed stabilizing appliances to be significantly (p<0.03) superior to relaxing appliances in the treatment of myofascial [sic] pain."</p>	<ul style="list-style-type: none"> • Randomization of patients. • No blinded evaluation. • Small sample size. • Significantly lower pretreatment CDI score (points and categories) in the anterior appliance group at the beginning of the study.

Table 3 (continued)

Study	Outcome variables for treatment success	Improvement of signs	Improvement of symptoms	Authors' conclusion	Reviewers' comments
Johansson <i>et al.</i> [47]	<p><i>Signs</i></p> <ul style="list-style-type: none"> - Mandibular mobility - Pain during mandibular movement - Mandibular deviation - Joint sounds - Tenderness on muscle palpation - Tenderness on TMJ palpation - Occlusal condition - CDI [38] <p><i>Symptoms</i></p> <ul style="list-style-type: none"> - Subjective dysfunction - Pain intensity - Subjective symptom improvement 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant in both treatment groups: CDI, tenderness to muscle palpation <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • Statistically not significant between treatment groups • Statistically significant improvement in both treatment groups vs. no treatment: CDI 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant in both treatment groups <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • Statistically not significant between treatment groups • Statistically significant improvement in both treatment groups vs. no treatment <p><i>Specific Symptom: Subjective dysfunction Treatment groups</i></p> <p>(1) 86% (2) 90% (3) < 15%</p>	<p>“The two examined treatment modalities have quite different ways of influencing the stomatognathic system..., but with practically the same success rate as measured by the subjective and clinical variables.”</p>	<ul style="list-style-type: none"> • Randomization of patients. • No blinded evaluation.
Dao <i>et al.</i> [17]	<p><i>Symptoms</i></p> <ul style="list-style-type: none"> - Pain intensity at rest and after chewing on wax - Pain unpleasantness at rest and after chewing on wax - Quality of life 	<p>– not recorded –</p>	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant in all treatment groups for all 5 outcome variables <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • No statistically significant differences found 	<p>“There were no significant treatment effects... for any of the variables under study. However, there was a general reduction in the pain ratings during treatment and an improvement in the quality of life.”</p> <p>“This study casts doubt on the therapeutic value of oral appliances.”</p>	<ul style="list-style-type: none"> • Randomization of patients. • Blinded evaluation.

Table 3 (continued)

Study	Outcome variables for treatment success	Improvement of signs	Improvement of symptoms	Authors' conclusion	Reviewers' comments
Cane <i>et al.</i> [7]	Signs (for masticatory muscles) <ul style="list-style-type: none"> - Presence of masticatory muscle pain and tension on palpation of the muscle belly and its insertion 	Within-treatment-groups pre-post-improvement <ul style="list-style-type: none"> • Reduction in the splint group only (5/18), not in the control group (0/15). Odds ratio: 11.5 (95% confidence interval: 0.6-231.0) Between-groups post-treatment differences <ul style="list-style-type: none"> • Statistically (marginally) significant differences 	– not recorded –		<ul style="list-style-type: none"> • Randomization of patients. • No blinded evaluation. • All patients suffered from cervical problems. • The outcome of this study does not add much to answer our question.
Truelove <i>et al.</i> [91] / Huggins <i>et al.</i> [44]	Signs <ul style="list-style-type: none"> - Muscle and TMJ palpation - Joint sounds - Maximum jaw opening Symptoms <ul style="list-style-type: none"> - Pain level - TMJ sounds - Eating difficulty - Tinnitus - Clenching/bruxism - Jaw locking/catching 	Within-treatment-groups pre-post-improvement <ul style="list-style-type: none"> • Statistically significant in the treatment groups Between-groups post-treatment differences <ul style="list-style-type: none"> • No statistically significant differences 	Within-treatment-groups pre-post-improvement <ul style="list-style-type: none"> • Statistically significant in the treatment groups Between-groups post-treatment differences <ul style="list-style-type: none"> • No statistically significant differences 	<p>“These data suggest that treatments using lower cost alternatives... to hard appliances provide levels of pain control and symptom reduction equivalent to more costly appliance therapy even over an extended period of time.”</p> <p>“These data indicate that for the long term clinical outcomes examined, neither the more costly [flat plane hard acrylic] appliance nor the less expensive [soft vinyl athletic mouthguard] is superior to non-appliance therapy.”</p>	<p>Only meeting abstract available.</p> <ul style="list-style-type: none"> • Randomization of patients. • No blinded evaluation (?)

Table 3 (continued)

Study	Outcome variables for treatment success	Improvement of signs	Improvement of symptoms	Authors' conclusion	Reviewers' comments
van der Glas <i>et al.</i> [98]	<p><i>Signs</i></p> <ul style="list-style-type: none"> - Pain intensity during mandibular movements - traction of the TMJs - palpation of the masticatory muscles and TMJs - clenching in centric occlusion and in eccentric mandibular position <p><i>Symptoms</i></p> <ul style="list-style-type: none"> - Pain intensity - Pain frequency - Masticatory muscle stiffness/tiredness - Limited jaw opening - Neck and shoulder pain - Pain in front of the ear - Headache - TMJ sounds 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant in the treatment groups <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • No significant differences between the therapies 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant in the treatment groups <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • No significant differences between the therapies 	<p>(a) Patients with low level myogenous TMD signs and symptoms: "Counseling [...] will most likely eliminate any further need for treatment [...]".</p> <p>(b) In more severe cases: "physiotherapy might be preferred as a starting option". Advantages as compared to appliance therapy: 1. similar efficacy; 2. shorter treatment duration; 3. lower costs.</p> <p>(c) In patients with pronounced occlusal interferences: occlusal adjustment. Advantages as compared to appliance therapy: 1. similar therapy efficacy; 2. shorter treatment duration; 3. lower costs.</p>	<ul style="list-style-type: none"> • Randomization of patients. • Blinded evaluation.

Table 3 (continued)

Study	Outcome variables for treatment success	Improvement of signs	Improvement of symptoms	Authors' conclusion	Reviewers' comments
Ekberg <i>et al.</i> [28]	<p><i>Signs</i></p> <ul style="list-style-type: none"> - Mandibular mobility - Pain during non-guided mandibular movements - Masticatory muscle and TMJ tenderness on palpation - CDI [38] <p><i>Symptoms</i></p> <ul style="list-style-type: none"> - Pain duration, frequency, intensity - Pain at rest and during mandibular movements - Improvement of overall subjective symptoms 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant in the stabilization appliance group only <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • Maximum jaw opening, masticatory muscle tenderness, CDI: statistically significant decrease in the appliance group as compared to the control group 	<p><i>Within-treatment-groups pre-post-improvement</i></p> <ul style="list-style-type: none"> • Statistically significant in both groups, but more pronounced in the stabilization appliance group <p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • Pain frequency and intensity, pain during mandibular movements, improvement of overall subjective symptoms: statistically significant decrease in the appliance group as compared to the control group 	<p>“the stabilization appliance can be recommended as a short-term treatment modality for TMD of mainly myogenous origin.”</p>	<ul style="list-style-type: none"> • Randomization of patients. • Blinded evaluation. • High pre-treatment pain intensity.
Sakuma <i>et al.</i> [84]	<p><i>Signs</i></p> <ul style="list-style-type: none"> - Maximum jaw opening <p><i>Symptoms</i></p> <ul style="list-style-type: none"> - Pain level 	<p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • Statistically significant 	<p><i>Between-groups post-treatment differences</i></p> <ul style="list-style-type: none"> • Statistically significant 	<p>“It is suggested that in patients with masticatory muscle pain, stabilization splints can be an effective therapy.” [Authors' translation from the original article]</p>	<p>Only meeting abstract available.</p> <ul style="list-style-type: none"> • Randomization of patients. • No blinded evaluation (?)

TMJ temporomandibular joint, MPD myofascial pain dysfunction, CDI Clinical Dysfunction Index [38], (?) not explicitly reported in the paper

Table 4 Assessment of the quality of the studies by using the quality score proposed by Jadad et al. [46]

Study	Study described as randomized?	Method described and appropriate?	Study described as (double) blind?	Method described and appropriate / inappropriate?	Description of withdrawals / dropouts?	Jadad score
Rubioff <i>et al.</i> [80]	Yes	Not described	Yes	Yes	Yes (n=4)	4
Siebert & Gundlach [86]	Yes	Yes	No	–	Yes (n=2)	3
Johansson <i>et al.</i> [47]	Yes	Not described	Yes	–	Yes (n=0)	3
Dao <i>et al.</i> [17]	Yes	Not described	Yes	Yes	Yes (n=2)	4
Cane <i>et al.</i> [7]	Yes	Not described	No	–	No withdrawal or dropout in the muscle group	2
Truelove <i>et al.</i> [91] / Huggins <i>et al.</i> [44]	Yes	Not described	No	–	No	1
van der Glas <i>et al.</i> [98]	Yes	Not described	Yes	No	No withdrawal or dropout	3
Ekberg <i>et al.</i> [28]	Yes	Yes	Yes	Yes	No withdrawal or dropout	5
Sakuma <i>et al.</i> [84]	Yes	Not described	No	–	No	1

stabilization splint. In two other studies of somewhat lower reporting quality and validity as compared to the trial by Ekberg *et al.* [28] (see Discussion), no statistically significant difference could be found between the two types of appliances [17, 80].

3. A stabilization appliance does not appear to yield a better clinical outcome than a soft splint [44, 91].
4. There is some evidence from one study with a small number of patients that with a (mandibular) stabilization appliance a statistically significant greater symptom improvement can be achieved than with an anteriorly occluding maxillary splint (“relaxing appliance”) [86].
5. Evidence is missing that treatment with a stabilization appliance leads to a statistically significant greater improvement of signs and symptoms than body acupuncture [47], physical therapy [98] and occlusal adjustment [98].

Discussion

The present investigation has focused on the effect of hard acrylic stabilization splints on pain located in the masticatory muscles. Our review is the first one that has looked specifically into this question. The available data indicate—at different levels of scientific evidence—that a hard acrylic stabilization splint does not yield a better clinical outcome than a non-occluding palatal appliance, a soft splint, or conservative treatment without a splint such as physical therapy and acupuncture. In the only controlled clinical study in which stabilization splints were compared with a partial-coverage anterior splint—a Sved appliance [88]—, there was a trend for greater symptom improvement in the stabilization splint group. This finding should be interpreted with caution, however, because the sample size was small [86]. In one study [98], occlusal

adjustment was carried out. In a recently published systematic review, Koh and Robinson [51] came to the conclusion that there is an absence of evidence that this invasive procedure is an effective therapeutic measure. Therefore, the systematic selective adjustment of the occlusal surface of teeth is not recommended [31, 51, 93].

Within the hierarchy of scientific evidence, systematic reviews are considered to have the highest quality level. These types of publication are carried out to answer one or more focused clinical questions about a topic related to health care [50]. In systematic reviews, it is not always possible to quantitatively combine the data from the identified studies (meta-analysis) [24]. Among other reasons, this may be due to the lack of reported original data or to methodological differences applied in the trials. In that case, a qualitative rather than a quantitative systematic review is carried out [49].

Kalso *et al.* [49] have noted that the strength of evidence lies in the quality of controlled trials: “Systematic reviews can only be as convincing as the quality of the controlled trials allows.” The 13 pertinent publications to answer our clinical question represent nine (mostly short-term) RCTs of varying reporting quality and validity. We have included the meeting abstracts of the RCT by Huggins/Truelove *et al.* [44, 91] and Sakuma *et al.* [84], although these reports have not yet been substantiated by more detailed articles in peer-reviewed journals. Of the 13 publications, three [17, 47, 80] were considered in the qualitative systematic reviews on occlusal splints published by Forssell *et al.* in 1999 [31], Kreiner *et al.* in 2001 [52], and Al-Ani *et al.* in 2004 [2], respectively.

We had to exclude a great number of articles. The reasons for exclusion were:

1. Study participants were not randomized (e.g., [8, 33, 36, 74]). Randomization is the most important proce-

- ture to avoid selection bias [48]. In general, non-randomized studies overestimate treatment effects [9].
2. Patients with masticatory muscle pain *and* TMJ pain were included in a trial / no clear distinction was made between muscle pain and TMJ pain (e.g., [8, 15, 16, 18, 20, 32, 35, 40, 41, 55, 56, 57, 60, 70, 73, 74, 77, 92, 94, 101]). (Nonetheless, inclusion of the results gained in these studies would *not* have altered the conclusions of the present review.)
 3. The effect of stabilization splints was evaluated in patients with TMJ pain (e.g., [25, 26, 27]) or TMJ disk displacement (e.g., [54, 59]).
 4. The effect of splint therapy on the accuracy of mandibular movements of patients with masticatory muscle pain was studied (e.g., [64]).
 5. The effect of soft occlusal splints on masticatory muscle pain was investigated [102].
 6. Multiple simultaneous treatments were carried out (e.g., [14, 37, 39, 69, 78]).
 7. No control group was used (e.g., [4, 8, 36, 100]). Without controls, it cannot be excluded that unspecific effects, e.g., spontaneous remission, natural course of the symptoms, regression to the mean, the placebo effect, the Hawthorne effect, biased (favorable) patient's answers, and accompanying therapeutic measures [95], may have been responsible for observed differences among groups.

When appraising articles on clinical studies in a qualitative systematic review, vote counting (determination of the number of articles showing that an intervention works or does not work) should be avoided [49]. Instead, more weight should be given to publications of (a) high reporting quality (i.e., Jadad score 3 to 5) and (b) high validity [49].

In our review, reporting quality of RCTs was assessed with the help of the Jadad score (range: 0–5). Among the many checklists, scales, and indexes that have been suggested for the evaluation of the (reporting) quality of randomized trials, the Jadad score was developed using standard scale development techniques [46]. Since it is the only known validated scale [6], it has been widely used by clinical researchers. Nonetheless, some critical voices have been raised lately alluding to the fact that the Jadad scale gives more weight to the quality of reporting than to actual methodological quality [49]. In addition, recent reports have pointed out that the inter-rater reliability of the Jadad score may be low [6, 12]. We have striven to avoid this latter problem by independent assessments of the identified articles and discussion between the two reviewers in case of inconsistencies of the scoring results. As Kalso et al. [49] point out, writing a qualitative systematic review requires at least two authors—it “is not a lonely (wo)man’s affair.”

Six of the nine studies considered had a Jadad scale score of 3 or more. In contrast, three studies had an unacceptably low reporting quality of 1 or 2. It has been shown that studies with a Jadad score of 2 or less tend to

give an overoptimistic picture of the real treatment effect than studies with a higher score (e.g., [29]).

There are indications that the reporting quality of a published article does not always correlate with the actual methodological quality of the trial [45]. Deficits in reporting about the results of RCTs have been mentioned by authors in dentistry [65, 87] as well as in medicine [1, 6, 19, 34, 45, 75]. For example, only 25.4% of the articles of RCTs published in the journal *Intensive Care Medicine* up to the year 2000 had a Jadad score of 3 or more [53]. In our review, there is reason to believe that the methodological quality of the studies by Huggins/Truelove [44, 91] and Sakuma et al. [84], which were awarded 1 point, respectively, is much better than the actual Jadad score suggests.

As far as the assessment of the second quality factor—high validity—is concerned, two important criteria are a sufficient baseline pain intensity [66] and an adequate number of patients in each group [67]. In the identified trials, recruiting of study participants took place by either resorting to patients seeking care at or being referred to an orofacial pain care center, or by placing announcements in local print media [17, 80]. It should be taken into consideration, however, that patients seeking TMD treatment by referrals are probably different from individuals recruited by a notice in a local newspaper or journal [28]: Rubinoff et al. [80] argued in the critical discussion of their study published in 1987 that the latter patients “may have been biased toward milder conditions that were tolerable to the patient until prompted by a media notice.” This assumption appears to be correct. When the baseline pain intensities of the three studies published in peer-reviewed journals in which a stabilization appliance were compared with a non-occluding appliance are analyzed, the following can be observed:

- a. In the trial by Rubinoff et al. [80], the majority of the study participants had low pain before the start of the study. All patients were recruited by a newspaper notice.
- b. In the investigation by Dao et al. [17], pre-treatment pain intensities were about 40 mm, which is equivalent to moderate pain [13], on a visual analog scale (VAS). Part of the participants were recruited through announcements published in local journals; the other part were referred by dentists.
- c. In the study by Ekberg et al. [28], the pain intensity prior to the start of the study was between moderate and very severe. All patients were referred for and requested treatment.

Hence, differences in the recruitment of patients may be one reason for different pre-treatment pain levels and the different results achieved in the three studies. The three trials also differ considerably with regard to the included number of patients (cf. Table 2). The result from this comparison is that two studies with an acceptable reporting quality and a moderate validity [17, 80] found no difference between a stabilization appliance and a non-

occluding appliance. These results differ from the recent study by Ekberg et al. [28] with both strong evidence and a strong validity supporting the efficacy of the Michigan splint compared with a palatal appliance.

Only two trials—those conducted by the Huggins/Truelove group [44, 91] and by van der Glas et al. [98], respectively—had an observation period of at least 1 year. In three studies [28, 47, 86], Helkimo's CDI [38] was used. However, the validity of this index has been shown to be doubtful [99].

Our search has also demonstrated the inconsistencies that exist among different authors with regard to the diagnosis "muscle pain" (Table 2). This variability is reflected in the number of diagnostic systems that have been proposed over the past decades for classifying the different subsets of TMDs (c.f. [71]). It was not before 1992 that biologically plausible classifications with specific diagnostic categories became available [22, 89]. As a result, comparisons among the studies identified in this search are difficult to make, and pooling of data is impossible. As far as the evaluation of trials carried out in the 1970s and 1980s is concerned (c.f. [8, 36, 80]), study participants diagnosed with "myofascial pain dysfunction syndrome" are not necessarily patients suffering from myalgia only. Although masticatory muscle pain is a symptom encountered in most, albeit not all of these patients, symptoms such as limited mandibular opening, TMJ sounds, and deviation on jaw opening may be present. Only one of these older articles allowed an explicit assessment of the symptom "muscular pain" [8].

Furthermore, our investigation questions the strategy of relying exclusively on Medline when looking for evidence. Of the 13 pertinent articles, seven were identified by the Ovid Medline search. Conversely, the meeting abstracts found by handsearching were not listed in any of the consulted electronic databases. Hence, as our search demonstrated, important study results may be missed if one relies on Medline as the sole information source [1, 21, 97]. Besides, limitation of the search to the English language, as it is often seen in reviews, may lead to different conclusions (English language bias) [23, 63].

Another point that can be made is that keywords are likely to change depending on the prevailing thinking in the field. For example, whereas "temporomandibular disorders" is a term that was agreed upon by most clinicians in the early 1990s [62], Medline lists articles about this topic under the medical subject heading "Temporomandibular Joint Dysfunction Syndrome."

Conclusions

The dearth of adequate clinical studies that are available to answer our clinical question mirrors the lack of hard data in an important and frequent scenario in clinical dentistry. We found this result astonishing because full-coverage heat-cured (or self-curing) acrylic resin occlusal splints have been in use since the 1960s [76], and these devices have been recommended by many clinicians

around the world for the management of patients with TMDs, including masticatory muscle pain [68]. Generally, the wealth of anecdotal reports and uncontrolled clinical observations tend to give a much more optimistic picture about the presumed effects of the stabilization splint therapy.

Due to the limited number of available studies, our clinical question can only be answered tentatively: based on the currently best available evidence it appears that most patients with masticatory muscle pain are helped by the incorporation of a stabilization splint. Nevertheless, evidence is equivocal that improvement of pain symptoms after incorporation of an intraoral appliance is caused by a specific effect of the splint [17, 28, 80]. In addition, there is not enough data about the long-term efficacy and effectiveness of these widely used therapeutic tools. It should be noted that a scarcity of prospective randomized controlled trials with high power does not discredit the concept or the applicability of EBM, because EBM is based on the *best available* evidence. However, by pointing out deficits in the quantity and quality of the evidence, EBM highlights the empirical nature of current management and emphasizes the need for more focused clinical research in dentistry.

Acknowledgements The authors are grateful to Dr. Hiyoyasu Komizumi (Nihon University, Tokyo, Japan) for the search in the Japanese database.

References

1. Adetugbo K, Williams H (2000) How well are randomized controlled trials reported in the dermatology literature? *Arch Dermatol* 136:381–385
2. Al-Ani MZ, Davies SJ, Gray RJM, Sloan P, Glenny AM (2004) Stabilization splint therapy for temporomandibular pain dysfunction syndrome (Cochrane review). The Cochrane Library, issue 1. John Wiley & Sons, Chichester, UK
3. Arbree NS, Campbell SD, Renner RP, Goldstein GR (1995) A survey of temporomandibular disorder conducted by the Greater New York Academy of Prosthodontics. *J Prosthet Dent* 74:512–516
4. Baldissara S, Mascellani SC, Catapano S, Baldissara P (1998) Effetti a breve termine della placca di Michigan sul dolore muscolare ed articolare. *Minerva Stomatol* 47:235–238
5. Bell WE (1989) Orofacial pains. Classification, diagnosis, management, 4th edn. Year Book Medical Publishers, Chicago
6. Bhandari M, Richards RR, Sprague S, Schemitsch EH (2001) Quality in the reporting of randomized trials in surgery: is the Jadad scale reliable? *Control Clin Trials* 22:687–688
7. Cane L, Schieroni MP, Ribero G, Ferrero M, Carossa S (1997) Effectiveness of the Michigan splint in reducing functional cervical disturbances: a preliminary study. *J Craniomand Pract* 15:132–135
8. Carraro JJ, Caffesse RG (1978) Effect of occlusal splints on TMJ symptomatology. *J Prosthet Dent* 40:563–566
9. Carroll D, Tramèr M, McQuay H, Nye B, Moore A (1996) Randomization is important in studies with pain outcomes: systematic review of transcutaneous electrical nerve stimulation in acute postoperative pain. *Br J Anaesth* 77:798–803
10. Clark GT (1984) A critical evaluation of orthopedic interocclusal appliance therapy: effectiveness for specific symptoms. *J Am Dent Assoc* 108:364–368

11. Clark GT (1984) A critical evaluation of orthopedic interocclusal appliance therapy: design, theory, and overall effectiveness. *J Am Dent Assoc* 108:359–364
12. Clark HD, Wells GA, Huet C, McAlister FA, Salmi LR, Fergusson D, Laupacis A (1999) Assessing the quality of randomized trials: reliability of the Jadad scale. *Control Clin Trials* 20:448–452
13. Collins SL, Moore RA, McQuay HJ (1997) The visual analogue pain intensity scale: what is moderate pain in millimetres? *Pain* 72:95–97
14. Crockett DJ, Foreman ME, Alden L, Blasberg B (1986) A comparison of treatment modes in the management of myofascial pain dysfunction syndrome. *Biofeedback Self Regul* 11:279–291
15. Dahlström L, Haraldson T (1985) Bite plates and stabilization splints in mandibular dysfunction. A clinical and electromyographic comparison. *Acta Odontol Scand* 43:109–114
16. Dahlström L, Carlsson GE, Carlsson SG (1982) Comparison of effects of electromyographic biofeedback and occlusal splint therapy on mandibular dysfunction. *Scand J Dent Res* 90:151–156
17. Dao TTT, Lavigne GJ, Charbonneau A, Feine JS, Lund JP (1994) The efficacy of oral splints in the treatment of myofascial pain of the jaw muscles: a controlled clinical trial. *Pain* 56:85–94
18. Davies SJ, Gray RJ (1997) The pattern of splint usage in the management of two common temporomandibular disorders. Part II: The stabilisation splint in the treatment of pain dysfunction syndrome. *Br Dent J* 183:247–251
19. Devereaux PJ, Manns BJ, Ghali WA, Quan H, Guyatt GH (2002) The reporting of methodological factors in randomized controlled trials and the association with a journal policy to promote adherence to the Consolidated Standards of Reporting Trials (CONSORT) checklist. *Control Clin Trials* 23:380–388
20. di Paolo C, Panti F, Rampello A (1998) Studio retrospettivo sull'applicazione di splint occlusali di 312 pazienti affetti da disordini temporo-mandibolari. *Minerva Stomatol* 47:187–195
21. Dickersin K, Scherer R, Lefebvre C (1994) Identifying relevant studies for systematic reviews. *Br Med J* 309:1286–1291
22. Dworkin SF, LeResche L (1992) Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord Facial Oral Pain* 6:301–355
23. Egger M, Zellweger-Zähner T, Schneider M, Junker C, Lengeler C, Antes G (1997) Language bias in randomised controlled trials published in English and German. *Lancet* 350:326–329
24. Egger M, Smith GD, Altman DG (2001) Systematic reviews in health care: meta-analysis in context, 2nd edn. BMJ Books, London
25. Ekberg E (1998) Treatment of temporomandibular disorders of arthrogenous origin. Controlled double-blind studies of a non-steroidal anti-inflammatory drug and a stabilisation appliance. *Swed Dent J Suppl* 131:1–57
26. Ekberg E, Nilner M (1999) The influence of stabilisation appliance therapy and other factors on the treatment outcome in patients with temporomandibular disorders of arthrogenous origin. *Swed Dent J* 23:39–47
27. Ekberg EC, Sabet ME, Petersson A, Nilner M (1998) Occlusal appliance therapy in a short-term perspective in patients with temporomandibular disorders correlated to condyle position. *Int J Prosthodont* 11:263–268
28. Ekberg EC, Vallon D, Nilner M (2003) The efficacy of appliance therapy in patients with temporomandibular disorders of mainly myogenous origin. A randomized, controlled, short-term study. *J Orofac Pain* 17:133–139
29. Ezzo J, Berman B, Hadhazy VA, Jadad AR, Lao L, Singh BB (2000) Is acupuncture effective for the treatment of chronic pain? A systematic review. *Pain* 86:217–225
30. Forssell H, Kalso E (2004) Application of principles of evidence-based medicine to occlusal treatment for temporomandibular disorders: are there lessons to be learned? *J Orofac Pain* 18:9–22
31. Forssell H, Kalso E, Koskela P, Vehmanen R, Puukka P, Alanen P (1999) Occlusal treatments in temporomandibular disorders: a qualitative systematic review of randomized controlled trials. *Pain* 83:549–560
32. Frahn G, John M (1996) Schmerzen im orofazialen System—eine kontrollierte Studie mit Stabilisierungsschiene und Ultraschall. *Dtsch Zahnärztl Z* 51:478–481
33. Gavish A, Winocur E, Ventura YS, Halachmi M, Gazit E (2002) Effect of stabilization splint therapy on pain during chewing in patients suffering from myofascial pain. *J Oral Rehabil* 29:1181–1186
34. Gluud C, Nikolova D (1998) Quality assessment of reports on clinical trials in the Journal of Hepatology. *J Hepatol* 29:321–327
35. Gray RJ, Davies SJ, Quayle AA, Wastell DG (1991) A comparison of two splints in the treatment of TMJ pain dysfunction syndrome. Can occlusal analysis be used to predict success of splint therapy? *Br Dent J* 170:55–58
36. Greene CS, Laskin DM (1972) Splint therapy for the myofascial pain-dysfunction (MPD) syndrome: a comparative study. *J Am Dent Assoc* 84:624–628
37. Greene CS, Laskin DM (1983) Long-term evaluation of treatment for myofascial pain-dysfunction syndrome: a comparative analysis. *J Am Dent Assoc* 107:235–238
38. Helkimo M (1974) Studies on function and dysfunction of the masticatory system. II. Index for anamnestic and clinical dysfunction and occlusal state. *Swed Dent J* 67:101–121
39. Helöe B, Heiberg AN (1980) A follow-up study of a group of female patients with myofascial pain-dysfunction syndrome. *Acta Odontol Scand* 38:129–134
40. Heuser MKH (1989) Vergleichende Untersuchung über die Behandlung des myofazialen Schmerzsyndroms mit Aufbißschienen, Neuraltherapie nach Huneke und Akupunktur. *Quintessenz* 40:531–542
41. Hijzen TH, Slangen JL, van Houweligen HC (1986) Subjective, clinical and EMG effects of biofeedback and splint treatment. *J Oral Rehabil* 13:529–539
42. Huggins KH, Truelove EL, Dworkin SF, LeResche L, Sommers E, Schubert M (1997) Randomized clinical trial (RCT) of a soft splint for TMD (Abstract 3005). *J Dent Res (spec issue)* 76:389
43. Huggins KH, Truelove EL, Dworkin SF, Mancil L, Sommers E, LeResche L (1998) Initial outcomes in RCT of splint for TMD: clinical findings (Abstract 152). *J Dent Res (spec issue A)* 77:112
44. Huggins KH, Truelove EL, Dworkin SF, Mancil L, Sommers E, LeResche L (1999) RCT for splints in TMD: clinical findings at 12 months (Abstract 1490). *J Dent Res (spec issue)* 78:292
45. Huwiler-Müntener K, Jüni P, Junker C, Egger M (2002) Quality of reporting of randomized trials as a measure of methodologic quality. *J Am Med Assoc* 287:2801–2804
46. Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJM, Gavaghan DJ, McQuay HJ (1996) Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Controlled Clin Trials* 17:1–12
47. Johansson A, Wenneberg B, Wagersten C, Haraldson T (1991) Acupuncture in treatment of facial muscular pain. *Acta Odontol Scand* 49:153–158
48. Jüni P, Altman DG, Egger M (2001) Assessing the quality of randomised controlled trials. In: Egger M, Smith GD, Altman DG (eds) *Systematic reviews in health care: meta-analysis in context*, 2nd edn. BMJ Books, London, pp 87–108
49. Kalso E, Edwards J, McQuay HJ, Moore RA (2002) Five easy pieces on evidence-based medicine (4). *Eur J Pain* 6:89–93
50. Khan KS, Kunz R, Kleijnen J, Antes G (2003) Systematic reviews to support evidence-based medicine. Royal Society of Medicine Press, London
51. Koh H, Robinson PG (2003) Occlusal adjustment for treating and preventing temporomandibular joint disorders (Cochrane

- Review). The Cochrane Library, issue 1. John Wiley & Sons, Chichester, UK
52. Kreiner M, Betancor E, Clark GT (2001) Occlusal stabilization appliances. Evidence of their efficacy. *J Am Dent Assoc* 132:770–777
 53. Latronico N, Botteri M, Minelli C, Zanotti C, Bertolini G, Candiani A (2002) Quality of reporting of randomised controlled trials in the intensive care literature. A systematic analysis of papers published in *Intensive Care Medicine* over 26 years. *Intensive Care Med* 28:1316–1323
 54. Linde C, Isacson G, Jonsson BG (1995) Outcome of 6-week treatment with transcutaneous electric nerve stimulation compared with splint on symptomatic temporomandibular joint disk displacement without reduction. *Acta Odontol Scand* 53:92–98
 55. List T, Helkimo M (1992) Acupuncture and occlusal splint therapy in the treatment of craniomandibular disorders. II. A 1-year follow-up study. *Acta Odontol Scand* 50:375–385
 56. List T, Helkimo M, Andersson S, Carlsson GE (1992) Acupuncture and occlusal splint therapy in the treatment of craniomandibular disorders. Part I. A comparative study. *Swed Dent J* 16:125–141
 57. List T, Helkimo M, Karlsson R (1993) Pressure pain thresholds in patients with craniomandibular disorders before and after treatment with acupuncture and occlusal splint therapy: a controlled clinical study. *J Orofac Pain* 7:275–282
 58. List T, Dworkin SF (1996) Comparing TMD diagnoses and clinical findings at Swedish and US TMD centers using research diagnostic criteria for temporomandibular disorders. *J Orofac Pain* 10:240–253
 59. Lundh H, Westesson PL, Eriksson L, Brooks SL (1992) Temporomandibular joint disk displacement without reduction. Treatment with flat occlusal splint versus no treatment. *Oral Surg Oral Med Oral Pathol* 73:655–658
 60. Manns A, Miralles R, Santander H, Valdivia J (1983) Influence of the vertical dimension in the treatment of myofascial pain-dysfunction syndrome. *J Prosthet Dent* 50:700–709
 61. Marbach JJ, Raphael KG (1996) Treatment of orofacial pain using evidence-based medicine: the case for intraoral appliances. In: Campbell JN (ed) *Pain 1996—an updated review*. IASP Press, Seattle, pp 413–422
 62. McNeill C (1992) Panel discussion: definition of terms. In: McNeill C (ed) *Current controversies in temporomandibular disorders*. Quintessence, Chicago, pp 15–17
 63. Moher D, Fortin P, Jadad AR, Jüni P, Klassen T, Le Lorier J, Liberati A, Linde K, Penna A (1996) Completeness of reporting of trials published in languages other than English: implications for conduct and reporting of systematic reviews. *Lancet* 347:363–366
 64. Monteiro AA, Clark GT (1988) Mandibular movement feedback vs. occlusal appliances in the treatment of masticatory muscle dysfunction. *J Craniomandib Disord* 2:41–47
 65. Montenegro R, Needleman I, Moles D, Tonetti M (2002) Quality of RCTs in periodontology—a systematic review. *J Dent Res* 81:866–870
 66. Moore A, Collins S, Carroll D, McQuay H (1997) Paracetamol with and without codeine in acute pain: a quantitative systematic review. *Pain* 70:193–201
 67. Moore RA, Gavaghan D, Tramer MR, Collins SL, McQuay HJ (1998) Size is everything—large amounts of information are needed to overcome random effects in estimating direction and magnitude of treatment effects. *Pain* 78:209–216
 68. National Institutes of Health (1996) National Institutes of Health Technology Assessment Conference Statement: management of temporomandibular disorders. *J Am Dent Assoc* 127:1595–1603
 69. Nel H (1978) Myofascial pain-dysfunction syndrome. *J Prosthet Dent* 40:438–441
 70. Nemcovsky CE, Gazit E, Serfati V, Gross M (1992) A comparative study of three therapeutic modalities in a temporomandibular disorder (TMD) population. *J Craniomand Pract* 10:148–155
 71. Ohrbach R, Stohler CS (1992) Review of the literature. A: Current diagnostic systems. *J Craniomand Disord Facial Oral Pain* 6:307–317
 72. Okeson JP (ed) (1996) *Orofacial pain. Guidelines for assessment, diagnosis, and management*, 3rd edn. Quintessence, Chicago
 73. Okeson JP, Moody PM, Kemper JT, Haley JV (1983) Evaluation of occlusal splint therapy and relaxation procedures in patients with temporomandibular disorders. *J Am Dent Assoc* 107:420–424
 74. Pettengill CA, Growney MR, Schoff R, Kenworthy CR (1998) A pilot study comparing the efficacy of hard and soft stabilizing appliances in treating patients with temporomandibular disorders. *J Prosthet Dent* 79:165–168
 75. Pienaar ED, Volmink J, Zwarenstein M, Swinger GH (2002) Randomised trials in the South African Medical Journal, 1948–1997. *S Afr Med J* 92:901–903
 76. Ramfjord SP, Ash MM (1994) Reflections on the Michigan occlusal splint. *J Oral Rehabil* 21:491–500
 77. Raphael KG, Marbach JJ (2001) Widespread pain and the effectiveness of oral splints in myofascial face pain. *J Am Dent Assoc* 132:305–316
 78. Rizzatti-Barbosa CM, Martinelli DA, Ambrosano GM, de Albergaria-Barbosa JR (2003) Therapeutic response of benzodiazepine, orphenadrine citrate and occlusal splint association in TMD pain. *J Craniomand Pract* 21:116–120
 79. Roldan OV, Maglione H, Carreira R, Mainieri S (1990) Piroxicam, diazepam y placebo en el tratamiento de las disfunciones de la articulación temporomandibular. Estudio doble ciego. *Rev Asoc Odontol Argent* 78:83–85
 80. Rubino MS, Gross A, McCall WD Jr (1987) Conventional and nonoccluding splint therapy compared for patients with myofascial pain dysfunction syndrome. *Gen Dent* 35:502–506
 81. Rudy T, Greco C, Turk D, Zaki H, Herlich A (1996) Empirically-derived RDC axis I classification of TMD patients (Abstract 1599). *J Dent Res* (spec issue) 75:217
 82. Sackett DL, Richardson WS, Rosenberg W, Haynes RB (1997) *Evidence-based medicine. How to practice and teach EBM*. Churchill Livingstone, New York
 83. Sackett DL, Straus SE, Richardson WS, Rosenberg W, Haynes RB (2000) *Evidence-based medicine. How to practice and teach EBM*, 2nd edn. Churchill Livingstone, Edinburgh
 84. Sakuma S, Aiji Y, Nakagawa M, Katsuse S, Suzuki J, Katsuno R, Nakamura H, Ko K, Takeuchi Y, Inugai Y, Noda K, Izumi M, Ogi M, Kurita K, Aiji E, Ito Y (2003) Therapy effect of splints for temporomandibular disorders RDC type I [in Japanese] (Abstract 2C4). *J Jap Soc Temporomandib Joint* 15:118
 85. Schulz KF, Chalmers I, Hayes RJ, Altman DG (1995) Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *J Am Med Assoc* 273:408–412
 86. Siegert R, Gundlach KKH (1989) Stabilisationsschiene versus Entspannungsbehelf zur Behandlung myofazialer Schmerzen. Erste Ergebnisse einer prospektiv randomisierten Studie. *Dtsch Zahnärztl Z* 44:S17–S19
 87. Sjögren P, Halling A (2002) Quality of reporting randomised clinical trials in dental and medical research. *Br Dent J* 192:100–103
 88. Sved A (1944) Changing the occlusal level and a new method of retention. *Am J Orthod* 30:527–535
 89. Truelove EL, Sommers EE, LeResche L, Dworkin SF, Von Korff M (1992) Clinical diagnostic criteria for TMD. New classification permits multiple diagnoses. *J Am Dent Assoc* 123:47–54
 90. Truelove EL, Huggins KH, Dworkin SF, Mancl L, Sommers E, LeResche L (1998) RCT splint treatment outcome in TMD: initial self report findings (Abstract 151). *J Dent Res* (spec issue A) 77:112
 91. Truelove EL, Huggins KH, Dworkin SF, Mancl L, Sommers E, LeResche L (1999) RCT for splint treatment in TMD:

- 12-month self-report outcomes (Abstract 1491). *J Dent Res* (spec issue) 78:292
92. Tsuga K, Akagawa Y, Sakaguchi R, Tsuru H (1989) A short-term evaluation of the effectiveness of stabilization-type occlusal splint therapy for specific symptoms of temporomandibular joint dysfunction syndrome. *J Prosthet Dent* 61:610–613
 93. Tsukiyama Y, Baba K, Clark GT (2001) An evidence-based assessment of occlusal adjustment as a treatment for temporomandibular disorders. *J Prosthet Dent* 86:57–66
 94. Turk DC, Zaki HS, Rudy TE (1993) Effects of intraoral appliance and biofeedback/stress management alone and in combination in treating pain and depression in patients with temporomandibular disorders. *J Prosthet Dent* 70:158–164
 95. Türp JC, Schwarzer G (2003) Zur Wirksamkeit therapeutischer Massnahmen: Der Post-hoc-ergo-propter-hoc-Trugschluss. *Schweiz Monatsschr Zahnmed* 113:36–46
 96. Türp JC, Kowalski CJ, Stohler CS (1998) Treatment-seeking patterns of facial pain patients: many possibilities, limited satisfaction. *J Orofac Pain* 12:61–66
 97. Türp JC, Schulte JM, Antes G (2002) Nearly half of dental randomized controlled trials published in German are not included in Medline. *Eur J Oral Sci* 110:405–411
 98. van der Glas HW, Buchner R, van Grootel RJ (2000) Vergelijking tussen behandelingsvormen bij myogene temporomandibulaire dysfunctie. *Ned Tijdschr Tandheelkd* 107: 505–512
 99. van der Weele LT, Dibbets JM (1987) Helkimo's index: a scale or just a set of symptoms? *J Oral Rehabil* 14:229–237
 100. Vauthier F, Spirgi M (1977) La gouttière occlusale et son influence sur l'enregistrement frontal du mouvement d'ouverture maximale et de fermeture dans les cas de troubles algodysfonctionnels des articulations temporo-mandibulaires. *Schweiz Monatsschr Zahnmed* 87:189–208
 101. Wahlund K, List T, Larsson B (2003) Treatment of temporomandibular disorders among adolescents: a comparison between occlusal appliance, relaxation training, and brief information. *Acta Odontol Scand* 61:203–211
 102. Wright E, Anderson G, Schulte J (1995) A randomized clinical trial of intraoral soft splints and palliative treatment for masticatory muscle pain. *J Orofac Pain* 9:192–199

Copyright of Clinical Oral Investigations is the property of Kluwer Academic Publishing / Academic and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.