

Periodontal inflamed surface area and C-reactive protein as predictors of HbA1c: a study in Indonesia

Hendri Susanto · Willem Nesse · Pieter U. Dijkstra ·
Evelien Hoedemaker · Yvonne Huijser van Reenen ·
Dewi Agustina · Arjan Vissink · Frank Abbas

Received: 2 July 2011 / Accepted: 2 October 2011 / Published online: 20 October 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract Periodontitis may exert an infectious and inflammatory burden, evidenced by increased C-reactive protein (CRP). This burden may impair blood glucose control (HbA1c). The aim of our study was to analyze whether periodontitis severity as measured with the periodontal inflamed surface area (PISA) and CRP predict HbA1c levels in a group of healthy Indonesians and a group of Indonesians treated for type 2 diabetes mellitus (DM2). A full-mouth periodontal examination, including probing pocket depth, gingival recession, clinical attachment loss, plaque index and bleeding on probing, was performed in 132 healthy Indonesians and 101 Indonesians treated for DM2. Using these data, PISA was calculated. In addition, HbA1c and CRP were analyzed. A validated questionnaire

was used to assess smoking, body mass index (BMI), education and medical conditions. In regression analyses, it was assessed whether periodontitis severity and CRP predict HbA1c, controlling for confounding and effect modification (i.e., age, sex, BMI, pack years, and education). In healthy Indonesians, PISA and CRP predicted HbA1c as did age, sex, and smoking. In Indonesians treated for DM2, PISA did not predict HbA1c. Periodontitis may impair blood glucose regulation in healthy Indonesians in conjunction with elevated CRP levels. The potential effect of periodontitis on glucose control in DM2 patients may be masked by DM2 treatment. Clinical relevance: periodontitis may impair blood glucose control through exerting an inflammatory and infectious burden evidenced by increased levels of CRP.

Hendri Susanto and Willem Nesse equally contributed to this study.

H. Susanto · D. Agustina
Department of Oral Medicine, Faculty of Dentistry,
Gadjah Mada University,
Denta Sekip Utara Yogyakarta 55281, Indonesia

W. Nesse · A. Vissink (✉)
Department of Oral and Maxillofacial Surgery, University
Medical Center Groningen, University of Groningen,
P.O. Box 30.001, 9700 RB Groningen, the Netherlands
e-mail: a.vissink@umcg.nl

P. U. Dijkstra
Center for Rehabilitation and Department of Oral
and Maxillofacial Surgery, University Medical Center Groningen,
University of Groningen,
P.O. Box 30.001, 9700 RB Groningen, the Netherlands

E. Hoedemaker · Y. H. van Reenen · F. Abbas
Center for Dentistry and Oral Hygiene, Department of
Periodontology,
University Medical Center Groningen, University of Groningen,
Ant. Deusinglaan 1,
9713 AV Groningen, the Netherlands

Keywords Periodontitis · Periodontal inflamed surface area/PISA · HbA1c · Type 2 diabetes mellitus

Introduction

Periodontal inflamed surface area (PISA) quantifies the amount of inflamed periodontal tissue and is supposed to quantify the inflammatory and infectious burden resulting from periodontitis [1]. Periodontitis may cause an inflammatory burden through the production of local inflammatory mediators entering the systemic circulation. This inflammatory burden is evidenced by increased serum C-reactive protein (CRP) levels found in patients suffering from periodontitis [2–7].

Periodontitis may also cause an infectious burden through bacteria and their products entering the systemic circulation. This burden may endanger overall health by for example causing atherosclerosis [8] and cardiovascular diseases [9].

Circulating oral bacteria and lipopolysaccharides are also able to stimulate hepatocytes to secrete CRP [10–12]. Thus, increased levels of CRP associated with periodontitis may be considered as a common pathway for both the inflammatory and infectious burden resulting from periodontitis.

The high levels of CRP accompanying periodontitis, may lead to insulin resistance and thereby to poor control of blood glucose in type 2 diabetes mellitus (DM2) and healthy subjects [13–16] resulting in increased levels of glycosylated hemoglobin (HbA1c) [17, 18]. Accordingly, high levels of CRP are correlated with increased HbA1c levels in DM2 patients [19, 20]. Increased CRP levels have also been associated with an increased risk of cardiovascular disease (CVD) in DM2 patients, the major cause of the increased mortality in DM2 patients [21, 22].

Given the importance of blood glucose control in preventing CVD in DM2 patients, relatively few studies assessed whether the specific combination of periodontitis and CRP as a reflection of inflammatory burden, predict HbA1c levels [23, 24]. Furthermore, no study has investigated this issue in Indonesia while this country is in the top ten of countries with the highest prevalence of DM (at 5.7%) in the world in 2005. It has been estimated that by the year 2030, the prevalence of DM in Indonesia will be about 10%, corresponding with 20 million DM patients [25, 26]. In addition to a high DM prevalence, a high prevalence of periodontitis has been reported in Indonesia [27]. Finally, depending on the definition used, our previous study showed that Indonesian DM2 patients have a prevalence of periodontitis at 72% to 93% and an increased severity of periodontitis [28]. Therefore, the aim of this study was to assess whether periodontitis severity, as measured with the PISA method and CRP predict HbA1c levels in healthy Indonesians and Indonesian treated for DM2 patients.

Materials and methods

The participants in this study were recruited from three different sites, namely (1) the Internal Medicine Department Dr. Sardjito Hospital, Yogyakarta, 2) Prof. Soedomo Dental Hospital, Faculty of Dentistry, Gadjah Mada University, Yogyakarta, and 3) Diabetes Center of Jogjakarta International Hospital, Yogyakarta, Indonesia. All participants had to be aged ≥ 18 years and had to have ≥ 8 remaining teeth. The latter inclusion criteria was proposed prior to commencing the study since current inflammatory burden posed by periodontitis requires the presence of at least a minimum number of teeth affected by periodontitis. This study was approved by the Ethical Committee for Research of the Medical Faculty of Gadjah Mada University, Yogyakarta, Indonesia.

All participants completed a validated general health assessment questionnaire [29, 30] to check for other medical conditions that might be a risk factor for periodontitis. Information on age, gender, height and weight (for body mass index (BMI) calculation), education, and smoking were obtained from each participant.

All participants underwent a periodontal examination including periodontal probing pocket depth (PD), gingival recession, plaque score, and bleeding on probing (BOP) measurements by trained and calibrated examiners (HS, YHR, and EH). All measurements were performed on all teeth, on six sites per tooth using a manual periodontal color-coded standard probe (Dentsply™, London, UK).

Clinical attachment loss (AL) was defined as the distance from the cemento-enamel junction to the bottom of the pocket/sulcus and calculated as the mathematical sum of the PD and gingival recession measurements [31]. Measurements were made in millimeters and were rounded off to the nearest millimeter. BOP was recorded as either present or absent within 30 s of probing at six sites per tooth. Plaque score was defined as being present or absent at six points on each tooth [32]. The number of missing teeth was also recorded.

Periodontitis extent and severity were operationalized using a variety of methods, all of which are currently used in literature studying the association between periodontitis and other diseases. All measurements were calculated using conventional clinical measurements obtained during the full-mouth periodontal examination (HS and WN). The number of sites with probing PD of ≥ 4 , ≥ 5 , and ≥ 6 mm, the numbers of sites with clinical AL of ≥ 3 , ≥ 4 , ≥ 5 , and ≥ 6 mm, mean PD, mean AL [33], and the percentage of sites with BOP were calculated. Additionally, a recently introduced measure of periodontitis severity, the PISA [1] was calculated, PISA quantifies the amount of inflamed periodontal tissue, and it is suggested that PISA thereby quantifies the inflammatory burden exerted by periodontitis [1].

Finally, all participants underwent a venapuncture to obtain a blood sample. Both blood glucose (fasting blood glucose), determined by glucose oxidase enzymatic method, and glycosylated/glycosylated hemoglobin (HbA1c) values, determined using low pressure cation ion exchange chromatography (DIASTAT™, Bio-Rad, USA), were determined for all participants. Additionally, CRP, determined by a high-sensitivity chemiluminescent immunometric assay (Immuline 2000™, Diagnostic Products Corp., Los Angeles, CA, USA), were determined for all participants.

Statistical analysis

Differences in periodontitis severity between DM2 and healthy subjects were analyzed first using univariate

analyses (independent sample *t* test, Mann–Whitney *U* test, or Chi-square test as appropriate). Likewise, differences in potential predictors of periodontitis (age, gender, BMI, education, smoking, plaque score, number of teeth, ethnicity, and other medical conditions) between DM2 patients and healthy subjects were tested for significance using univariate analyses. Since periodontitis severity was operationalized in 12 different ways, significance level α of 0.05 was corrected for multiple comparisons according to the Bonferroni–Holm method.

To assess periodontitis and CRP as potential predictors of HbA1c in subjects with and without DM2, a multiple linear regression analysis was performed, using a backward stepwise method. HbA1c was the dependent. As independent variables, i.e., predictors of HbA1c: age, sex, BMI, pack years, education, CRP, and all measures of periodontitis severity. The latter were introduced in the model one by one. Statistics were calculated using SPSS 16.0.

Results

In a total of 101 participants, 40 men and 61 women, with a mean age of 54 years treated for DM2, diagnosed according to World Health Organization criteria [34] were included

(Table 1). Most of the participants were of Javanese origin (93.0%). The mean BMI in this group was 25.5, labeling them as being overweight according to the World Health Organization classification (overweight—BMI, 23–27.5) [35]. The DM2 patients had an average of 24 teeth with a mean plaque score of 91% (Table 2). The mean HbA1c in the DM2 group was 8.9%. In addition, a total of 132 healthy controls was enrolled, 34 men and 98 women, with a mean age of 48 years. Again, the healthy subjects were mainly of Javanese origin (90%) and were on average ranked as being overweight according to the BMI index for Asian populations (BMI, 24.4) [35]. The healthy subjects had an average of 26 teeth and a mean plaque score of 93%. The average HbA1c level of this group was 5.5% (below 6.5% classified as healthy/non diabetic).

The severity of periodontitis was significantly higher in participants with DM2 when compared to controls, again independent of the method used to operationalize periodontitis severity (Table 2). In the DM2 patients, none of the measures of periodontitis severity predicted HbA1c. Only SES ($\beta=-0.752$; 95% CI=-1.345 to -0.158) was a predictor of HbA1c, (r^2 of 6.0%; data not shown). By contrast, in healthy Indonesians, PISA ($\beta=0.0004$; 95% CI=0.00004–0.00076) was a predictor of HbA1c together with age, sex, smoking and CRP, ($r^2=21\%$) in the model (Table 3).

Table 1 Characteristics of the participants

Variables	DM2 (<i>n</i> =101)	Healthy controls (<i>n</i> =132)	<i>p</i> value
Demography			
Age (years) mean (SD)	54.4 (10.7)	47.9 (10.1)	<0.001
Smoking (pack year) median (IQR) ^b	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.109
BMI (kg/m ²) mean (SD)	25.5(4.2)	24.4 (3.8)	0.054
Java origin (<i>n</i> (%)) ^a	94 (93%)	119 (90%)	0.431
Gender (<i>n</i> (%)) ^a			<0.05
Male	40 (40%)	34 (26%)	
Female	60 (60%)	98 (74%)	
Education (<i>n</i> (%)) ^a	2.18 (0.79)	2.13 (0.77)	0.662
Low (6–9 years)	32 (24%)	24 (23%)	
Middle (9–12 years)	51 (39%)	34 (34%)	
High (12–17 years)	49 (37%)	43 (43%)	
Hypertension (yes) (<i>n</i> (%)) ^a	26 (26%)	24 (18%)	0.164
Medication (<i>n</i> (%))			
Sulfonylurea	55 (54%)	NA	NA
Insulin	41 (41%)	NA	
Metformin	37 (37%)	NA	
Acarbose	33 (33%)	NA	
Laboratory serum markers			
HbA1c (%) mean (SD)	8.9 (2.4)	5.5 (0.4)	<0.001
CRP (mg/l) median (IQR) ^b	2.8(1.6–7.0)	1.1 (0.6–2.3)	<0.001

p values were Bonferroni–Holm corrected. A *p* value of <0.05 was considered statistically significant

p probability, *BMI* body mass index, *CRP* C-reactive protein, *DM2* diabetes mellitus type 2, *HbA1c* glycated hemoglobin, *IQR* interquartile range, *n* number of participants, *NA* not applicable, *SD* standard deviation, low, elementary and junior school (education), *middle* high school (education), *high* university (education)

^aResult of Chi-square test

^bResult of Mann–Whitney *U* test; other results are the results of the independent samples *t* test

Table 2 Periodontal status of healthy controls and DM2

Variables	Healthy controls (n=132)	DM2 (n=101)	p value
Periodontal prevalence (n (%)) ^a			
PD (≥4 mm) and AL (≥3 mm)	89 (67%)	89 (88%)	<0.001
PD (≥5 mm) AL (≥2 mm)	43 (33%)	67 (66%)	<0.001
Periodontal severity (median (IQR)) ^b			
PISA (mm ²)	83.9 (35.2–206.4)	170.4 (91.5–392.6)	<0.017
AL (mm)	1.8 (1.7–2.3)	2.4 (2.0–3.4)	<0.001
PD (mm)	1.7 (1.6–1.9)	1.9 (1.7–2.4)	<0.001
Number of sites (median (IQR)) ^b			
AL (≥3 mm)	30.0 (17.0–47.0)	54.0 (32.50–74.50)	<0.001
AL (≥4 mm)	5.0 (1.0–15.8)	25.0 (9.0–48.0)	<0.001
AL (≥5 mm)	1.0 (0.0–4.8)	13.0 (2.0–31.50)	<0.001
AL (≥6 mm)	0.0 (0.0–1.0)	4.0 (0.0–19.5)	<0.001
PD (≥4 mm)	2.0 (0.0–4.0)	6.0 (2.0–18.0)	<0.001
PD (≥5 mm)	0.0 (0.0–1.0)	2.0 (0.0–8.0)	<0.001
PD (≥6 mm)	0.0 (0.0–0.0)	1.0 (0.0–3.0)	0.001
BOP (n (%))	9.0 (4.0–20.0)	18.0 (10.0–31.5)	<0.001
Mean (SD)			
Plaque score (%)	93% (8.5)	91% (7.9)	0.059
Number of teeth	25.7 (4.7)	23.6 (6.2)	0.005

p values were Bonferroni–Holm corrected. A p value of <0.05 was considered statistically significant

p probability, BOP bleeding on probing, AL clinical attachment loss, DM2 diabetes mellitus type 2, IQR interquartile range, n number of participants, PISA periodontal inflamed surface area, PD probing pocket depth, SD standard deviation

^aResult of Chi-square test

^bResult of Mann–Whitney U test; other results are the results of the independent samples t test

Discussion

This study showed that PISA is a predictor of HbA1c, in conjunction with CRP, age, sex, and smoking, in healthy Indonesians. Thus, periodontitis may play a role in inducing impaired blood glucose control. Indeed, there is evidence that periodontitis may induce insulin resistance [18]. It is thought that periodontitis exerts an inflammatory and infectious burden as evidenced by increased levels of

CRP [3, 5, 6] which may increase HbA1c levels [19, 20, 36]. Accordingly, PISA predicted HbA1c together with CRP.

It is striking that out of the various methods to operationalize periodontitis severity, only the PISA emerged as a predictor of HbA1c. The other methods for determining the severity and extent of periodontitis did not contribute significantly to a model that predicts HbA1c. It may be that the PISA is a predictor of HbA1c because it reflects the amount of inflamed periodontal tissue, thereby predicting both infectious and inflammatory burden more accurately than other methods used to operationalize periodontitis [1, 18].

In Indonesian DM2 patients, no measure for periodontitis severity predicted HbA1c. The observed differences between DM2 patients and healthy Indonesians may be explained as follows. First, almost all DM2 patients used a combination of medication to control blood sugar, drugs that healthy Indonesians obviously did not take [37]. Use such medication may mask an effect of periodontitis on the HbA1c level in DM2 patients. Second, DM2 patients may have dietary restrictions, e.g., avoiding sugar rich foods. Such a diet may impact blood sugar and may mask an affect of periodontitis on HbA1c level.

Previously, a dose–response relationship was observed between PISA and HbA1c in DM2 patients from the Caribbean island, Curacao [18]. Additionally, an increased HbA1c was observed to be associated with increased severity of periodontitis [38]. Moreover, it has been

Table 3 Results of the multiple linear regression analysis healthy control group

Model predictors	β	p value of β	R^2	95% confidence interval of β
Model			0.21	
PISA (mm ²) ^a	0.0004	<0.05		0.00004 to 0.00076
Age (year)	0.0105	0.001		0.00431 to 0.01674
Male/female (1/0)	−0.1378	0.08		−0.29170 to 0.01595
Smoking (pack year)	0.0086	0.07		−0.00074 to 0.01805
CRP (mg/l)	0.0212	0.06		−0.00093 to 0.04343
Constant	4.9591	<0.001		4.62706 to 5.29121

A p value of <0.05 was considered statistically significant. Dependent variable—HbA1c; independent variables: PISA, age, sex, and smoking/pack years. Constant

p probability, β unstandardized coefficient, HbA1c glycated hemoglobin, periodontal inflamed surface area, CRP C-reactive protein

^a Other measures of periodontitis severity were not predictors of HbA1c

reported that periodontal treatment leads to improvement of glycemic control in DM2 patients [39]. These findings appear to be in sharp contrast with the findings in this study, i.e., PISA is not a predictor of HbA1c in a group of Indonesian DM2 patients. The difference between the current study and the study in Curacao [18] could be explained by differences in populations. First, substantial differences were observed in the frequency of antidiabetics used between DM2 from Indonesia and Curacao. Acarbose (33% vs. 5%), insulin (41% vs. 21%), and sulfonylurea (54% vs. 35%) were used more often in DM2 patients from Indonesia, while metformin 37 vs. 67% was used more often in DM2 patients from Curacao. Additionally, 16% of DM2 patients from Curacao used tolbutamide, while none of Indonesian DM2 patients used it. Differences in the frequency of antidiabetics used, may also explain the differences in results. As reported by Teeuw et al. [38] and Simpson et al. [39], HbA1c levels might be affected by the treatment DM2 patients receive, i.e., both the drugs they use and the periodontal treatment the patients have received. Thus, a drug-related reduction in HbA1c levels in patients not yet adequately periodontally treated might be responsible for the failure of PISA to predict HbA1c levels in our Indonesian DM2. Second, the average BMI of DM2 patients from Curacao was 31 while the average BMI of Indonesian DM2 patients was substantially lower, namely 25. BMI may be an effect modifier, i.e., a higher BMI may increase the potential insulin resistance inducing effect of periodontitis in DM2 patients [40, 41]. Third, there is clear difference in ethnicity between the two groups, i.e., one from Indonesia of Javanese origin and the other from Curacao of African-American origin. Possible differences in (health promoting) behavior, diet, and genes, related to the differences in ethnicity, may be another explanation for the differences in results [42, 43].

Conclusions

PISA was shown to be a predictor of HbA1c in healthy Indonesians in conjunction with CRP, age, sex, and smoking. This implies that periodontitis might contribute to insulin resistance through the inflammatory and infectious burden leading to DM2. In Indonesian patients, treated for DM2, PISA was not able to predict HbA1c.

Acknowledgments We would like to thank the following institutions for their cooperation in Yogyakarta, Indonesia: the Internal Medicine Department and the Dental Clinic of the Dr. Sardjito Hospital, the Prof. Soedomo Dental Hospital of the Faculty of Dentistry of Gadjah Mada University, the Diabetes Center of Yogyakarta International Hospital, and the Prodia Laboratory. This study was supported by a Bernoulli Foundation grant from the University of Groningen.

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Nesse W, Abbas F, van der Ploeg I, Spijkervet FK, Dijkstra PU, Vissink A (2008) Periodontal inflamed surface area: quantifying inflammatory burden. *J Clin Periodontol* 35:668–673
2. Noack B, Genco RJ, Trevisan M, Grossi S, Zambon JJ, De Nardin E (2001) Periodontal infections contribute to elevated systemic C-reactive protein level. *J Periodontol* 72:1221–1227
3. Craig RG, Yip JK, So MK, Boylan RJ, Socransky SS, Haffajee AD (2003) Relationship of destructive periodontal disease to the acute-phase response. *J Periodontol* 74:1007–1016
4. D'Aiuto F, Parkar M, Andreou G, Suvan J, Brett PM, Ready D, Tonetti MS (2004) Periodontitis and systemic inflammation: control of the local infection is associated with a reduction in serum inflammatory markers. *J Dent Res* 83:156–160
5. Salzberg TN, Overstreet BT, Rogers JD, Califano JV, Best AM, Schenkein HA (2006) C-reactive protein levels in patients with aggressive periodontitis. *J Periodontol* 77:933–939
6. Linden GJ, McClean K, Young I, Evans A, Kee F (2008) Persistently raised C-reactive protein levels are associated with advanced periodontal disease. *J Clin Periodontol* 35:741–747
7. Pitiphat W, Savetsilp W, Wara-Aswapati N (2008) C-reactive protein associated with periodontitis in a Thai population. *J Clin Periodontol* 35:120–125
8. Nakajima T, Yamazaki K (2009) Periodontal disease and risk of atherosclerotic coronary heart disease. *Odontology* 97:84–91
9. Persson GR, Persson RE (2008) Cardiovascular disease and periodontitis: an update on the associations and risk. *J Clin Periodontol* 35:362–379
10. Mealey BL, Oates TW (2006) American academy of periodontology. Diabetes mellitus and periodontal diseases. *J Periodontol* 77:1289–1303
11. Li X, Kolltveit KM, Tronstad L, Olsen I (2000) Systemic diseases caused by oral infection. *Clin Microbiol Rev* 13:547–558
12. Soell M, Hassan M, Miliauskaitė A, Haikel Y, Selimovic D (2007) The oral cavity of elderly patients in diabetes. *Diabetes Metab* 33 (Suppl 1):S10–S18
13. Hu FB, Meigs JB, Li TY, Rifai N, Manson JE (2004) Inflammatory markers and risk of developing type 2 diabetes in women. *Diabetes* 53:693–700
14. Wang Z, Hoy WE (2007) C-reactive protein and the risk of developing type 2 diabetes in Aboriginal Australians. *Diabetes Res Clin Pract* 76:37–43
15. Dehghan A, van Hoek M, Sijbrands EJ, Stijnen T, Hofman A, Witteman JC (2007) Risk of type 2 diabetes attributable to C-reactive protein and other risk factors. *Diabetes Care* 30:2695–2699
16. Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM (2001) C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA* 286:327–334
17. Taylor GW (1999) Periodontal treatment and its effects on glycemic control: a review of the evidence. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 87:311–316
18. Nesse W, Linde A, Abbas F, Spijkervet FK, Dijkstra PU, de Brabander EC, Gerstenbluth I, Vissink A (2009) Dose-response

- relationship between periodontal inflamed surface area and HbA1c in type 2 diabetics. *J Clin Periodontol* 36:295–300
19. Rodriguez-Moran M, Guerrero-Romero F (1999) Increased levels of C-reactive protein in noncontrolled type II diabetic subjects. *J Diabetes Complications* 13:211–215
 20. Gomez JM, Vila R, Catalina P, Soler J, Badimon L, Sahun M (2008) The markers of inflammation and endothelial dysfunction in correlation with glycated haemoglobin are present in type 2 diabetes mellitus patients but not in their relatives. *Glycoconj J* 25:573–579
 21. Pfutzner A, Forst T (2006) High-sensitivity C-reactive protein as cardiovascular risk marker in patients with diabetes mellitus. *Diabetes Technol Ther* 8:28–36
 22. Flores-Alfaro E, Parra-Rojas I, Salgado-Bernabe AB, Chavez-Maldonado JP, Salazar-Martinez E (2008) Cardiovascular risk evaluated by C-reactive protein levels in diabetic and obese Mexican subjects. *Circ J* 72:1170–1174
 23. Takeda M, Ojima M, Yoshioka H, Inaba H, Kogo M, Shizukuishi S, Nomura M, Amano A (2006) Relationship of serum advanced glycation end products with deterioration of periodontitis in type 2 diabetes patients. *J Periodontol* 77:15–20
 24. Lim LP, Tay FB, Sum CF, Thai AC (2007) Relationship between markers of metabolic control and inflammation on severity of periodontal disease in patients with diabetes mellitus. *J Clin Periodontol* 34:118–123
 25. Mihadja L, Delima Manz HS, Ghani L, Soegondo S (2009) Prevalence and determinants of diabetes mellitus and impaired glucose tolerance in indonesia (a part of basic health research/risekdas). *Acta Med Indones* 41:169–174
 26. Wild S, Roglic G, Green A, Sicree R, King H (2004) Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 27:1047–1053
 27. Timmerman MF, Van der Weijden GA, Armand S, Abbas F, Winkel EG, Van Winkelhoff AJ, Van der Velden U (1998) Untreated periodontal disease in Indonesian adolescents. Clinical and microbiological baseline data. *J Clin Periodontol* 25:215–224
 28. Susanto H, Nesse W, Dijkstra PU, Agustina D, Vissink A, Abbas F (2011) Periodontitis prevalence and severity in Indonesians with type 2 diabetes. *J Periodontol* 82:550–557
 29. de Jong KJ, Oosting J, Peters GJ, Abraham-Inpijn L (1992) Detecting medical problems in dentistry: a survey of 4,087 patients in The Netherlands. *Eur J Med* 1:23–29
 30. de Jong KJ, Abraham-Inpijn L (1994) A risk-related patient-administered medical questionnaire for dental practice. *Int Dent J* 44:471–479
 31. Armitage GC (2004) The complete periodontal examination. *Periodontol* 2000(34):22–33
 32. Mercado FB, Marshall RI, Klestov AC, Bartold PM (2001) Relationship between rheumatoid arthritis and periodontitis. *J Periodontol* 72:779–787
 33. Vettore MV, Leal M, Leao AT, da Silva AM, Lamarca GA, Sheiham A (2008) The relationship between periodontitis and preterm low birthweight. *J Dent Res* 87:73–78
 34. Alberti KG, Zimmet PZ (1998) Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 15:539–553
 35. WHO Expert Consultation (2004) Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 363:157–163
 36. Nabipour I, Vahdat K, Jafari SM, Beigi S, Assadi M, Azizi F, Sanjdideh Z (2008) Elevated high sensitivity C-reactive protein is associated with type 2 diabetes mellitus: the Persian Gulf Healthy Heart Study. *Endocr J* 55:717–722
 37. Kooy A, de Jager J, Leher P, Bets D, Wulffele MG, Donker AJ, Stehouwer CD (2009) Long-term effects of metformin on metabolism and microvascular and macrovascular disease in patients with type 2 diabetes mellitus. *Arch Intern Med* 169:616–625
 38. Simpson TC, Needleman I, Wild SH, Moles DR, Mills EJ (2010) Treatment of periodontal disease for glycaemic control in people with diabetes. *Cochrane Database Syst Rev* 5: CD004714
 39. Teeuw WJ, Gerdes VE, Loos BG (2010) Effect of periodontal treatment on glycemic control of diabetic patients: a systematic review and meta-analysis. *Diabetes Care* 33:421–427
 40. Genco RJ, Grossi SG, Ho A, Nishimura F, Murayama Y (2005) A proposed model linking inflammation to obesity, diabetes, and periodontal infections. *J Periodontol* 76:2075–2084
 41. King GL (2008) The role of inflammatory cytokines in diabetes and its complications. *J Periodontol* 79:1527–1534
 42. Petersen PE (2004) Challenges to improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. *Int Dent J* 54:329–343
 43. Siriwardena AN (2004) Specific health issues in ethnic minority groups. *Clin Cornerstone* 6:34–40, discussion 41–2

Copyright of Clinical Oral Investigations is the property of Springer Science & Business Media B.V. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.