Abnormal mandibular growth and the condylar cartilage

Pertti Pirttiniemi*, Timo Peltomäki**, Lukas Müller** and Hans. U. Luder*** *Department of Oral Development and Orthodontics, Institute of Dentistry, University of Oulu, Finland, **Department of Orthodontics and Pediatric Dentistry and ***Institute of Oral Biology, University of Zurich, Switzerland

SUMMARY Deviations in the growth of the mandibular condyle can affect both the functional occlusion and the aesthetic appearance of the face. The reasons for these growth deviations are numerous and often entail complex sequences of malfunction at the cellular level.

The aim of this review is to summarize recent progress in the understanding of pathological alterations occurring during childhood and adolescence that affect the temporomandibular joint (TMJ) and, hence, result in disorders of mandibular growth. Pathological conditions taken into account are subdivided into (1) congenital malformations with associated growth disorders, (2) primary growth disorders, and (3) acquired diseases or trauma with associated growth disorders.

Among the congenital malformations, hemifacial microsomia (HFM) appears to be the principal syndrome entailing severe growth disturbances, whereas growth abnormalities occurring in conjunction with other craniofacial dysplasias seem far less prominent than could be anticipated based on their oftendisfiguring nature. Hemimandibular hyperplasia and elongation undoubtedly constitute the most obscure conditions that are associated with prominent, often unilateral, abnormalities of condylar, and mandibular growth. Finally, disturbances of mandibular growth as a result of juvenile idiopathic arthritits (JIA) and condylar fractures seem to be direct consequences of inflammatory and/or mechanical damage to the condylar cartilage.

Introduction

The size of the mandible, including the corpus, ramus, and condyle, as well as the timing and amount of condylar growth, vary considerably between individuals. Factors potentially contributing to this individual variation are the extent of masticatory action related to the consistency of the diet (Kiliaridis *et al.*, 1999) and, as shown recently (Van Erum *et al.*, 1997; Zhou *et al.*, 2005), genetic predisposition.

Distinct from even extreme cases of individual variation, the deviations from normality considered in this review constitute examples of abnormal mandibular and/or condylar growth that are related to truly pathological alterations. Based on their aetiology and time of appearance, they can be classified as (1) congenital malformations with associated growth disorders, (2) primary growth disorders, and (3) acquired diseases or trauma with associated growth disorders.

Congenital malformations with associated growth disorders

The most prevalent craniofacial malformation involving mostly unilateral condylar and ramal underdevelopment, as well as greatly variable abnormalities of the external and middle ear, is hemifacial microsomia [HFM; (Online Mendelian Inheritance in Man database of Johns Hopkins University: http://www.ncbi.nlm.nih.gov/sites/entrez?db= omim) OMIM 164210] with an estimated frequency of about 1/5000–6000 live births. Because of the similarity with the

manifestations of Goldenhar syndrome that is characterized additionally by vertebral defects and epibulbar dermoids, the two conditions have been combined under the term oculoauriculo-vertebral (OAV) spectrum (Gorlin et al., 2001). Its aetiology seems to be heterogeneous. Genetic factors, in particular chromosomal anomalies such as deletions of 5p, 6q, 8q, 18q, and 22q; duplications of 22q; and trisomies 7, 9, 18, and 22, have been implicated with the OAV spectrum. On the other hand, anomalies similar to those of HFM and Goldenhar syndrome resulted from early foetal exposure to thalidomide or retinoic acid (Gorlin et al., 2001). In fact, a HFM-like phenotype could be reproduced using triazine, an anti-cancer drug, in mice and thalidomide in monkeys (Poswillo, 1973). Poswillo (1973) suggested that haemorrhages at the anastomosis of the external carotid and stapedial artery were responsible for the anomalies that seemed to increase in severity with the extension of local tissue damage. However, there is also considerable overlap in manifestations of the OAV spectrum, the retinoic acid syndrome, and the DiGeorge syndrome (OMIM 188400), a condition attributed to the loss of the T-box transcription factor TBX1 due to chromosomal deletion 22q11.2 (Botto et al., 2003; Packham and Brook, 2003). Common to all of these three entities are cono-truncal cardiovascular defects, which led Johnston and Bronsky (1995) to suspect that defective neural crest cell development could be the primary cause of HFM.

The extent of temporomandibular joint (TMJ) involvement in the OAV spectrum largely determines the timing and type of treatment (Caccamese *et al.*, 2006). Therefore, HFM is often classified based on the degree of TMJ dysmorphology (Mulliken and Kaban, 1987). The mildest forms (Kaban type I), such as the example shown in Figure 1a,b, appear to be characterized by a slightly hypoplastic mandibular condyle and thinner than normal condylar cartilage, but fairly normal hypertrophy of the chondrocytes and endochondral ossification (Figure 1c,d). Hence, mandibular growth can be expected to be only slightly deficient, justifying the recommendation to treat these cases at skeletal maturity using conventional orthognathic procedures (Caccamese et al., 2006). In contrast, the severe forms of HFM (Kaban type III), such as the case illustrated in Figure 1e,f, exhibit aplasia or severe hypoplasia of the condyle. Even if present, these condyles seem to completely lack condylar cartilage and endochondral ossification (Figure 1g,h). Hence, mandibular growth on the affected side most likely comes to an early standstill; facial asymmetry must be assumed to worsen progressively, and early intervention utilizing, for example, rib grafts seems indicated to avoid secondary tilting of the maxilla.

Unlike the manifestations of the OAV spectrum, those of mandibulofacial dysostosis (MFD) are always bilateral. The prototype MFD, Treacher Collins syndrome (TCS; mainly in Europe also known as Franceschetti-Zwahlen-Klein syndrome; OMIM 154500), occurs at a frequency of about 1/25 000–50 000 live births. Abnormalities include underdeveloped supraorbital ridges, downward sloping of the palpebral fissures, and hypoplasia of the zygomatic bones, as well as of the mandibular rami and condyles (Gorlin *et al.*, 2001).

TCS is an autosomal dominant disorder with variable expressivity. Although MFD can also be caused by chromosomal abnormalities (Stevenson *et al.*, 2007), most individuals with TCS bear a mutation of the *TCOF1* gene. However, even among carriers of such a specific genetic defect, the phenotype varies markedly and is sometimes so mild that the disorder may go undetected clinically (Teber *et al.*, 2004). *TCOF1* encodes Treacle, a nucleolar protein. When it is defective, the biogenesis of



Figure 1 Hemifacial microsomia (HFM). (a–d) A 1.5-year-old girl with marked chin deviation (a), but mild involvement of the external ear (b) associated with HFM on the left side; (c) overview micrograph of the left condyle replaced by a costochondral graft and (d) detail of condylar cartilage marked by the rectangle in (c); note the diameter of the condyle (c) and the thickness of the hypertrophic cartilage (HC; d) that are approximately two-thirds of the dimensions seen in an age-matched healthy specimer; in contrast, endochondral ossification (d; arrows) is inconspicuous. (e–h) A 14-year-old girl with marked chin deviation (e) and severe involvement of the external ear (f) due to HFM on the left side. The overview micrograph of the resected condyle (g) and the detail of the condylar articular surface (h) marked by the rectangle in (g) reveal that the condyle is small for the age and condylar cartilage is completely missing. (c, d, g, and h) toluidine blue; original magnifications (c and g) ×6.3 and (d and h) ×80.

mature ribosomes in neuroepithelial and neural crest cells is impaired, the formation and proliferation of neural crest cells is disturbed and, finally, the number of neural crest cells that migrate into the branchial arches is deficient (Dixon *et al.*, 2006).

In agreement with this pathogenetic mechanism, the examination of a male foetus of 130 mm crown-rump length diagnosed as having TCS revealed that at this early pre-natal stage, the developing maxillary and zygomatic bones were hypoplastic. As far as the mandible was concerned, the rami and corpus were unusually short and abnormally shaped, while condylar cartilage was missing on one side and markedly deficient on the other side (Behrents et al., 1977). On the other hand, clinical and cephalometric findings from TCS patients repeatedly show that the craniofacial skeletal pattern observed during infancy remained fairly stable during further development (Roberts et al., 1975; Posnick and Ruiz, 2000). In fact, longitudinal changes in the length of the mandible reported by Roberts et al. (1975) appear to deviate little from control values, suggesting that the disorder of post-natal condylar growth associated with TCS may be far less extensive than the primary, pre-natal establishment of an abnormal skeletal pattern.

A clinically significant entity, also involving potential condylar growth problems, is the Pierre Robin sequence (PRS; OMIM 261800) with a prevalence of about 1/2000–8500 live births. It is characterized by mandibular micrognathia, cleft palate, and glossoptosis (Gorlin *et al.*, 2001). Although the aetiology of the disorder so far has not been elucidated, a recent systematic analysis of several PRS patients and the finding of a familial translocation of chromosomes 2 and 17 in one case suggested that genetic defects of *SOX9*, a gene encoding a transcription factor that is crucial for the differentiation of chondrocytes and chondrogenesis, could play a role in the development of PRS (Jakobsen *et al.*, 2007).

A first issue of ongoing debate with respect to PRS applies to the role played by tongue interposition as the cause for the formation of isolated palatal clefts. It has been argued that palatal clefts result from insufficient downwardforward displacement of the tongue due to a retroposition of the mandible. Alternatively, development of the tongue itself could be defective and result in insufficient displacement of the mandible. From the analysis of mice carrying a genetic defect predisposing to the development of a phenotype resembling PRS, Schubert et al. (2005) concluded that primary mandibular retrognathia, rather than deficient growth of the tongue, was responsible for the formation of the clefts. It should be borne in mind, however, that downward and forward growth of the mandibular arch prior to the fusion of the secondary palate is accounted for by Meckel's cartilage, rather than secondary condylar cartilage that develops considerably later (Diewert, 1982, 1985). When considering this, the suggested aetiological

involvement of defects in *SOX9* (Jakobsen *et al.*, 2007) appears quite conceivable.

A second controversial issue related to PRS is the question as to whether catch-up mandibular growth is able to compensate, at least in part, for the mandibular deficiency. While Figueroa *et al.* (1991) supported the possibility of catch-up growth, Laitinen and Ranta (1992), Hermann *et al.* (2004), and, recently, Eriksen *et al.* (2006) showed that post-natal growth of the mandible in children with PRS is normal and comparable with that of other children with clefts. Also, children with PRS associated with mandibular hypodontia had smaller mandibles than subjects with PRS and all permanent mandibular teeth, and this pattern did not improve with further growth (Suri *et al.*, 2006).

In addition to these relatively frequent syndromes, there are additional, but very rare conditions affecting the size and shape of the mandible and possibly also condylar growth. Examples are acrofacial dysostosis 1 (OMIM 154400) or Nager syndrome, that can be associated with condylar ankylosis (Halonen et al., 2006), and Turner syndrome with a short mandibular body and posterior rotation of the mandible (Peltomäki et al., 1989; Rongen-Westerlaken et al., 1993; Babić et al., 1997; Perkiömäki et al., 2005). Hemifacial atrophy (OMIM 141300) or Parry-Romberg syndrome may affect mandibular growth and lead to progressive facial asymmetry (Buonaccorsi et al., 2005), while Hallerman-Streiff syndrome (OMIM 234100) is characterized by a narrow upper dental arch, extensive hypodontia, and a very small, posteriorly rotated mandible, giving the impression of a 'bird face' (Defraia et al., 2005).

In Silver-Russell syndrome (OMIM 180860), the mandible as well as the maxilla are small and retrognathic. Left– right differences in mandibular growth causing facial asymmetries are common (Kotilainen *et al.*, 1995). Similarly, children with Marfan syndrome (OMIM 154700), a connective tissue disorder resulting from mutations in the *FBN1* gene that encodes the microfibril component fibrillin-1, often have a retrognathic maxilla and mandible (Westling *et al.*, 1998).

Although less common than mandibular hypoplasia, oversized mandibles do occur in association with some syndromes. Acromegaly is a condition of growth hormone overproduction. When the disease becomes manifest, the mandible enlarges rapidly and a progenic habit develops, sometimes in combination with sleep appoea syndrome due to changes in the laryngeal soft tissues. Recently, an inherited form of acromegaly due to a germ-line mutation in the AIP gene (OMIM 605555) has been detected. As a result, the defective aryl hydrocarbon receptor-interacting protein leads to a predisposition for pituitary adenoma (Georgitsi et al., 2007). In Proteus syndrome (OMIM 176920) that is characterized by striking facial abnormalities, one of two subforms shows unilateral condylar overgrowth causing progressive craniofacial asymmetry (Kreiborg et al., 1991). The stimulating influence of sex chromosomes on the growth of the mandible can be clearly seen in patients with Klinefelter's syndrome (47,XXY). In comparison with normal control females, these individuals have larger mandibles, in particular larger mandibular bodies, which cause marked mandibular prognathism.

Finally, there are craniofacial malformations that do not affect the jaws directly, but lead to indirect, presumably compensatory, alterations of otherwise relatively normal condylar growth. For example, in cases of craniosynostoses such as Crouzon (OMIM 123500), Pfeiffer (OMIM 101600), and Apert (OMIM 101200) syndromes, unusual transverse mandibular growth may be regarded as an attempt at adapting to the impaired expansion of the cranial vault (Boutros *et al.*, 2007).

In summary, disorders of mandibular and condylar growth associated with craniofacial malformations appear far less extensive than could be expected, when considering the mostly markedly disfiguring conditions. Rather than disturbed growth, the primary establishment of an aberrant craniofacial skeletal pattern seems to account for the clinical impression of abnormality. Obvious exceptions from this rule are, however, severe cases of HFM, where unilaterally deficient or completely absent condylar growth results in progressive facial asymmetry.

Primary growth disorders

Condylar hyperactivity can be clearly identified only when it occurs unilaterally. Bilateral symmetric cases are very difficult to delineate against mandibular prognathism and seem to be very rare (Obwegeser, 2001). Unilateral condylar hyperactivity typically becomes apparent at some time during the growth period, most often during childhood. Hence, it constitutes a true disorder of growth (Obwegeser, 2001). The aetiology is largely unknown. There are reports of cases where condylar trauma during childhood later manifested itself as hyperplastic growth (Jacobsen and Lund, 1972; Rubenstein and Campbell, 1985). Other possible causes taken into consideration, but so far not substantiated, are inflammation, hypervascularization, and unspecified genetic factors (Obwegeser, 2001).

In the literature, condylar hyperactivity is commonly referred to as condylar hyperplasia. This term was coined by Rushton (1944, 1946), although it did not go unnoticed that apart from the condyle, the entire ramus and corpus of the hemimandible on the affected side were enlarged, resulting in gross distortion of the lower face without significant deviation of the chin (Figure 2a). However, unilateral condylar hyperactivity can also manifest itself in elongation, rather than increase in volume, of the condylar neck, ramus, and corpus, leading to facial asymmetry with marked deviation of the chin to the unaffected side (Figure 2b). For this form of growth anomaly, Obwegeser and Makek (1986) introduced the term hemimandibular elongation and referred to the disorder described by Rushton (1944, 1946) as 'hemimandibular hyperplasia'.

Considering that the entire hemimandible, rather than only the mandibular head, is affected in cases of unilateral condylar hyperactivity irrespective of the clinical classification, the question arises whether the causative disturbance really lies in the condyle. Most previous reports on histological findings, reviewed recently by Luder (2001), mentioned alterations of the condylar cartilage that seemed to indicate unusually rapid growth and/or an abnormally long duration of growth. Slootweg and Müller (1986) even suggested a classification of condylar hyperplasia based on the appearance of the cartilage. However, compared with normal age-matched specimens, differences in cartilage morphology and thickness proved rather small (Luder, 2001). Thus, given the variation in the timing of mandibular growth (Björk, 1963), caution seems to be warranted when drawing conclusions as to individual growth velocity based on the appearance of the condylar cartilage.

Another structural feature observed consistently in hyperplastic condyles is the distribution of cartilage rests in the subchondral spongiosa. As noted by Rushton (1944), remnants of cartilage matrix occur at abnormally large distances from the front of the erosion. When comparing the various forms of condylar hyperactivity, this seems to be particularly true for cases of hemimandibular hyperplasia that additionally reveal a conspicuous arrangement of mixed cartilaginous-bony trabeculae lacking a clear orientation (Figure 2c). In contrast, condyles from subjects with hemimandibular elongation exhibited cartilage of normal structure and thickness, subchondral cartilage rests at normal distances from the zone of erosion, and a spongiosa comprising well-orientated bone trabeculae (Figure 2d; Luder, 2001).

These findings indicate that in cases of condylar hyperactivity, the primary vascular invasion and resorption of cartilage, that determine its thickness, are fairly normal or at least keep pace with cartilage growth. In contrast, remodelling of the primary spongiosa, that is responsible for the ultimate removal of cartilage remnants and arrangement of the bone trabeculae, may be disturbed or unbalanced in hemimandibular hyperplasia, but normal in hemimandibular elongation (Luder, 2001). It is, thus, possible that distinct pathogenetic mechanisms account for the two clinical forms of condylar hyperactivity, although this has yet to be confirmed by additional, more comprehensive studies.

The question as to whether condylar growth is active or has ceased is critical for selecting the appropriate treatment procedure. When growth is still ongoing, high condylectomy, although controversial, is considered an option to avoid secondary, adaptive deformation of the maxilla. On the other hand, corrective orthopaedic surgery should be envisaged only when condylar growth has ceased (Marchetti *et al.*, 2000; Obwegeser, 2001; Deleurant *et al.*, 2008). As



Figure 2 Hemimandibular hyperplasia (HH; a, c, and e) and hemimandibular elongation (HE; b, d, and f). Dental pantomographs of a 12-year-old boy (a) and a 13-year-old girl (b); note the distortion of the entire right hemimandible associated with HH (a) and the midline deviation to the left due to HE on the right side (b). (c–f) Overview micrographs of the condyles (c and d) resected in the subjects shown in (a) and (b) and details of the condylar cartilage (e and f) marked by the rectangles in (c) and (d), respectively; note the arrangement of the subchondral bone trabeculae (c and d; arrows) and the front of endochondral ossification (e and f; arrowheads). (c–f) toluidine blue; original magnifications (c and d) ×6.3 and (e and f) ×80.

the method of choice for the assessment of condylar growth activity, planar or three-dimensional quantification of 99m technetium methylene diphosphonate (99m Tc-MDP) uptake using plane scintigraphy or single photon emission computed tomography are strongly recommended (Chan *et al.*, 2000; Pripatnanont *et al.*, 2005). However, it should be borne in mind that all these methods of bone scanning, although highly sensitive, are non-specific, that is they do not yield any clue as to the reason for an observed asymmetry in condylar activity (Obwegeser, 2001).

Acquired diseases or trauma with associated growth disorders

Juvenile idiopathic arthritis (JIA) is a chronic inflammatory disease of unknown aetiology, which is present for longer than 6 weeks and starts before the age of 16 years (Petty *et al.*, 1998). It comprises seven subtypes (systemic arthritis, rheumatoid factor-negative and -positive polyarthritis,

oligoarthritis, enthesitis-related arthritis, psoriatic arthritis, and others) based on clinical symptoms during the first 6 months of the disease. Overall, JIA is the most common form of arthritis in children. Its frequency is about 1-2/1000in most populations (Saurenmann *et al.*, 2007) with a ratio of girls to boys of about 3:2 (Andersson Gäre *et al.*, 1987). The disease is characterized by variable degrees of joint inflammation, joint destruction, and progressive disability (Palmisani *et al.*, 2006).

Although the aetiology of JIA is unknown, it has clear autoimmunological characteristics. There is, however, no evidence indicating that some specific antigen is responsible. Rather, it is presumed that many different factors, to varying degrees, lead to the outbreak of the disease (Sen, 2005). For this reason, the serological demonstration of HLA-B27, antinuclear antibodies, and rheumatoid factor are of little prognostic value for the course of JIA (Ilowite, 2002).

Histopathologically, JIA is characterized by hypertrophic inflammatory synovitis with cellular infiltration and

proliferation of blood vessels. In the chronic state of the disease, the articular surfaces are covered by a pannus, a tumour-like mass of inflamed granulation tissue. The expansion of the pannus is enhanced by extensive formation of new blood vessels and results in cartilage and joint destruction that may occur early in the course of the disease (Yang *et al.*, 2002).

Reported frequencies of TMJ involvement in JIA vary from approximately 17 to 87 per cent, probably depending on whether all subtypes of the disease have been taken into account and whether the diagnoses have been based on a purely clinical, a radiological, or even a magnetic resonance imaging (MRI) examination (Mayne and Hatch, 1969; Rönning et al., 1974; Küseler et al., 1998; Billiau et al., 2007). In all subtypes, one or both TMJs can be affected and may even be the initial joints involved (Karhulahti et al., 1990; Küseler et al., 1998; Martini et al., 2001). While Billiau et al. (2007) claimed that condylar damage was unrelated to JIA subtype and disease activity, severity, or duration, Säilä et al. (2004) reported that patients reacting positive for antinuclear antibodies more frequently exhibited TMJ involvement than histocompatability locus antigenpositive individuals, and Pedersen et al. (2001), based on a study using enhanced MRI, concluded that TMJ involvement was particularly frequent in polyarticular onset JIA. Also based on MRI, Küseler et al. (2005) found a prevalence of 26 per cent for pannus formation in the TMJ and a frequency of 71 per cent regarding condylar erosions.

Because of its suitability to detect even small initial destructive lesions, MRI has become the gold standard for TMJ examinations in subjects with JIA (Figure 3). It has even been suggested that a panographic examination is not indicated when a MRI is obtained (Arabshahi and Cron, 2006; Helenius *et al.*, 2006). High-resolution ultrasonography (US) has been proposed as an alternative to MRI, because it can be used while the joint is functioning, thus allowing

assessment of disc movements (Jank *et al.*, 2007). However, a closer look at the publication of those authors shows that only severe destructive TMJ changes had been investigated. In agreement with this observation, a recent, unpublished pilot study indicated that US is not able to detect early TMJ arthritis before destructive alterations have occurred. These results suggest that while US is a valuable diagnostic imaging method for the TMJ, it cannot yet replace a MRI investigation.

Histopathologically, JIA seems to affect the lower and upper TMJ compartments to markedly different degrees. In TMJs exhibiting severe radiographic alterations where condylar replacement with costochondral grafts seemed indicated, the entire lower joint compartments were filled with masses of granulomatous tissue, while the upper compartments appeared macroscopically healthy. In contrast, condyles showed localized to total cartilage destruction and prominent inflammatory infiltrates of the subchondral bone marrow (Svensson et al., 2001). Evidence regarding the pathogenesis of cartilage destruction is available only from investigations of rheumatoid arthritis and osteoarthritis, both of which occur in adulthood and, therefore, do not result in growth disturbances, although they may entail marked remodelling of articular tissues. According to these studies (Kanyama et al., 2000; Miyamoto et al., 2002; Gepstein et al., 2003; Goldring, 2003; Tiilikainen et al., 2005; Malemud, 2007), inflammatory cells populating the synovial membrane release pro-inflammatory cytokines, chemokines, angiogenic factors, and proteinases. Among the secreted products that appear to exert the greatest effect on cartilage loss are interleukin-1 (IL-1) and tumour necrosis factor α (TNF α). The most potent angiogenic factor is vascular endothelial growth factor which plays a critical role, because it enhances growth of the vasculature in the inflamed synovial tissue. IL-1 stimulates chondrocytes to produce matrix metalloproteinases (MMPs) which, in their active



Figure 3 Juvenile idiopathic arthritis (JIA). Lateral magnetic resonance views of the right temporomandibular joint affected by JIA (a) and the contralateral healthy joint (b) of a 13-year-old girl; note the erosion of the right condyle (a; arrowhead) and the position of the intermediate zone of the articular discs (a and b; arrows).

forms, effectively degrade collagens and proteoglycans. In addition, members of the aggrecanase (ADAMTS) family, particularly ADAMTS-4 and ADAMTS-5, seem to participate in cartilage destruction (Yoshida *et al.*, 2006). Interestingly, both MMPs and aggrecanases are also involved in remodelling processes elicited by experimental changes in dietary loading (Pirttiniemi *et al.*, 2004; Yu *et al.*, 2007).

JIA involving the TMJ is associated with characteristic facial changes, in particular a short mandibular ramus and backward-rotated mandibular corpus, prominent antegonial notching, and mandibular retrognathia (Rönning et al., 1974; Björk and Skieller, 1985; Hanna et al., 1996; Mericle et al., 1996; Kjellberg, 1998). Whereas Twilt et al. (2006) found that in comparison with age-matched healthy individuals, patients with JIA, regardless of their TMJ status, exhibited retrognathia and posterior rotation of the mandible, craniofacial alterations were reported by Billiau et al. (2007) to be related to the presence of radiographic condylar damage, even if this may be mild. Notably, however, bony erosion seems to occur later in JIA than in adult rheumatoid arthritis (Barriga et al., 1974), and condylar resorption seems to be present for some time before bone destruction is radiographically detectable (Küseler et al., 1998; Twilt et al., 2006). Thus, the destruction of condylar cartilage, irrespective of its severity, appears to entail significant disturbances of mandibular growth. On the other hand, recent longitudinal studies (Twilt et al., 2007, 2008) revealed that radiographic signs of condylar damage completely disappeared in about half and improved in another fifth of examined patients, while they worsened in a few subjects that exhibited particularly high disease activity.

Abnormal position or displacement of the TMJ disc has emerged relatively recently as a possible cause of mandibular growth disturbances. Disc disorders are anything but rare in childhood. While no cases of abnormal disc position could be detected in a sample of children ranging in age from 2 months to 5 years (Paesani *et al.*, 1999), Ribeiro *et al.* (1997) observed dislocated discs in 11 per cent of children aged from 6 to 11 years. Also, audible joint sounds, which may be associated with disc disorders, are relatively frequent, even in clinically healthy children. Heikinheimo *et al.* (1989) reported a frequency of 25–27 per cent, while Henrikson *et al.* (2000), as well as Henrikson and Nilner (2000), noticed joint sounds, although with fluctuating intensity, in 39 per cent of the children examined.

In an experiment using growing rabbits, the surgical induction of unilateral anterior disc displacement resulted in an asymmetric reduction of ramal growth and mandibular length, before visible osteoarthrotic TMJ changes developed (Legrell and Isberg, 1998, 1999; Legrell *et al.*, 1999). The authors suggested that disc dislocation *per se* had a primary adverse effect on condylar growth. Supporting these findings, Bryndahl *et al.* (2006) showed that

bilateral anterior disc displacement in a growing animal causes significant mandibular retrognathia. As complete maxillomandibular immobilization does not seem to affect mandibular growth in spite of significant histological alterations in condylar cartilage (Isacsson et al., 1993), it is doubtful that the observed disturbances of condylar growth resulted from a mechanical impairment of anterior condylar excursions, which conceivably could have been caused by the displaced disc. Alternatively, the adverse effects on condylar growth could also have been the consequence of an altered masticatory function (Legrell and Isberg, 1998, 1999). On the other hand, an experimental disc perforation created in growing rabbits initially led to increased cell proliferation in the condylar cartilage. At longer intervals following surgery, distinct arthrotic changes, comprising osteophytes and flattening of the condyle developed, but at still later time points, these changes were milder, suggesting some local adaptation to the disc perforation (Narinobou et al., 2000).

Mandibular trauma during childhood involves the condylar region in 36–50 per cent of subjects (Baumann *et al.*, 2004). Mandibular fractures are estimated to be about twice as frequent as noticed or diagnosed, as many of them occur during early childhood and often pass with little discomfort. The consequences of a trauma to the condyle depend on its location. In the case of intracapsular fractures, the fragments are seldom severely dislocated, but there is an increased risk for ankylosis, particularly in children younger than 3 years of age (Baumann *et al.*, 2004). If the fracture affects the condylar neck and, thus, is extracapsular, the condylar head often becomes dislocated, almost always in a forward–medial direction.

Long-term complications of both intra- and extracapsular fractures, such as the development of facial asymmetry or mandibular retrognathism and anterior open bite as well as TMJ ankylosis or painful temporomandibular disorders, seem to be rare and, if present, mild (Kellenberger et al., 1996; Marianowski et al., 2003). A new condyle can even be generated and facial symmetry may thus be recovered. An ankylosis of the TMJ as a consequence of a childhood condylar trauma is very rare (Marianowski et al., 2003). In a prospective study of 38 growing patients with fractures of the condylar neck, Lund (1974) found that in a majority of the subjects, greater than normal, compensatory growth occurred on the affected side, and no significant facial asymmetry developed. In the same study, a remarkable potential for post-traumatic condylar remodelling was noticed, which in some instances resulted in close to complete regeneration of a new mandibular head.

This potential for remodelling is demonstrated in the case of a 22-year-old female illustrated in Figure 4. As a consequence of a bicycle accident, she suffered a fracture of the left mandibular neck with forward–medial displacement of the condyle (Figure 4a,b). Conservative functional treatment was started only 1 week later, when she finally



Figure 4 Condylar fracture. Dental pantomographs (a and b) of a fracture of the left condyle suffered by a 22-year-old female as a consequence of a bicycle accident. Microradiograph of the left condyle (c), overview micrograph of the left temporomandibular joint (d), and detail of condylar cartilage (e) marked by the rectangle in (d) 2.5 months after the accident; note the old (light) and newly formed (dark) condylar bone (c), the dislocation of the condyle relative to the mandibular fossa (d) as well as in (e) the unusual vascularization of the articular fibrous layer (arrows) and presence of hypertrophic cartilage and endochondral ossification (arrowheads). (d and e) Toluidine blue; original magnifications (c) $\times 6.3$, (d) $\times 3$, and (e) $\times 80$.

went to see an emergency doctor. Two and a half months after the accident, she died from a cause unrelated to the first event, and an autopsy was carried out. The dislocated condyle exhibited conspicuous vascularized connective tissue underneath the articular surface and hypertrophic cartilage involved in endochondral ossification (Figure 4d,e), which could not be expected in a normal mandibular head of a female of this age. As a result of endochondral ossification, a remarkable amount of new bone had apparently been formed at the insertion of the lateral pterygoid muscle as well as at the zenith and posterior slope of the condyle (Figure 4c). The location of the hypertrophic cartilage and new bone formation gave the impression that reactivated condylar growth tended to re-establish a normal condyle-fossa relationship. Although no final conclusion can be drawn, this conceivably could have occurred in response to tension exerted by the stretched capsular apparatus, to post-traumatic inflammatory processes, or to a combination of both.

Owing to the significant progress in research techniques, our knowledge regarding the aetiology and pathogenesis of disorders affecting condylar cartilage has increased considerably during the last decades. At the same time, the diagnostic possibilities of the TMJ have improved greatly and allow early diagnoses of initial pathological processes. It is to be hoped that continuing efforts in basic and clinical research will eventually shed light on the remaining obscure forms of mandibular growth disorders and allow the design of new strategies for their prevention and treatment.

Address for correspondence

Pertti Pirttiniemi Department of Oral Development and Orthodontics Institute of Dentistry Box 5281 90014 university of Oulu Finland E-mail: pertti.pirttiniemi@oulu.fi

References

- Andersson Gäre B *et al.* 1987 Incidence and prevalence of juvenile chronic arthritis: a population survey. Annals of the Rheumatic Diseases 46: 277–281
- Arabshahi B, Cron R Q 2006 Temporomandibular joint arthritis in juvenile idiopathic arthritis: the forgotten joint. Current Opinion in Rheumatology 18: 490–495
- Babić M, Glišić B, Šćepan I 1997 Mandibular growth pattern in Turner's syndrome. European Journal of Orthodontics 19: 161–164

- Barriga B, Lewis T M, Law D B 1974 An investigation of the dental occlusion in children with juvenile rheumatoid arthritis. The Angle Orthodontist 44: 329–335
- Baumann A, Troulis M J, Kaban L B 2004 Facial trauma II: dentoalveolar injuries and mandibular fractures. In: Kaban L B, Troulis M J (eds). Pediatric oral and maxillofacial surgery Saunders, Philadelphia, pp. 441–462
- Behrents R G, McNamara J A, Avery J K 1977 Prenatal mandibulofacial dysostosis (Treacher Collins syndrome). The Cleft Palate Journal 14: 13–34
- Billiau A D, Hu Y, Verdonck A, Carels C, Wouters C 2007 Temporomandibular joint arthritis in juvenile idiopathic arthritis: prevalence, clinical and radiological signs, and relation to dentofacial morphology. The Journal of Rheumatology 34: 1925–1933
- Björk A 1963 Variations in the growth pattern of the human mandible: longitudinal radiographic study by the implant method. Journal of Dental Research 42 (Supplement): 400–411
- Björk A, Skieller V 1985 Contrasting mandibular growth and facial development in long face syndrome, juvenile rheumatoid polyarthritis, and mandibulofacial dysostosis. Journal of Craniofacial Genetics and Developmental Biology 1 (Supplement): 127–138
- Botto L D *et al.* 2003 A population-based study of the 22q11.2 deletion: phenotype, incidence, and contribution to major birth defects in the population. Pediatrics 112: 101–107
- Boutros S, Shetye P R, Ghali S, Carter C R, McCarthy J G, Grayson B H 2007 Morphology and growth of the mandible in Crouzon, Apert, and Pfeiffer syndromes. Journal of Craniofacial Surgery 18: 146–150
- Bryndahl F, Eriksson L, Legrell P E, Isberg A 2006 Bilateral TMJ disk displacement induces mandibular retrognathia. Journal of Dental Research 85: 1118–1123
- Buonaccorsi S, Leonardi A, Covelli E, Indrizzi E, Perdicchi A, Fini G 2005 Parry-Romberg syndrome. Journal of Craniofacial Surgery 16: 1132–1135
- Caccamese Jr J F, Costello B J, Mooney M P 2006 Novel deformity of the mandible in oculo-auriculo-vertebral spectrum: case report and literature review. Journal of Oral and Maxillofacial Surgery 64: 1278–1282
- Chan W L, Carolan M G, Fernandes V B, Abbati D P 2000 Planar versus SPET imaging in the assessment of condylar growth. Nuclear Medicine Communications 21: 285–290
- Defraia E, Marinelli A, Antonini A, Giuntini V 2005 Abnormal mandibular growth after craniovertebral surgery in Morquio syndrome type A. The Angle Orthodontist 75: 461–464
- Deleurant Y, Zimmermann A, Peltomäki T 2008 Hemimandibular elongation: treatment and long-term follow-up. Orthodontics & Craniofacial Research 11: 172–179
- Diewert V M 1982 Contributions of differential growth of cartilages to changes in craniofacial morphology. In: Dixon A D, Sarnat B G (eds). Factors and mechanisms influencing bone growth Alan R. Liss, Inc., New York, pp. 229–242
- Diewert V M 1985 Development of human craniofacial morphology during the late embryonic and early fetal periods. American Journal of Orthodontics 88: 64–76
- Dixon J et al. 2006 Tcof1/Treacle is required for neural crest cell formation and proliferation deficiencies that cause craniofacial abnormalities. Proceedings of the National Academy of Sciences of the United States of America 103: 13403–13408
- Eriksen J, Hermann N V, Darvann T A, Kreiborg S 2006 Early postnatal development of the mandible in children with isolated cleft palate and children with nonsyndromic Robin sequence. The Cleft Palate-Craniofacial Journal 43: 160–167
- Figueroa A A, Glupker T J, Fitz M G, BeGole E A 1991 Mandible, tongue, and airway in Pierre Robin sequence: a longitudinal cephalometric study. The Cleft Palate-Craniofacial Journal 28: 425–434
- Georgitsi M *et al.* 2007 Molecular diagnosis of pituitary adenoma predisposition caused by aryl hydrocarbon receptor-interacting protein gene mutations. Proceedings of the National Academy of Sciences of the United States of America 104: 4101–4105

- Gepstein A, Arbel G, Blumenfeld I, Peled M, Livne E 2003 Association of metalloproteinases, tissue inhibitors of matrix metalloproteinases, and proteoglycans with development, aging, and osteoarthritis processes in mouse temporomandibular joint. Histochemistry and Cell Biology 120: 23–32
- Goldring S R 2003 Inflammatory mediators as essential elements in bone remodeling. Calcified Tissue International 73: 97–100
- Gorlin R J, Cohen Jr M M, Hennekam R C M 2001 Syndromes of the head and neck, 4th edn. Oxford University Press, Oxford
- Halonen K, Hukki J, Arte S, Hurmerinta K 2006 Craniofacial structures and dental development in three patients with Nager syndrome. Journal of Craniofacial Surgery 17: 1180–1187
- Hanna V E *et al.* 1996 Effects of systemic onset juvenile rheumatoid arthritis on facial morphology and temporomandibular joint form and function. The Journal of Rheumatology 23: 155–158
- Heikinheimo K, Salmi K, Myllärniemi S, Kirveskari P 1989 Symptoms of craniomandibular disorder in a sample of Finnish adolescents at the ages of 12 and 15 years. European Journal of Orthodontics 11: 325–332
- Helenius L M et al. 2006 Clinical, radiographic and MRI findings of the temporomandibular joint in patients with different rheumatic diseases. International Journal of Oral and Maxillofacial Surgery 35: 983–989
- Henrikson T, Nilner M 2000 Temporomandibular disorders and the need for stomatognathic treatment in orthodontically treated and untreated girls. European Journal of Orthodontics 22: 283–292
- Henrikson T, Nilner M, Kurol J 2000 Signs of temporomandibular disorders in girls receiving orthodontic treatment. A prospective and longitudinal comparison with untreated Class II malocclusions and normal occlusion subjects. European Journal of Orthodontics 22: 271–281
- Hermann N V, Darvann T A, Jensen B L, Dahl E, Bolund S, Kreiborg S 2004 Early craniofacial morphology and growth in children with bilateral complete cleft lip and palate. The Cleft Palate-Craniofacial Journal 41: 424–438
- Ilowite N T 2002 Current treatment of juvenile rheumatoid arthritis. Pediatrics 109: 109–115
- Isacsson G, Carlson D S, McNamara Jr J A, Isberg A M 1993 Effect of maxillomandibular fixation on condylar growth in juvenile *Macaca mulatta*: a cephalometric and histologic study. Scandinavian Journal of Dental Research 101: 103–109
- Jacobsen P U, Lund K 1972 Unilateral overgrowth and remodeling processes after fracture of the mandibular condyle. A longitudinal radiographic study. Scandinavian Journal of Dental Research 80: 68–74
- Jakobsen L P et al. 2007 Pierre Robin sequence may be caused by dysregulation of SOX9 and KCNJ2. Journal of Medical Genetics 44: 381–386
- Jank S *et al.* 2007 Sonographic investigation of the temporomandibular joint in patients with juvenile idiopathic arthritis: a pilot study. Arthritis and Rheumatism 57: 213–218
- Johnston M C, Bronsky P T 1995 Prenatal craniofacial development: new insights on normal and abnormal mechanisms. Critical Reviews in Oral Biology and Medicine 6: 368–422
- Kanyama M et al. 2000 Matrix metalloproteinases and tissue inhibitors of metalloproteinases in synovial fluids of patients with temporomandibular joint osteoarthritis. Journal of Orofacial Pain 14: 20–30
- Karhulahti T, Rönning O, Jämsä T 1990 Mandibular condyle lesions, jaw movements, and occlusal status in 15-year-old children with juvenile rheumatoid arthritis. Scandinavian Journal of Dental Research 98: 17–26
- Kellenberger M, von Arx T, Hardt N 1996 Results of follow-up of temporomandibular joint fractures in 30 children. Fortschritte der Kiefer- und Gesichts-Chirurgie 41: 138–142
- Kiliaridis S, Thilander B, Kjellberg H, Topouzelis N, Zafiriadis A 1999 Effect of low masticatory function on condylar growth: a morphometric study in the rat. American Journal of Orthodontics and Dentofacial Orthopedics 116: 121–125
- Kjellberg H 1998 Craniofacial growth in juvenile chronic arthritis. Acta Odontologica Scandinavica 56: 360–365

- Kotilainen J, Hölttä P, Mikkonen T, Arte S, Sipilä I, Pirinen S 1995 Craniofacial and dental characteristics of Silver-Russell syndrome. American Journal of Medical Genetics 56: 229–236
- Kreiborg S, Cohen Jr M M, Skovby F 1991 Craniofacial characteristics of Proteus syndrome: two modes of abnormal growth. Proceedings of the Finnish Dental Society 8: 183–188
- Küseler A, Pedersen T K, Gelineck J, Herlin T 2005 A 2 year followup study of enhanced magnetic resonance imaging and clinical examination of the temporomandibular joint in children with juvenile idiopathic arthritis. The Journal of Rheumatology 32: 162–169
- Küseler A, Pedersen T K, Herlin T, Gelineck J 1998 Contrast enhanced magnetic resonance imaging as a method to diagnose early inflammatory changes in the temporomandibular joint in children with juvenile chronic arthritis. The Journal of Rheumatology 25: 1406–1412
- Laitinen S H, Ranta R E 1992 Cephalometric measurements in patients with Pierre Robin syndrome and isolated cleft palate. Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery 26: 177–183
- Legrell P E, Isberg A 1998 Mandibular height asymmetry following experimentally induced temporomandibular joint disk displacement in rabbits. Oral Surgery, Oral Medicine, and Oral Pathology 86: 280–285
- Legrell P E, Isberg A 1999 Mandibular length and midline asymmetry after experimentally induced temporomandibular joint disk displacement in rabbits. American Journal of Orthodontics and Dentofacial Orthopedics 115: 247–253
- Legrell P E, Reibel J, Nylander K, Hörstedt P, Isberg A 1999 Temporomandibular joint condyle changes after surgically induced nonreducing disk displacement in rabbits: a macroscopic and microscopic study. Acta Odontologica Scandinavica 57: 290–300
- Luder H U 2001 Histology of condyles in mandibular growth anomalies. In: Obwegeser H L (ed). Mandibular growth anomalies: terminology, aetiology, diagnosis, treatment Springer-Verlag, Berlin, pp. 347–353
- Lund K 1974 Mandibular growth and remodelling processes after condylar fracture. A longitudinal roentgencephalometric study. Acta Odontologica Scandinavica 32 (Supplement): 643–117
- Malemud C J 2007 Growth hormone, VEGF and FGF: involvement in rheumatoid arthritis. Clinica Chimica Acta 375: 10–19
- Marchetti C, Cocchi R, Gentile L, Bianchi A 2000 Hemimandibular hyperplasia: treatment strategies. Journal of Craniofacial Surgery 11: 46–53
- Marianowski R, Martins Carvalho C, Potard G, Pondaven S, Jézéquel J-A 2003 Mandibular fractures in children: long term results. International Journal of Pediatric Otorhinolaryngology 67: 25–30
- Martini G, Bacciliero U, Tregnaghi A, Montesco M C, Zulian F 2001 Isolated temporomandibular synovitis as unique presentation of juvenile idiopathic arthritis. The Journal of Rheumatology 28: 1689–1692
- Mayne J G, Hatch G S 1969 Arthritis of the temporomandibular joint. Journal of the American Dental Association 79: 125–130
- Mericle P M et al. 1996 Effects of polyarticular and pauciarticular onset juvenile rheumatoid arthritis on facial and mandibular growth. The Journal of Rheumatology 23: 159–165
- Miyamoto K, Ishimaru J, Kurita K, Goss A N 2002 Synovial matrix metalloproteinase-2 in different stages of sheep temporomandibular joint osteoarthrosis. Journal of Oral and Maxillofacial Surgery 60: 66–72
- Mulliken J B, Kaban L B 1987 Analysis and treatment of hemifacial microsomia in childhood. Clinics in Plastic Surgery 1: 91–100
- Narinobou M, Takatsuka S, Nakagawa K, Kubota Y, Terai K, Yamamoto E 2000 Histological changes in the rabbit condyle following posterolateral disk perforation. Journal of Cranio-Maxillo-Facial Surgery 28: 345–351
- Obwegeser H L (ed.) 2001 Mandibular growth anomalies: terminology, aetiology, diagnosis, treatment. Springer-Verlag, Berlin
- Obwegeser H L, Makek M S 1986 Hemimandibular hyperplasia hemimandibular elongation. Journal of Maxillofacial Surgery 14: 183–208

- Packham E A, Brook J D 2003 T-box genes in human disorders. Human Molecular Genetics 12: R37–R44
- Paesani D, Salas E, Martinez A, Isberg A 1999 Prevalence of temporomandibular joint disk displacement in infants and young children. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics 87: 15–19
- Palmisani E *et al.* 2006 Correlation between juvenile idiopathic arthritis activity and damage measures in early, advanced, and longstanding disease. Arthritis and Rheumatism 55: 843–849
- Pedersen T K, Jensen J J, Melsen B, Herlin T 2001 Resorption of the temporomandibular condylar bone according to subtypes of juvenile chronic arthritis. The Journal of Rheumatology 28: 2109–2115
- Peltomäki T, Alvesalo L, Isotupa K 1989 Shape of the craniofacial complex in 45,X females: cephalometric study. Journal of Craniofacial Genetics and Developmental Biology 9: 331–338
- Perkiömäki M R, Kyrkanides S, Niinimaa A, Alvesalo L 2005 The relationship of distinct craniofacial features between Turner syndrome females and their parents. European Journal of Orthodontics 27: 48–52
- Petty R E *et al.* 1998 Revision of the proposed classification criteria for juvenile idiopathic arthritis: Durban, 1997. The Journal of Rheumatology 25: 1991–1994
- Pirttiniemi P, Kantomaa T, Sorsa T 2004 Effect of decreased loading on the metabolic activity of the mandibular condylar cartilage in the rat. European Journal of Orthodontics 26: 1–5
- Posnick J C, Ruiz R L 2000 Treacher Collins syndrome: current evaluation, treatment, and future directions. The Cleft Palate-Craniofacial Journal 37: 434
- Poswillo D 1973 The pathogenesis of the first and second branchial arch syndrome. Oral Surgery 35: 302–328
- Pripatnanont P, Vittayakittipong P, Markmanee U, Thongmak S, Yipintsoi T 2005 The use of SPECT to evaluate growth cessation of the mandible in unilateral condylar hyperplasia. International Journal of Oral and Maxillofacial Surgery 34: 364–368
- Ribeiro R F et al. 1997 The prevalence of disc displacement in symptomatic and asymptomatic volunteers aged 6 to 25 years. Journal of Orofacial Pain 11: 37–47
- Roberts F G, Pruzansky S, Aduss H 1975 An X-radiocephalometric study of mandibulofacial dysostosis in man. Archives of Oral Biology 20: 265–281
- Rongen-Westerlaken C et al. 1993 Effect of growth hormone treatment on craniofacial growth in Turner's syndrome. Acta Paediatrica 82: 364–368
- Rönning O, Väliaho M-L, Laaksonen A-L 1974 The involvement of the temporomandibular joint in juvenile rheumatoid arthritis. Scandinavian Journal of Rheumatology 3: 89–96
- Rubenstein L K, Campbell R L 1985 Acquired unilateral condylar hyperplasia and facial asymmetry: report of case. ASDC Journal of Dentistry for Children 52: 114–120
- Rushton M A 1944 Growth at the mandibular condyle in relation to some deformities. British Dental Journal 76: 57–68
- Rushton M A 1946 Unilateral hyperplasia of the mandibular condyle. Proceedings of the Royal Society of Medicine 39: 431–438
- Säilä H *et al.* 2004 HLA and susceptibility to juvenile idiopathic arthritis: a study of affected sibpairs in an isolated Finnish population. The Journal of Rheumatology 31: 2281–2285
- Saurenmann R K *et al.* 2007 Epidemiology of juvenile idiopathic arthritis in a multiethnic cohort: ethnicity as a risk factor. Arthritis and Rheumatism 56: 1974–1984
- Schubert J, Jahn H, Berginski M 2005 Experimental aspects of the pathogenesis of Robin sequence. Cleft Palate-Craniofacial Journal 42: 372–376
- Sen M 2005 Wnt signalling in rheumatoid arthritis. Rheumatology 44: 708–713
- Slootweg P J, Müller H 1986 Condylar hyperplasia. A clinico-pathological analysis of 22 cases. Journal of Maxillofacial Surgery 14: 209–214

- Stevenson D A, Bleyl S B, Maxwell T, Brothman A R, South S T 2007 Mandibulofacial dysostosis in a patient with a *de novo* 2;17 translocation that disrupts the *HOXD* gene cluster. American Journal of Medical Genetics 143A: 1053–1059
- Suri S, Ross R B, Tompson B D 2006 Mandibular morphology and growth with and without hypodontia in subjects with Pierre Robin sequence. American Journal of Orthodontics and Dentofacial Orthopedics 130: 37–46
- Svensson B, Larsson Å, Adell R 2001 The mandibular condyle in juvenile chronic arthritis patients with mandibular hypoplasia. A clinical and histological study. International Journal of Oral and Maxillofacial Surgery 30: 300–305
- Teber Ö A *et al.* 2004 Genotyping in 46 patients with tentative diagnosis of Treacher Collins syndrome revealed unexpected phenotypic variation. European Journal of Human Genetics 12: 879–890
- Tiilikainen P, Pirttiniemi P, Kainulainen T, Pernu H, Raustia A 2005 MMP-3 and -8 expression is found in the condylar surface of temporomandibular joints with internal derangement. Journal of Oral Pathology & Medicine 34: 39–45
- Twilt M, Arends L R, ten Cate R, van Suijlekom-Smit L W A 2007 Incidence of temporomandibular involvement in juvenile idiopathic arthritis. Scandinavian Journal of Rheumatology 36: 184–188
- Twilt M, Schulten A J M, Nicolaas P, Dülger A, van Suijlekom-Smit L W A 2006 Facioskeletal changes in children with juvenile idiopathic arthritis. Annals of the Rheumatic Diseases 65: 823–825

- Twilt M, Schulten A J M, Verschure F, Wisse L, Prahl-Andersen B, van Suijlekom-Smit L W A 2008 Long-term followup of temporomandibular joint involvement in juvenile idiopathic arthritis. Arthritis and Rheumatism 59: 546–552
- Van Erum R, Mulier M, Carels C, Verbeke G, de Zegher F 1997 Craniofacial growth in short children born small for gestational age: effect of growth hormone treatment. Journal of Dental Research 76: 1579–1586
- Westling L, Mohlin B, Bresin A 1998 Craniofacial manifestations in the Marfan syndrome: palatal dimensions and a comparative cephalometric analysis. Journal of Craniofacial Genetics and Developmental Biology 18: 211–218
- Yang L, Thornton S, Grom A A 2002 Interleukin-15 inhibits sodium nitroprusside-induced apoptosis of synovial fibroblasts and vascular endothelial cells. Arthritis and Rheumatism 46: 3010–3014
- Yoshida K et al. 2006 Expression of matrix metalloproteinases and aggrecanase in the synovial fluids of patients with symptomatic temporomandibular disorders. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics 102: 22–27
- Yu D, Tiilikainen P, Raustia A, Pirttiniemi P 2007 Dietary loading and aggrecanase-1/TIMP-3 expression in rat mandibular condylar cartilage. Journal of Orofacial Pain 21: 232–238
- Zhou J *et al.* 2005 The growth hormone receptor gene is associated with mandibular height in a Chinese population. Journal of Dental Research 84: 1052–1056

Copyright of European Journal of Orthodontics is the property of Oxford University Press / UK and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.