

# Patient and parent motivation for orthodontic treatment—a questionnaire study

Barbara Wędrychowska-Szulc and Maria Syryńska

Department of Orthodontics, Pomeranian Medical University, Szczecin, Poland

**SUMMARY** The aims of this investigation were to examine patients' and parents/guardians' motivation for seeking orthodontic treatment and to determine the influence of age and gender in this process. The sample comprised 674 subjects (365 girls and 309 boys) aged 7–18 years and 674 of their parents/guardians, as well as 86 adult patients aged 19–42 years (57 females and 29 males). Similar questionnaires, designed to assess motivation for orthodontic treatment, were completed by all subjects. Statistical analysis was undertaken using Pearson's chi-square test.

In the patient groups, a desire to improve aesthetics was the main motivational factor for undergoing treatment. With increasing age, 13 per cent more girls ( $P=0.039$ ) were more aware of their malocclusion. The influence of their surroundings on the uptake of treatment decreased with increasing age. Less than 5 per cent of the examined subjects started treatment because other children made fun of them, and only 3 per cent of older patients were motivated by future improvements in health. Between 63 and 67 per cent of parents reported pressurizing their children to seek orthodontic treatment in order to avoid possible future claims of neglect. No statistically significant dependence on gender or age of the children was found.

Improvement in dental aesthetics was the principle motivational factor for the children (29–48 per cent), their parents/guardians (54 per cent), and adult patients (55 per cent) seeking orthodontic treatment.

## Introduction

Facial aesthetics seems to be a significant determinant of self and social perceptions (Bos *et al.*, 2003). Optimal facial appearance is not only seen as being more attractive but also more socially accepted by peers, teachers, employers, and others (Phillips *et al.*, 1998; Tung and Kiyak, 1998; Trulsson *et al.*, 2002). Smiling is still considered to be one of the most effective methods of influencing people (Kokich *et al.*, 1999, 2006).

Some individuals with dentofacial disharmonies seeking treatment experience a level of psychological distress that may warrant psychological/psychiatric intervention (Phillips *et al.*, 1998). However, the most common reasons for seeking professional help by orthognathic patients were due problems in biting and chewing, temporomandibular disorders, and headache (Nurminen *et al.*, 1999).

Correlations between satisfaction with dental and facial appearance and expectations of orthodontic treatment appear to be age, but not gender, related (Bos *et al.*, 2003).

Although many different factors motivate patients in their decision to undergo orthodontic treatment, many adolescents are not fully conscious of such external influences, such as social norms, and the beauty culture in their reference group and in society (Trulsson *et al.*, 2002). Their opinion, as a group, was that they had made an independent decision to undergo orthodontic treatment. Thus, it is essential to understand subjective motives for undergoing orthodontic therapy and thereby set realistic treatment goals (Trulsson *et al.*, 2002).

According to Sheats *et al.* (1998), eighth graders with no history of orthodontic treatment were generally satisfied with the appearance of their teeth and perceived less treatment need than clinicians. Tung and Kiyak (1998) reported that although Caucasian and minority children were similar in their self-ratings and expectations from orthodontics, the former were more critical in their aesthetic judgement.

However, it has been reported that up to 75 per cent of adult patients are dissatisfied with their dental aesthetics and that this is their prime motive for seeking treatment (Birkeland *et al.*, 1999; Riedmann *et al.*, 1999). Bergström *et al.* (1998) found that three out of four individuals considered orthodontic treatment important, even when irregularities of the teeth were minor. A majority of the individuals stated that they would have been able to wear visible fixed appliances if needed, even in adulthood.

Females demonstrated less satisfaction than males with the appearance of their dentition and were more likely to perceive a treatment need. Asians and females had higher Index of Orthodontic Treatment Need Dental Health Component scores, but a better aesthetic appearance than Caucasians and males (Sheats *et al.*, 1998; Mandall *et al.*, 1999).

Adult patients, including those who had been previously treated (during childhood or adolescence), reported satisfaction with their decision either to choose to undergo orthodontic treatment or to decline it (Bergström *et al.*, 1998). They were also satisfied with their dental appearance.

Dental professionals were considered to have had the greatest influence on this decision (Bergström *et al.*, 1998).

Kokich *et al.* (1999) reported that even if both the orthodontist and the general dentist detect specific dental aesthetic discrepancies, which may help the dental professional in their treatment recommendations, orthodontists were more critical than dentists and laypeople. The desire for treatment may be guided to a considerable extent by the orthodontist (Stenvik *et al.*, 1996; Bergström *et al.*, 1998; Birkeland *et al.*, 1999; Espeland and Stenvik, 1999). It has also been noted that patients treated by specialists were more satisfied than individuals treated by general practitioners (Bergström *et al.*, 1998).

It is of note that individuals with malocclusion and treatment need, but who refused treatment, were, in general, more unhappy with their dental appearance (Bergström *et al.*, 1998). More than half regretted their decision. Thus, the perceptions of orthodontic treatment need are multifactorial and that they are also influenced by factors other than health measures of normative orthodontic treatment need and perception of aesthetics (Hamdan, 2004).

Kerosuo *et al.* (2002) stated that patients from rural areas expressed a treatment need less often than those from more urban areas. Orthodontic treatment is often undertaken on the assumption that an improved dental appearance will benefit a patient by increasing his/her social acceptance and self-concept (Dann *et al.*, 1995). On the other hand, an objectively favourable occlusal treatment result may lead to patient dissatisfaction, particularly when there is a difference between the patient's motivation and expectations (Bos *et al.*, 2003).

Thus, it is important to determine prior to treatment both the patient's and their parents/guardians' motivations for and expectations of treatment. A well-designed questionnaire may prove to be a valuable tool (Arnett and Worley, 1999).

The aims of this study were to examine and compare patients' and their parents/guardians' reasons for seeking orthodontic treatment, and to explore and analyse the conformity for orthodontic treatment in relation to age and gender, and to examine the influence of specialists and parents in motivating patients.

## Subjects and methods

This questionnaire-based study was carried out between June and December 2006 in north-west Poland among 674 (365 girls and 309 boys) children aged 7–18 years and 674 of their parents/guardians. In addition, 86 adult patients aged 19–42 years (57 females and 29 males) were evaluated using a similar questionnaire. The patients were recruited from the Department of Orthodontics of the Pomeranian Medical University and four private orthodontic practices in Szczecin and Kołobrzeg. All subjects were Caucasian.

Seven hundred and sixty patients (674 children and 86 adult patients) who applied for orthodontic treatment completed a questionnaire, which contained nine items on motivation for starting treatment (Appendix 1). The 674 parents/guardians completed a similar questionnaire, containing 10 items (Appendix 2). The subjects could, if applicable, select more than one answer to each question.

To determine any influence of the patient's age and/or gender, Pearson's chi-square independent test for contingency tables was used. The calculations were made using the Statistica program for Windows version 6.0 (StatSoft, Tulsa, Oklahoma, USA).

## Results

### Subject demographics

The age and gender distribution of the study sample are shown in Table 1. With the exception of the 7- to 9-year-olds, more females than males sought orthodontic treatment. The 10- to 12-year-old group demonstrated the greatest percentage of males and females presenting for treatment.

### Patients' reasons for seeking orthodontic treatment

The majority of patients selected aesthetics as the main reason for their decision to undergo orthodontic treatment (Table 2; Appendix 1: answers a = 64 per cent and b = 42 per cent). The intention to improve self-image increased with age ( $P < 0.001$ ) and gender ( $P = 0.029$ ). In addition to increasing age, a significantly higher percentage of females

**Table 1** Age and gender of the respondents.

Patient's age (years)	Patient's gender, <i>n</i> (%)			Parents/guardians, <i>n</i> (%)
	Females	Males	Total	
7–9	72 (17)	73 (22)	145 (19)	145 (21.5)
10–12	164 (39)	148 (44)	312 (41)	312 (46)
13–15	89 (21)	58 (17)	147 (19.5)	147 (22)
16–18	40 (9.5)	30 (8.5)	70 (9)	70 (10.5)
More than 18	57 (13.5)	29 (8.5)	86 (11.5)	—
Total patients	422	338	760	
Total parents/guardians	365	309		674

**Table 2** The reasons for coming to the orthodontist—responses of the patients.

The selected answer	<i>n</i> (%) of the selected answers														
	Patient's age (years) and gender (M—boys/males, F—girls/females)														
	7–9			10–12			13–15			More than 16			Total		
	F	M	F + M	F	M	F + M	F	M	F + M	F	M	F + M	F	M	F + M
a **( <i>P</i> =0.039)	41 (57)	39 (53)	80 (55)	111 (68)	93 (63)	204 (65)	64 (72)	36 (62)	100 (68)	68 (70)	35 (59)	103 (66)	284 (67)	203 (60)	487 (64)
b *( <i>P</i> <0.001)**( <i>P</i> =0.029)	23 (32)	20 (27)	43 (29)	69 (42)	52 (35)	121 (38)	45 (50)	26 (45)	71 (48)	56 (57)	30 (50)	86 (55)	193 (45)	128 (38)	321 (42)
c *( <i>P</i> =0.007)	35 (49)	34 (47)	69 (48)	64 (39)	56 (38)	120 (38)	32 (35)	26 (45)	58 (39)	25 (26)	19 (32)	44 (28)	156 (57)	135 (40)	291 (38)
d	4 (5)	6 (8)	10 (7)	10 (6)	7 (5)	17 (5)	1 (1)	2 (3)	3 (2)	1 (1)	3 (5)	4 (3)	16 (6)	18 (5)	34 (4)
e *( <i>P</i> <0.001)**( <i>P</i> =0.026)	31 (43)	34 (47)	65 (45)	80 (49)	79 (53)	159 (51)	37 (42)	31 (53)	68 (46)	19 (20)	17 (29)	36 (23)	36 (23)	161 (48)	328 (43)
f	4 (5)	8 (11)	12 (8)	6 (4)	10 (7)	16 (5)	2 (2)	1 (2)	3 (2)	11 (11)	5 (8)	16 (10)	23 (5)	23 (7)	46 (6)
g *( <i>P</i> =0.000001)	0 (0)	0 (0)	0 (0)	1 (0.6)	2 (1)	3 (1)	0 (0)	0 (0)	0 (0)	12 (12)	8 (14)	20 (13)	13 (3)	10 (3)	23 (3)
h	0 (0)	0 (0)	0 (0)	1 (0.6)	0 (0)	1 (0.3)	0 (0)	0 (0)	0 (0)	2 (2)	0 (0)	2 (1)	3 (0.7)	0 (0)	3 (0.4)
i *( <i>P</i> =0.0001)	2 (3)	3 (4)	5 (3)	5 (3)	7 (5)	12 (4)	5 (6)	5 (9)	19 (7)	14 (14)	8 (14)	22 (14)	26 (6)	22 (7)	48 (6)

Statistically significant depending on the patient's \*age and \*\*gender.

**Table 3** The reasons for coming to the orthodontist—responses of the parents/guardians.

The selected answer	<i>n</i> (%) of the selected (by parents/guardians) answers														
	Patient's age (years) and gender (M—boys/males, F—girls/females)														
	7–9			10–12			13–15			More than 16			Total		
	F	M	F + M	F	M	F + M	F	M	F + M	F	M	F + M	F	M	F + M
a	51 (71)	58 (79)	109 (75)	133 (81)	118 (80)	251 (80.5)	64 (72)	45 (78)	109 (74)	30 (75)	22 (73)	52 (74)	278 (76)	243 (79)	521 (77)
b	40 (55)	43 (59)	83 (57)	95 (58)	76 (51)	171 (55)	46 (52)	27 (47)	73 (50)	21 (52)	14 (47)	35 (50)	202 (55)	160 (52)	362 (54)
c*( <i>P</i> <0.001)	35 (49)	44 (60)	79 (54)	75 (46)	63 (43)	138 (44)	34 (38)	24 (41)	58 (39)	9 (22.5)	9 (30)	18 (26)	153 (42)	140 (45)	293 (43)
d	3 (4)	3 (4)	6 (4)	10 (6)	5 (3)	15 (5)	0 (0)	1 (2)	1 (0.7)	1 (2.5)	3 (1)	4 (6)	14 (4)	12 (4)	26 (4)
e	42 (58)	50 (68)	92 (63)	105 (64)	97 (66)	202 (65)	56 (63)	37 (64)	93 (63)	26 (65)	21 (70)	47 (67)	229 (63)	205 (66)	434 (64)
f	6 (8)	11 (15)	17 (12)	16 (10)	15 (10)	31 (10)	5 (6)	4 (7)	9 (6)	1 (3)	2 (7)	3 (4)	28 (8)	32 (10)	60 (9)
g*( <i>P</i> =0.049)	0 (0)	0 (0)	0 (0)	2 (1)	0 (0)	2 (0.6)	0 (0)	0 (0)	0 (0)	0 (0)	2 (7)	2 (3)	0 (0.5)	2 (0.6)	4 (0.6)
h	0 (0)	1 (1)	1 (0.6)	4 (2)	0 (0)	4 (1)	1 (1)	0 (0)	1 (0.7)	0 (0)	2 (7)	2 (3)	5 (1.4)	3 (1)	8 (1)
i	13 (18)	13 (18)	26 (18)	31 (19)	27 (18)	58 (18.5)	22 (25)	13 (22)	35 (24)	6 (15)	5 (17)	11 (16)	72 (20)	58 (19)	130 (19)
j	2 (3)	3 (4)	5 (3.5)	9 (5)	13 (9)	22 (7)	5 (6)	5 (9)	10 (7)	1 (2.5)	3 (10)	4 (6)	17 (5)	24 (8)	41 (6)

\*Statistically significant depending on the patient's age.

( $P=0.039$ ) expressed dissatisfaction with their dental appearance, whereas the number of subjects who wanted to improve their appearance increased more slowly among males ( $P=0.029$ ). On the contrary, the percentage of patients whose decision to undergo the treatment might be guided by dental professionals decreased by 20 per cent with increasing age ( $P=0.007$ ).

The influence of parents/guardians on the patient decision to undergo orthodontic treatment (Appendix 1: answer e) decreased with patient age ( $P<0.001$ ). Less than 5 per cent of patients in all age groups sought orthodontic treatment as a result of the influence of their peers. Health did not appear to be a key motivational factor in seeking treatment among adults [Appendix 1: answers f ( $P=0.018$ ) and g ( $P<0.001$ )].

#### *Parent/guardian reasons for their child seeking orthodontic treatment*

The percentage of parents/guardians that selected irregular teeth and a wish for their child to look 'pretty' as reasons for seeking orthodontic treatment was 77 and 54, respectively (Table 3). No statistically significant differences relating to age or gender of the child were found.

Along with increasing patient age, the percentage of the parents/guardians whose children were referred for treatment by other clinicians decreased ( $P<0.001$ ).

Sixty-four per cent of parents/guardians reported that they sought treatment for their child out of fear of any possible future blame from the child that their parents/guardians neglected their duty to make sure they underwent orthodontic treatment.

No relationship was found between the age or gender of the patients and conformity of the motivations for the treatment stated by the patient and/or parents/guardians.

With reference to the question h (Appendix 2), only 16 parents/guardians reported any influence of the school in their decision to seek treatment for their child.

## **Discussion**

### *Patient's reasons for seeking orthodontic treatment*

Taking into consideration the relationship between a patient's age and their motivation for seeking orthodontic treatment, the present study found that the number of patients dissatisfied with the appearance of their teeth increased with age. Similar results have been reported by Birkeland *et al.* (1999) and Bos *et al.* (2003). Espeland and Stenvik (1991) noted that among young adults who were characterized as having an almost ideal occlusion or only minor deviations, almost all (98 per cent) expressed satisfaction with their actual occlusal status.

In relation to gender (while assessing themselves subjectively), females in the present study demonstrated more concern for appearance (56 per cent) than males (44 per cent), which corresponds with the findings of previous

studies (Sheats *et al.*, 1998; Tung and Kiyak, 1998; Mandall *et al.*, 1999; Nurminen *et al.*, 1999; Riedmann *et al.*, 1999; Kerosuo *et al.*, 2002; Bos *et al.*, 2003; Hamdan, 2004).

The patient's decision in relation to starting orthodontic treatment was based on the recommendation of either the dentist or their parents/guardians. The influence of these advisors decreased with increasing patient age. This corresponds to the findings of others (Dann *et al.*, 1995; Birkeland *et al.*, 1996; Bergström *et al.*, 1998).

The influence of social stereotypes based on facial appearance as a major factor in total life adjustment is one reason for seeking orthodontic treatment. Dental anomalies have been reported to be the cause of teasing and 'general playground harassment' among children and are associated with lowered social attractiveness (Phillips *et al.*, 1998; Hamdan, 2004). In the present study, less than 5 per cent of patients in all age groups sought orthodontic treatment as a result of the influence of their peers. In addition, negative attitudes of school authorities and employers were not found to be a significant factor.

Problems with biting and chewing, temporomandibular disorders, and headache have been reported as reasons for seeking professional advice (Phillips *et al.*, 1998; Nurminen *et al.*, 1999). In the present study, health did not appear to be a key motivational factor for the adults to seek treatment. However, according to Nurminen *et al.* (1999) and Trulsson *et al.* (2002), functional reasons were reported by 47–68 per cent of patients who had undergone orthognathic surgery.

### *Parent/guardian reasons for their child seeking orthodontic treatment*

Previous studies have shown that not only parents' motivation, especially that of the mother, is the most important factor in initiating orthodontic treatment, but also that treatment is considered to be more important by the parents than by the child (Birkeland *et al.*, 1996; Bergström *et al.*, 1998; Daniels and Richmond, 2000; Kerosuo *et al.*, 2002; Hamdan, 2004; Zhang *et al.*, 2007). Thus, parents appear to make the final decision about treatment, although they may have different motives to their children (Birkeland *et al.*, 1996; Hamdan, 2004; Hamdan *et al.*, 2007). In line with the findings of Birkeland *et al.* (1996) and Tung and Kiyak (1998), the number of parents/guardians dissatisfied with the appearance of their children's teeth was 75 per cent, of which 54 per cent wanted their children 'to look pretty'. The current study also found that the final decision of parents/guardians for their child to undergo orthodontic treatment was not solely their own but also influenced by the referring dentists, other physicians, and speech therapists. However, this influence decreased significantly ( $P<0.001$ ) with increasing age of the children. Similar findings were noted by Trulsson *et al.* (2002) and Bergström *et al.* (1998). Interestingly, response to the question 'I want my child not to blame me in the future that I have neglected his/her treatment for his/her dental trouble'



revealed a positive response of 64 per cent, which was not influenced by either the child's age or the gender.

The results of the current study are in agreement with the findings of Dann *et al.* (1995) that dentofacial appearance undoubtedly plays an important part in establishing the overall attractiveness of individuals.

### Limitations of the study

There is evidence that socio-economic background may also play a role in the self-perception of malocclusion, with individuals in a higher social class considered to be more critical of their dental aesthetics (Lewit and Virolainen, 1968; Al-Sarheed *et al.*, 2003; Reichmuth *et al.*, 2005). The current study did not take account of the influence of socio-economic factors.

### Conclusions

1. The principle motivational factor in children and adults seeking orthodontic treatment was the desire to improve aesthetics.
2. From the parents/guardians perspective, the most important reason for seeking orthodontic treatment for their child was also the improvement of dental aesthetics.
3. The percentage of patients whose decision to undergo treatment might be guided by dental or other professionals, or their parents/guardians decreased with increasing patient age.

### Address for correspondence

Barbara Wędrychowska-Szulc  
Department of Orthodontics  
Pomeranian Medical University  
Al. Powstańców Wielkopolskich 72  
PL – 70-111 Szczecin  
Poland  
E-mail: bws@sci.pam.szczecin.pl

### Acknowledgements

We would like to express our special gratitude to Ewa Młodzińska, Hanna Morzycka, Małgorzata Piela, and Bohdan Krawczyk for their considerable help in data collection.

### References

- Al-Sarheed M, Bedi R, Hunt N P 2003 Orthodontic treatment need and self-perception of 11–16-year-old Saudi Arabian children with a sensory impairment attending special schools. *Journal of Orthodontics* 30: 39–44
- Arnett G W, Worley C M Jr 1999 The treatment motivation survey: defining patient motivation for treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 115: 233–238
- Bergström K, Halling A, Wilde B 1998 Orthodontic care from the patients' perspective: perceptions of 27-year-olds. *European Journal of Orthodontics* 20: 319–329
- Birkeland K, Bøe O E, Wisth P J 1996 Orthodontic concern among 11-year-old children and their parents compared with orthodontic treatment need assessed by Index of Orthodontic Treatment Need. *American Journal of Orthodontics and Dentofacial Orthopedics* 110: 197–205
- Birkeland K, Kattø A, Løvgreen S, Bøe O E, Wisth P J 1999 Factors influencing the decision about orthodontic treatment. A longitudinal study among 11- and 15-year-olds and their parents. *Journal of Orofacial Orthopedics* 60: 292–307
- Bos A, Hoogstraten J, Prah Andersen B 2003 Expectations of treatment and satisfaction with dentofacial appearance in orthodontic patients. *American Journal of Orthodontics and Dentofacial Orthopedics* 123: 127–132
- Daniels C, Richmond S 2000 The development of the Index of Complexity, Outcome and Need (ICON). *Journal of Orthodontics* 27: 149–162
- Dann C, Phillips C, Broder H L, Tulloch C 1995 Self-concept, Class II malocclusion, and early treatment. *The Angle Orthodontist* 65: 411–416
- Espeland L, Stenvik A 1999 Residual need in orthodontically untreated 16–20-year-olds from areas with different treatment rates. *European Journal of Orthodontics* 21: 523–531
- Espeland L V, Stenvik A 1991 Perception of personal dental appearance in young adults: relationship between occlusion, awareness, and satisfaction. *American Journal of Orthodontics and Dentofacial Orthopedics* 100: 234–241
- Hamdan A M 2004 The relationship between patient, parent and clinician perceived need and normative orthodontic treatment need. *European Journal of Orthodontics* 26: 265–271
- Hamdan A M, Al-Omari I K, Al-Bitar Z B 2007 Ranking dental aesthetics and thresholds of treatment need: a comparison between patients, parents, and dentists. *European Journal of Orthodontics* 29: 366–371
- Kerosuo H, Abdulkarim E, Kerosuo E 2002 Subjective need and orthodontic treatment experience in a middle east country providing free orthodontic services: a questionnaire survey. *The Angle Orthodontist* 72: 565–570
- Kokich V O, Kiyak H A, Shapiro P A 1999 Comparing the perception of dentists and lay people to altered dental esthetics. *Journal of Esthetic Dentistry* 11: 311–324
- Kokich V O, Kokich V G, Kiyak H A 2006 Perceptions of dental professionals and laypersons to altered dental esthetics: asymmetric and symmetric situations. *American Journal of Orthodontics and Dentofacial Orthopedics* 130: 141–151
- Lewit D W, Virolainen K 1968 Conformity and independence in adolescents' motivation for orthodontic treatment. *Child Development* 39: 1189–1200
- Mandall N A, McCord J F, Blinkhorn A S, Worthington H V, O'Brien K D 1999 Perceived aesthetic impact of malocclusion and oral self-perceptions in 14–15-year-old Asian and Caucasian children in Greater Manchester. *European Journal of Orthodontics* 21: 175–183
- Nurminen L, Pietilä T, Vinkka-Puhakka H 1999 Motivations for and satisfaction with orthodontic-surgical treatment: a retrospective study of 28 patients. *European Journal of Orthodontics* 21: 79–87
- Phillips C, Bennett M E, Broder H L 1998 Dentofacial disharmony: psychological status of patients seeking treatment consultation. *The Angle Orthodontist* 68: 547–556
- Reichmuth M, Greene K A, Orsini M G, Cisneros G J, King G, Kiyak H A 2005 Occlusal perceptions of children seeking orthodontic treatment: impact of ethnicity and socioeconomic status. *American Journal of Orthodontics and Dentofacial Orthopedics* 128: 575–582
- Riedmann T, Georg T, Berg R 1999 Adult patients' view of orthodontic treatment outcome compared to professional assessments. *Journal of Orofacial Orthopedics* 60: 308–320
- Sheats R D, McGorray S P, Keeling S D, Wheeler T T, King G J 1998 Occlusal traits and perception of orthodontic need in eighth grade students. *The Angle Orthodontist* 68: 107–114
- Stenvik A, Espeland L, Berset G P, Eriksen H M, Zachrisson B U 1996 Need and desire for orthodontic (re-)treatment in 35-year-old Norwegians. *Journal of Orofacial Orthopedics* 57: 334–342

- Trulsson U, Strandmark M, Modlin B, Berggren U 2002 A qualitative study of teenagers' decisions to undergo orthodontic treatment with fixed appliance. *Journal of Orthodontics* 29: 197–204
- Tung A W, Kiyak H A 1998 Psychological influences on the timing of orthodontic treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 113: 29–39
- Zhang M, McGrath C, Hägg U 2007 Who knows more about the impact of malocclusion on children's quality of life, mothers or fathers? *European Journal of Orthodontics* 29: 180–185

## Appendix 1

### Questionnaire

Patient's age. . . . . Patient's gender: male/female  
[the correct is to be underlined] Child's father/mother/  
guardian  
[the correct is to be underlined]  
The following questions to be answered:

Patient:

1. The reason for which the patient has applied to the orthodontist:
  - a. I can see that I have 'irregular teeth' [malocclusion]
  - b. I want to correct my appearance
  - c. I have been referred by: my dentist—another physician—speech therapist [the correct is to be underlined]
  - d. The other children laugh at my 'irregular teeth'
  - e. My parents wanted me to be treated
  - f. I have some difficulties while: speaking—eating [the correct is to be underlined]

- g. I feel some pain/clicking around my ears (at the temporomandibular joint) [the correct is to be underlined]
- h. I have been ordered by employer – school authorities [the correct is to be underlined]
- i. Other reasons . . .

## Appendix 2

### Child's father/mother/guardian

2. The reason of coming with the child to the orthodontist:
  - a. I have realized that my child has got 'irregular teeth' [malocclusion]
  - b. I want my child to look pretty
  - c. Child has been sent for treatment by: his dentist—another physician—speech therapist [the correct is to be underlined]
  - d. Other children are laughing at my child's 'irregular teeth'
  - e. I do not want my child to blame me in the future that I have neglected his/her treatment for his/her dental trouble
  - f. Child has some difficulties while: speaking – biting solid food [the correct is to be underlined]
  - g. My child has complained of some pain around his/her ears (at the temporomandibular joint)
  - h. Such are the school requirements
  - i. I think that to correct my child's malocclusion could help him/her in the future to find a better job
  - j. Other reasons . . .

Copyright of European Journal of Orthodontics is the property of Oxford University Press / UK and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.