Expressions of RANKL/RANK and M-CSF/*c-fms* in root resorption lacunae in rat molar by heavy orthodontic force

Yoko Nakano, Masaru Yamaguchi, Shoji Fujita, Masaki Asano, Kayo Saito and Kazutaka Kasai

Department of Orthodontics, Nihon University School of Dentistry at Matsudo, Japan

Correspondence to: Masaru Yamaguchi, Department of Orthodontics, Nihon University School of Dentistry at Matsudo, 2-870-1 Sakaecho-Nishi, Matsudo, Chiba 271-8587, Japan. E-mail: yamaguchi.masaru@nihon-u.ac.jp

SUMMARY The differentiation and functions of osteoclasts are regulated by receptor activator of nuclear factor- κ B (RANK)/receptor activator of nuclear factor- κ B ligand (RANKL) system that stimulates osteoclasts formation. Macrophage colony-stimulating factor (M-CSF) is also essential for osteoclastogenesis. A recent immunocytochemical study reported that RANKL/RANK and M-CSF/*c-fms* were localized in the periodontal ligament of rat molars during experimental orthodontic tooth movement. The present study focused on the expressions of RANKL/RANK and M-CSF/*c-fms* in root resorption area during experimental tooth movement in rats.

Forty 6-week-old male Wistar rats were subjected to an orthodontic force of 10 or 50 g with a closed coil spring (wire size: 0.005 inch, diameter: 1/12 inch) ligated to the maxillary first molar cleat by a 0.008 inch stainless steel ligature wire to induce a mesial tipping movement of the upper first molars. Experimental tooth movement was undertaken for 10 days. Each sample was sliced into 6 μ m continuous sections in a horizontal direction and prepared for haematoxylin and eosin (H and E) and immunohistochemistry staining for tartrate-resistant acid phosphatase (TRAP), RANK, RANKL M-CSF, and *c-fms* in root resorption area. Statistical analysis was carried out using a Mann–Whitney *U*-test with a significance level of *P* < 0.01.

On days 7 and 10, immunoreactivity for RANKL/RANK and M-CSF/*c-fms* was detected in odontoclasts with an orthodontic force of 50 g, but not 10 g. Therefore, RANKL/RANK and M-CSF/*c-fms* systems may be involved in the process of root resorption by heavy orthodontic force.

Introduction

Two cytokines are essential for osteoclastogenesis, the receptor activator of nuclear factor-kB ligand (RANKL; Lacey et al., 1998; Yasuda et al., 1998) and macrophage colony-stimulating factor (M-CSF), also called colonystimulating factor-1, with the former being the key osteoclastogenic cytokine and the latter contributing to proliferation, survival, and differentiation of early precursors (Udagawa et al., 1999). According to Soedarsono et al. (2006), recent advances in bone cell biology have allowed elucidation of the crucial roles of RANK/RANKL/osteoprotegerin (OPG) system in osteoclast differentiation and function. RANKL is a cytokine that belongs to the tumour necrosis factor (TNF) family and is essential for the induction of osteoclastogenesis. Osteoblasts and bone marrow stromal cells produce this cytokine, and its signals are transduced by the specific receptor RANK, which is localized on the cell surface of osteoclast progenitors. The RANKL/RANK system has been suggested to play an integral role in osteoclast activation during orthodontic tooth movement. Shiotani et al. (2001) observed RANKL in osteoblasts, osteocytes, fibroblasts, and osteoclasts during the application of orthodontic forces. The RANKL/RANK system may

regulate the natural process of root resorption in exfoliated primary teeth (Lossdörfer *et al.*, 2002).

M-CSF has also been identified as a molecule that mediates the survival and proliferation of precursors of the monocyte/macrophagelineage, as well as their differentiation into phagocytes (Hamilton and Anderson, 2004). The finding that cells of the myeloid lineage are osteoclast precursors suggests that M-CSF plays an important role in osteoclast biology. Cellular response to M-CSF is regulated by the M-CSF receptor, which is a single-chain molecule encoded by the *c-fms* proto-oncogene that consists of an extracellular ligand-binding domain, single membranespanning segment, and intercellular tyrosine kinase domain. There is some evidence of a role for M-CSF/*c-fms* system in osteoclast activation during orthodontic tooth movement. Kaku et al. (2008) observed the expression of M-CSF mRNA in osteoblasts and fibroblasts during experimental tooth movement. Kitaura et al. (2008) reported that the anti*c-fms* antibody inhibited tooth movement by inhibiting osteoclastogenesis. M-CSF/c-fms system may involve the velocity of tooth movement (Yamaguchi et al., 2007).

Orthodontically induced inflammatory root resorption (OIIRR) is an unavoidable pathological consequence of orthodontic tooth movement. It can be defined as an iatrogenic disorder that occurs, unpredictably, after orthodontic treatment, whereby the resorbed apical root portion is replaced with normal bone. OIIRR is a sterile inflammatory process that is extremely complex and involves various disparate components, including mechanical forces, tooth roots, bone, cells, surrounding matrix, and certain known biological messengers (Krishnan and Davidovitch, 2006; Meikle, 2006). A recent study demonstrated that mRNA of RANK was detected in tissues involved in root resorption following application of orthodontic forces (Low et al., 2005). Furthermore, Yamaguchi et al. (2006) reported that compressed periodontal ligament (PDL) cells obtained from tissues with severe external apical root resorption produced a large amount of RANKL that up-regulated osteoclastogenesis in vitro. However, few studies have attempted to identify the concomitant expressions of RANKL/RANK and M-CSF/cfms during root resorption triggered by orthodontic force.

In the present study, the expressions of RANKL/ RANK and M-CSF/*c-fms* in rat root resorption during experimental tooth movement were investigated using immunohistochemical analysis.

Materials and methods

Animals

The animal experimental protocol in this study was approved by the Ethics Committee for Animal Experiments at Nihon University School of Dentistry at Matsudo (approval No. ECA-05-0025).

A total of 40 male, 6-week-old, Wistar strain rats (Sankyo Labo Service Co., Tokyo, Japan) weighting 180 ± 10 g were used for the experiment. The rats were divided into eight groups (five rats each), according to the magnitude and duration of the applied force. Four time groups were set at 0, 3, 7, and 10 days and two magnitude groups at 10 and 50 g. The contralateral molars served as the controls (0 g). The rats were kept in the animal centre of Nihon University School of Dentistry at Matsudo in separate cages, in a 12 hour light/dark environment at a constant temperature of 23°C, and provided with food and water *ad libitum*. The health status of each rat was evaluated by daily body weight monitoring for 1 week before the start of the experiments.

Application of orthodontic devices and tissue harvesting

The animals were anaesthetized with ketamine (90 mg/kg body weight) and rompum (10 mg/kg body weight) for insertion of orthodontic devices. Experimental tooth movement was performed using the method of Fujita *et al.* (2008), with a closed coil spring (wire size: 0.005 inch, diameter: 1/12 inch; Accurate Sales Co., Chiba, Japan) ligated to the maxillary first molar cleat by a 0.008 inch stainless steel ligature wire (Tomy International Inc., Tokyo,

Japan). The other side of the coil spring was also ligated, with the holes in the maxillary incisors drilled laterally just above the gingival papilla with a no. 1/4 round bur, using the same ligature wire. The upper first molar was moved mesially by the closed coil spring with a force of 10 or 50 g (Figure 1). The period of the experiment was 10 days.

Measurement of tooth movement

Measurement of tooth movement was performed according to the method of Fujita *et al.* (2008). To determine the amount of tooth movement, plaster models of the maxillae were made using a silicone impression material (Dent Silicone-V; Shofu Inc., Kyoto, Japan) before (day 0) and after tooth movement (days 3, 7, or 10). The models were scanned using a contact-type three-dimensional (3D) measurement apparatus (3D-picza; Roland Co., Hamamatsu, Japan) by setting the plane to pass through three points, i.e the bilateral interpapillary crests between the first and second molars and the interpapillary crest between the second and third molars.

Using 3D morphological analysis software (3D-Rugle; Medic Engineering Inc., Kyoto, Japan), the distance between the central fossa of the first molar and the mesial surface of the second molar was measured to determine tooth movement.

Tissue preparation

The experimental periods were set at 0, 3, 7, and 10 days after tooth movement. The animals were deeply anaesthetized by pentobarbital sodium and perfused with 4 per cent paraformaldehyde in 0.1 M phosphate buffer transcardially, after which the maxilla was immediately dissected and immersed in the same fixative overnight at 4°C. The specimens were decalcified in 10 per cent disodium ethylenediaminetetraacetic acid (pH 7.4) solution for 4



Figure 1 Experimental tooth movement was performed with a closed coil spring (wire size: 0.005 inch, diameter: 1/12 inch) ligated to the maxillary first molar cleat by a 0.008 inch stainless steel ligature wire. The other side of the coil spring was also ligated, with the holes in the maxillary incisors drilled laterally just above the gingival papilla with a no. 1/4 round bur, using the same ligature wire. The upper first molar was moved mesially by the closed coil spring at 10 or 50 g. The period of the experiment was 10 days.

weeks and then the decalcified specimens were dehydrated through an ethanol series and embedded in paraffin. Each sample was sliced into 6 µm continuous sections in the horizontal direction and prepared for haematoxylin and eosin (H&E) and immunohistochemistry staining for tartrate-resistant acid phosphatase (TRAP), RANK, RANKL M-CSF, and *c-fms*. The periodontal tissues in the mesial part of the mesial root of an upper first (M1) were observed.

Immunohistochemistry

Immunohistochemical staining was performed as follows: the sections were deparaffinized and endogenous peroxidase activity was quenched by incubation in 3 per cent H₂O₂ in methanol for 30 minutes at room temperature. After washing in Tris-buffered saline (TBS), the sections were incubated with polyclonal anti-rabbit TRAP (Santa Cruz Biotechnology, Inc., Santa Cruz, California, USA; working dilution, 1:400), polyclonal anti-rabbit RANK (Santa Cruz biotechnology; working dilution, 1:100), polyclonal anti-goat RANKL (Santa Cruz Biotechnology; working dilution, 1:100), polyclonal anti-rabbit M-CSF (Santa Cruz Biotechnology; working dilution: 1:300), and polyclonal anti-rabbit c-fms (Santa Cruz Biotechnology; working dilution: 1:300) overnight at 4°C. TRAP, RANK, M-CSF, and c-fms were stained using Histofine Simple Stain MAX-Po kits (Nichirei, Tokyo, Japan), and RANKL was stained using a Histofine SAB-PO (G) kit (Nichirei) according to the manufacturer's protocol.

The sections were rinsed with TBS and final colour reactions were performed using the substrate reagent 3, 3'-diaminobenzidine tetra-hydrochloride and aminoethylcarbazole and then counter-stained with haematoxylin. As immunohistochemical controls, some sections were incubated in the same way and then with either non-immune rabbit IgG or 0.01 M (phosphate-buffered saline) alone, instead of the primary antibody.

Quantitative assessment

Quantitative measurements were conducted at the root-PDL–alveolar bone complex close to the apical root, where orthodontic root resorption usually occurs (Chutimanutskul *et al.*, 2006). The slides were evaluated at a magnification of ×400 with a fixed measurement frame ($450 \times 450 \mu m$). TRAP-positive cells were counted as multinucleate odontoclasts, which were observed on the surface of cementum. The data were then processed with a Mann–Whitney *U*-test. A value of P < 0.01 was considered to indicate a significant difference.

Results

Tooth movement during the experimental period

The body weight of the rats in both force groups decreased transiently on day 1 and then recovered. No significant

differences between the two groups were observed (data not shown). The amount of tooth movement also was not significantly different between the two groups on day 10 (data not shown).

Histological changes in periodontal tissues during tooth movement (H&E staining)

In the control group (0 g) on days 3 and 7 after tooth movement, PDL specimens were composed of relatively dense connective tissue fibres and fibroblasts that regularly ran in a horizontal direction from the root cementum towards the alveolar bone. Blood capillaries were mainly recognized near the alveolar bone in the PDL. The alveolar bone and root surface were relatively smooth, with a few mononuclear and multinucleate osteoclasts, and resorption lacunae were observed on the alveolar bone surface (Figure 2a and 2b).

In the 10 g group, the arrangement of the fibres and fibroblasts became coarse and irregular, and blood capillaries were compressed on day 3. Resorption lacunae with a few multinucleate osteoclasts were observed on the surface of the alveolar bone surface (Figure 2c). On day 7, on the surface of the alveolar bone, bone resorption lacunae with multinucleate osteoclasts were observed. Osteoclasts on the alveolar bone were increased in comparison with those on day 3 (Figure 2d).

The PDL in the 50 g group was composed of a coarse arrangement of fibres and expanded blood capillaries. Many resorption lacunae with multinucleate osteoclasts were observed on the alveolar bone on day 3 (Figure 2e). On day 7, many root resorption lacunae with multinucleated odontoclasts were recognized on the surface of the root (Figure 2f).

TRAP immunohistochemical findings

In the control group (0 g) on days 3 and 7, a few resorption lacunae with TRAP-positive multinucleated osteoclasts were observed on the surfaces of the alveolar bone and root (Figure 3A-a and b).

In the 10 g group, some resorption lacunae with TRAPpositive multinucleated osteoclasts appeared on the surface of the alveolar bone on day 3 (Figure 3A-c). On day 7, on the alveolar bone surface, bone resorption lacunae with multinucleated TRAP-positive osteoclasts were recognized. The multinucleate TRAP-positive osteoclasts on the alveolar bone were increased in comparison with those on day 3 (Figure 3A-d).

In the 50 g group, on the surface of the alveolar bone, many bone resorption lacunae with multinucleate TRAPpositive osteoclasts were observed on day 3 (Figure 3A-e) and TRAP-positive odontoclasts on day 7 (Figure 3A-f).

Quantitative evaluation showed the number of TRAPpositive odontoclasts to be significantly increased in the 50 g group on day 7 (6.0-fold) as compared with the 10 g group (*P < 0.01, Figure 3B).



Figure 2 Light macroscopic images of the effect of different orthodontic forces on the multinucleate osteoclasts (haematoxylin and eosin: original magnification $\times 400$). Osteoclasts (thin arrows) appeared on the alveolar bone surface in both groups on days 3 (c and e) and 7 (d and f). Odontoclasts (thick arrows) on the cementum in the 50 g group was more than that of the 10 g on day 7 (f). AB: alveolar bone, PDL: periodontal ligament, C: cementum, and D: dentine; bar = 50 μ m. The direction of applied force is indicated by the dotted arrow.

Protein expressions of RANKL/RANK and M-CSF/c-fms

Immunoreactivity for RANKL/RANK and M-CSF/c-fms was performed on days 7 and 10 after tooth movement. In the control (0 g) group, the immunoreactivity of RANKL and M-CSF were weakly localized in cytoplasm of some fibroblasts and pericytes near the alveolar bone surface (Figures 4A-a, 4B-a, 5A-a, and 5B-a). RANK- and c-fmspositive osteoclasts were not observed (Figures 4A-b, 4B-b, 5A-b, and 5B-b). In the 10 g group, a few RANKL- and M-CSF-positive fibroblasts in the PDL and osteoblasts at the root surface were increased (Figures 4A-c, 4B-c, 5A-c, and 5B-c). RANK- and *c-fms*-positive odontoclasts were rarely observed (Figures 4A-d, 4B-d, 5A-d, and 5B-d). In the 50 g group, many RANKL- and M-CSF-positive odontoclasts and fibroblasts were observed (Figures 4A-e, 4B-e, 5A-e, and 5B-e). Many odontoclasts showed strong immunoreactivity of RANK and *c-fms* (Figures 4A-f, 4B-f, 5A-f, and 5B-f).

Discussion

Considering the method used to induce root resorption, many resorption lacunae with multinucleated osteoclasts and odontoclasts appeared on the alveolar bone and root on days 3 and 7 after tooth movement (Figures 2 and 3). Many investigators have reported that root resorption is aggravated by increasing force magnitudes (Reitan, 1985; Chan and Darendeliler, 2005). Gameiro *et al.* (2008) demonstrated osteoclastic resorption of roots on the mesial surfaces of teeth subjected to heavy orthodontic force (50 g). Gonzales *et al.* (2008) showed that 10 g of light force application produced significantly greater tooth movement with significantly less root resorption over a period of 28 days in relation to heavier force application in rats. The optimum force for the movement of the rat upper molar may be less than the 10 g previously suggested (Kohno *et al.*, 2002).

On days 7 and 10, immunoreactivity for RANKL/RANK was detected in odontoclasts in the 50 g group (Figure 4A and 4B). Oshiro et al. (2001) suggested that osteoblast/ stromal cells and PDL fibroblasts are involved in supporting osteoclast differentiation during tooth movement. Further, Low et al. (2005) reported that mRNA of RANKL was detected in tissues including root resorption subjected to orthodontic forces. Taken together, those findings and the present results suggest that osteoclast and odontoclasts differentiation in root resorption areas are critically regulated by RANKL, which is produced as a local factor by PDL cells in response to heavy orthodontic force. RANKL has been identified to be an osteoclast differentiation factor that binds to RANK on the cell surface of osteoclastic cells (Myers et al., 1999). On the other hand, OPG is a novel secreted member of the TNF receptor superfamily that works as a decoy receptor of the RANKL-RANK signalling



Figure 3 (A) Effect of different orthodontic forces on tartrate-resistant acid phosphatase (TRAP)-positive odontoclasts by immunohistochemistry (original magnification ×400). The osteoclasts (thin arrows) appeared on the alveolar bone surface in both groups on days 3 (c and e) and 7 (d and f). Immunoreactivity of TRAP positive was observed in the odontoclasts (thick arrows) on the cementum in the 50 g group (f), but not in the 10 g on day 7. AB: alveolar bone, PDL: periodontal ligament, C: cementum, and D: denine; bar = 50 µm. The direction of applied force was indicated by the dotted arrow. (B) Diagram depicting quantitative assessment of cellular changes. The number of TRAP-positive odontoclasts in the 50 g group was more than in the 10 g on days 7 and 10 (*P < 0.01).

system, thereby inhibiting osteoclastogenesis (Burgess *et al.*, 1999). Thus, OPG, RANKL, and RANK form a critical system that controls bone resorption by regulating the number and activity of osteoclasts. In relation to the OPG/RANKL/RANK system and orthodontic tooth movement, since RANKL expression was observed in compressed PDL cells, it is possible that osteoclastogenesis is promoted on the compression side during orthodontic tooth movement (Garlet *et al.*, 2007). Oshiro *et al.* (2001) reported that in OPG-deficient mice, there was a greater induction of TRAP-positive multinucleated cells in the PDL

on the compressed side and in the adjacent alveolar bone after experimental application of orthodontic force than in their wild-type OPG (+/+) littermates. Kanzaki *et al.* (2004) stated that OPG gene transfer to periodontal tissue inhibited RANKL-mediated osteoclastogenesis and inhibited experimental tooth movement. Nishijima *et al.* (2006) found an increased concentration of RANKL in gingival crevicular fluid (GCF) during orthodontic tooth movement, and the ratio of concentration of RANKL to that of OPG in the GCF was significantly higher than at the control sites. These studies suggest that during orthodontic tooth movement, the



Figure 4 Effect of different orthodontic forces on RANKL- and RANK-positive odontoclasts by immunohistochemistry on days 7 (Figure 4A) and 10 (Figure 4B; original magnification ×400). Immunoreactivity of RANKL and RANK was observed in the odontoclasts (arrow) on the cementum in the 50 g group (4AB-e, f), but not in the 10 g (4AB-c, d) on days 7 and 10. PDL: periodontal ligament, C: cementum, D: dentine; bar = 50 μ m. The direction of applied force is indicated by the dotted arrow.

OPG/RANKL/RANK system in the periodontal tissues is an important determinant regulating balanced alveolar bone resorption. Taken together, those findings and the present results suggest that the OPG/RANKL/RANK system may play an important role in orthodontically induced root resorption. Further studies are necessary to confirm the relationship between OPG and RANKL/RANK in root resorption.

On the other hand, M-CSF stimulates not only the proliferation of osteoclast progenitors but also their



Figure 5 Effect of different orthodontic forces on macrophage colony stimulating factor (M-CSF)- and *c-fms*-positive odontoclasts by immunohistochemistry on days 7 (Figure 4A) and 10 (Figure 4B; original magnification ×400). Immunoreactivity of M-CSF and *c-fms* were observed in the odontoclasts (arrows) on the cementum in the 50 g group (4AB-e, f), but not in the 10 g (4AB-c, d) on days 7 and 10. PDL: periodontal ligament, C: cementum, D: dentine; bar = 50 μ m. The direction of applied force is indicated by the dotted arrow.

differentiation into mature osteoclasts *in vivo* (Felix *et al.*, 1994) and *in vitro* (Tanaka *et al.*, 1993). Yang *et al.* (1996) reported that the presence of the M-CSF receptor on preformed osteoclasts is carried over from the developing

cells and plays a functional role in the modulation of bone resorption by mature cells. *c-fms*, the cellular homologue of the feline transforming virus v-FMS, is the sole receptor for M-CSF (Stanley *et al.*, 1997). The functional linkage

between M-CSF and its receptor has been established by the finding that mice lacking csflr, the gene coding for c-fms, exhibit the same major phenotype as op/op mice, namely a marked decrease in tissue macrophages and severe osteopetrosis due to a lack of osteoclasts (Dai *et al.*, 2002).

Previous studies have demonstrated that a severe deficiency of osteoclasts in *op/op* mice can be reversed by injections of recombinant human M-CSF (Kodama et al., 1991a: Wiktor-Jedrzeiczak et al., 1991), and the direct action of M-CSF on osteoclast lineage cells was exhibited by expression of the receptors for M-CSF and c-fms in osteoclasts both in vitro (Kodama et al., 1991b) and in vivo (Hofstetter et al., 1995). Those findings indicate that M-CSF plays an essential role in the differentiation of osteoclasts. In addition, it has become clear that M-CSF supports osteoclast differentiation in co-operation with RANKL (Lacey et al., 1998; Yasuda et al., 1998, Udagawa et al., 1999). Kimoto et al. (1999) suggested that M-CSF may affect bone remodelling and root resorption in PDL cells derived from human primary teeth in response to mechanical stress in vitro. On days 7 and 10, immunoreactivity for M-CSF/c-fms was also detected in odontoclasts in the 50 g group (Figure 5A and 5B). Therefore the M-CSF and *c-fms* system may also play an important role in orthodontic root resorption.

Conclusions

On days 7 and 10, immunoreactivity for RANKL/RANK and M-CSF/*c-fms* was detected in rat odontoclasts and PDL fibroblasts with an orthodontic force of 50 g. Therefore, RANKL/RANK and M-CSF/*c-fms* systems may be involved in the process of root resorption by excessive orthodontic force.

Funding

Japan Society for the Promotion of Science (C:18592252, C:22592297).

References

- Burgess T L et al. 1999 The ligand for osteoprotegerin (OPGL) directly activates mature osteoclasts. Journal of Cell Biology 145: 527–538
- Chan E, Darendeliler M A 2005 Physical properties of root cementum: Part 5. Volumetric analysis of root resorption craters after application of light and heavy orthodontic forces. American Journal of Orthodontics and Dentofacial Orthopedics 127: 186–195
- Chutimanutskul W, Darendeliler M A, Shen G, Petocz P, Swain M V 2006 Changes in the physical properties of human premolar cementum after application of 4 weeks of controlled orthodontic forces. European Journal of Orthodontics 28: 313–318
- Dai X M *et al.* 2002 Targeted disruption of the mouse colony-stimulating factor 1 receptor gene results in osteopetrosis, mononuclear phagocyte deficiency, increased primitive progenitor cell frequencies, and reproductive defects. Blood 99: 111–120

- Felix R, Hofstetter W, Wetterwald A, Cecchini M G, Fleisch H 1994 Role of colony-stimulating factor-1 in bone metabolism. Journal of Cellular Biochemistry 55: 340–349
- Fujita S, Yamaguchi M, Utsunomiya T, Yamamoto H, Kasai K 2008 Low-energy laser stimulates tooth movement velocity via expression of RANK and RANKL. Orthodontics and Craniofacial Research 11: 143–155
- Gameiro G H *et al.* 2008 Evaluation of root resorption associated with orthodontic movement in stressed rats. Minerva Stomatologica 57: 569–575
- Garlet T P, Coelho U, Silva J S, Garlet G P 2007 Cytokine expression pattern in compression and tension sides of the periodontal ligament during orthodontic tooth movement in humans. European Journal of Oral Science 115: 355–362
- Gonzales C, Hotokezaka H, Yoshimatsu M, Yozgatian J H, Darendeliler M A, Yoshida N 2008 Force magnitude and duration effects on amount of tooth movement and root resorption in the rat molar. Angle Orthodontist 78: 502–509
- Hamilton J A, Anderson G P 2004 GM-CSF biology. Growth Factors 22: 225–231
- Hofstetter W, Wetterwald A, Cecchini M C, Felix R, Fleisch H, Mueller C 1995 Detection of transcripts for the receptor for macrophage colonystimulating factor, c-fms, in murine osteoclasts. Bone 17: 145–151
- Kaku M et al. 2008 VEGF and M-CSF levels in periodontal tissue during tooth movement. Biomedical Research 29: 181–187
- Kanzaki H, Chiba M, Takahashi I, Haruyama N, Nishimura M, Mitani H 2004 Local OPG gene transfer to periodontal tissue inhibits orthodontic tooth movement. Journal of Dental Research 83: 920–925
- Kimoto S *et al.* 1999 Cytokine secretion of periodontal ligament fibroblasts derived from human deciduous teeth: effect of mechanical stress on the secretion of transforming growth factor-beta 1 and macrophage colony stimulating factor. Journal of Periodontal Research 34: 235–243
- Kitaura H *et al.* 2008 An anti-c-Fms antibody inhibits orthodontic tooth movement. Journal of Dental Research 87: 396–400
- Kodama H *et al.* 1991a Congenital osteoclast deficiency in osteopetrotic (op/op) mice is cured by injections of macrophage colony-stimulating factor. Journal of Experimental Medicine 173: 269–272
- Kodama H, Nose M, Niida S, Yamasaki A 1991b Essential role of macrophage colony-stimulating factor in the osteoclast differentiation supported by stromal cells. Journal of Experimental Medicine 173: 1291–1294
- Kohno T, Matsumoto Y, Kanno Z, Warita H, Soma K 2002 Experimental tooth movement under light orthodontic forces: rates of tooth movement and changes of the periodontium. Journal of Orthodontics 29: 129–135
- Krishnan V, Davidovitch Z 2006 Cellular, molecular, and tissue-level reactions to orthodontic force. American Journal of Orthodontics and Dentofacial Orthopedics 129: 469.e1–469.e32
- Lacey L D et al. 1998 Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. Cell 93: 165–176
- Lossdörfer S, Götz W, Jäger A 2002 Immunohistochemical localization of receptor activator of nuclear factor kappaB (RANK) and its ligand (RANKL) in human deciduous teeth. Calcified Tissue International 71: 45–52
- Low E, Zoellner H, Kharbanda O P, Darendeliler M A 2005 Expression of mRNA for osteoprotegerin and receptor activator of nuclear factor kappa β ligand (RANKL) during root resorption induced by the application of heavy orthodontic forces on rat molars. American Journal of Orthodontics and Dentofacial Orthopedics 128: 497–503
- Meikle M C 2006 The tissue, cellular, and molecular regulation of orthodontic tooth movement: 100 years after Carl Sandstedt. European Journal of Orthodontics 28: 221–240
- Myers D E *et al.* 1999 Expression of functional RANK on mature rat and human osteoclasts. FEBS Letters 463: 295–300
- Nishijima Y, Yamaguchi M, Kojima T, Aihara N, Nakajima R, Kasai K 2006 Levels of RANKL and OPG in gingival crevicular fluid during orthodontic tooth movement and effect of compression force on releases from periodontal ligament cells *in vitro*. Orthodontics and Craniofacial Research 9: 63–70

RANKL/RANK AND M-CSF/C-FMS IN ROOT RESORPTION

- Oshiro T, Shibasaki Y, Martin T J, Sasaki T 2001 Immunolocalization of vacuolar-type H+-ATPase, cathepsin K, matrix metalloproteinase-9, and receptor activator of NFkappaB ligand in odontoclasts during physiological root resorption of human deciduous teeth. Anatomical Record 264: 305–311
- Reitan K 1985 Biomechanical principles and reactions. In: Graber T M, Swain B F (eds). Orthodontics: current principles and techniques Mosby, St Louis, pp. 101–192
- Shiotani A, Shibasaki Y, Sasaki T 2001 Localization of receptor activator of NFkappaB ligand, RANKL, in periodontal tissues during experimental movement of rat molars. Journal of Electron Microscopy (Tokyo) 50: 365–369
- Soedarsono N *et al.* 2006 Evaluation of RANK/RANKL/OPG gene polymorphisms in aggressive periodontitis. Journal of Periodontal Research 41: 397–404
- Stanley E R et al. 1997 Biology and action of colony—stimulating factor-1. Molecular Reproduction and Development 46: 4–10
- Tanaka S*etal.* 1993 Macrophage colony-stimulating factor is indispensable for both proliferation and differentiation of osteoclast progenitors. Journal of Clinical Investigation 91: 257–263

- Udagawa N *et al.* 1999 Osteoblasts/stromal cells stimulate osteoclast activation through expression of osteoclast differentiation factor/ RANKL but not macrophage colony-stimulating factor: receptor activator of NF-kappa B ligand. Bone 25: 517–523
- Wiktor-Jedrzejczak W et al. 1991 Correction by CSF-1 of defects in the osteopetrotic op/op mouse suggests local, developmental, and humoral requirements for this growth factor. Experimental Hematology 19: 1049–1054
- Yamaguchi M, Aihara N, Kojima T, Kasai K 2006 RANKL increase in compressed periodontal ligament cells from root resorption. Journal of Dental Research 85: 751–756
- Yamaguchi M et al. 2007 Low-energy laser irradiation stimulates the tooth movement velocity via expression of M-CSF and c-fms. Orthodontic Waves 66: 139–148
- Yang S, Zhang Y, Rodriguiz R M, Ries W L, Key Jr L L 1996 Functions of the M-CSF receptor on osteoclasts. Bone 18: 355–360
- Yasuda H et al. 1998 Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. Proceedings of the National Academy of Sciences the United States of America 95: 3597–3602

Copyright of European Journal of Orthodontics is the property of Oxford University Press / UK and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.