The perception of facial aesthetics in a young Spanish population

Ana B. Macías Gago*, Martín Romero Maroto* and Antonio Crego** Departments of *Orthodontics and **Psychology, Rey Juan Carlos University, Madrid, Spain

Correspondence to: Ana Belén Macías Gago, Private practice, C/Ortega y Gasset nº1 1º H, 24400 Ponferrada, León, Spain. E-mail: ana.macias@urjc.es

SUMMARY Improved facial aesthetics is one aim of orthodontic treatment. This study was designed to determine if the faces considered more beautiful in a young population exhibit the same parameters used by orthodontists to assess successful results.

A panel of 34 laypeople (30 females and 4 males) evaluated a set containing one frontal, one frontal during smiling, and one profile photograph of 89 students (77 females and 12 males) on a 5-point attractiveness scale, in relation to a set of reference photographs. For each photographic set, the mean and final scores were calculated. Once the sample was established, 11 subjects (9 females and 2 males) with the highest final facial aesthetic score were selected and cephalometric analysis was performed.

All cephalometric measurements were within the norm for the total sample. When the sample was divided by gender, Wilcoxon's *W* non-parametric test showed significant differences between the male and female photographs; while females tended to a Class II malocclusion, with the mandible slightly retrusive to the maxilla, males tended to a Class III and showed a straighter profile with a prominent chin; the face height ratio was higher in males. There were no significant differences between genders for lower lip to E plane. The findings show that the faces considered more attractive fulfilled the cephalometric and facial norms.

Introduction

The face has been exhaustively studied by scientists, doctors, and artists; most of whom have tried to measure and reproduce facial features, especially those related to beauty.

Beauty and physical attractiveness are of importance to humans; social acceptance, popularity, mate selection, and careers are all affected by an individual's physical attractiveness (Dipboye et al., 1977; Langlois and Roggman, 1990; O'Doherty et al., 2003). But what makes a face attractive? is beauty 'altogether in the eye of the beholder?' Aesthetic criteria appear to have been defined in almost all cultures (Peck and Peck, 1970; Matoula and Pancherz, 2006), but several findings suggest that the perception of beauty may be innate and, additionally, universal or crosscultural (Langlois and Roggman, 1990; Langlois et al., 1991, 1994; Cunningham et al., 1995). Accordingly, Langlois and Roggman (1990) and Langlois et al. (1994) proposed that averageness is attractive (a face is perceived as attractive when its facial gestalt is close to the average or mean of a population of faces).

Improved facial aesthetics is one of the main aims of orthodontic treatment, and, in recent years, it has acquired even more importance both for patients and orthodontists. Orthodontic patients and their parents believe that having well-aligned teeth is an important factor in facial appearance; they hope that orthodontic treatment will improve their dental, dentofacial, and facial aesthetics and consequently their popularity and social success (Shaw *et al.*, 1985; Kerosuo *et al.*, 1995; Birkeland *et al.*, 2000; Kiekens *et al.*, 2006). A significant correlation has been found between orthodontic treatment and facial appeal (Tatarunaite *et al.*, 2005). However, orthodontists are not always aware of the difference in perception between patients and clinicians as to which result defines treatment success. Whereas patients hope for results determined by social and cultural rules of beauty in their reference group and in society in general, orthodontists prefer to use parameters to determine diagnosis and subsequent treatment planning (Spyropoulos and Halazonetis, 2001; Kiekens *et al.*, 2006; McKoy-White *et al.*, 2006; Miner *et al.*, 2007). This study was designed to determine if faces considered more beautiful in a young population exhibit the same parameters used by orthodontists to assess ideal treatment and successful results.

Subjects and methods

Approval for this study was obtained from the ethical committee of Rey Juan Carlos University and all subjects gave informed consent to participation.

Reference set

A total of 91 (78 females and 13 males, between 20 and 34 years of age) third- and fourth-year students pursuing a degree in Dentistry at Rey Juan Carlos University, who fulfilled the following criteria, were included: no facial or dental trauma, no congenital defects, and who did not wear glasses.

A frontal relaxed, a frontal during smile, and profile photographs were taken of each subject, using a Coolpix 5700 digital camera (5 MP; Nikon, Tokyo, Japan). The photographs were obtained with the Frankfort plane parallel to the floor and perpendicular to the body axis (natural position) and with the interpupilar parallel to the floor. The participants stood against a white background at a distance of 1.5 m from the camera. The photographs were sorted using Microsoft Power Point for Windows XP (2003 version) and a digital photographic panel was produced, in which three views of each individual were displayed simultaneously (Figure 1) (Kiekens *et al.*, 2006).

Judges

Fifty-one fifth-year students (43 females and 8 males) from the dentistry programme of Rey Juan Carlos University, between 22 and 26 years of age, scored the photographs of the 91 subjects. Rating of facial aesthetics was performed on a 5-point attractiveness scale with values from 1 (very unattractive) to 5 (very attractive) (Ong *et al.*, 2006).

The mean and standard deviation (values indicating the final facial aesthetic score for each individual) were calculated for each slide using Microsoft Excel for Windows XP. The mean scores ranged from 1.32 to 3.8, median 2.42. A slide of a male and female (with a final facial aesthetic score close to the median value) was then selected to serve as a reference set for the measuring system (Figure 2) (Shell and Woods, 2004; Kiekens *et al.*, 2005).

Evaluation of the sample

The photographs (frontal relaxed, frontal during smiling, and profile) of each subject, together with a set of reference photographs, were shown to a group of judges formed of 34 students (30 females and 4 males, between 20 and 26 years of age) from the second year of the Physiotherapy Diploma Course of the Rey Juan Carlos University. From the initial sample, the two slides comprising the reference set were eliminated, so the number of slides shown was 89. The slides were randomly placed, showing every female face in relation to the female reference set.

Each slide was shown for 15 seconds (Kiekens *et al.*, 2005). The rating system was the same as used for the reference set. For each slide, the mean and standard deviation were again calculated as the final facial aesthetic score. For the 11 subjects with highest facial aesthetic final score, lateral cephalometric radiographs were obtained and the following objective parameters were measured (Table 1): SNA, SNB, ANB, facial angle, and facial convexity; these values indicate the maxillomandibular relationship in the antero-posterior plane or skeletal Class, face height ratio (Jarabak); this proportion indicates vertical discrepancy; and lower lip to the E plane.

A. B. MACÍAS GAGO ET AL.



Figure 1 Digital picture panel.



Figure 2 Reference photographs for a male and female. The final facial aesthetic score was 2.38 and 2.5, respectively.

 Table 1
 Cephalometric measurements.

1. SNA: 82 ± 2° (Steiner, 1953) 2. SNB: 80 ± 2° (Steiner, 1953)	Sella–nasion–subnasale (point A) angle Sella–nasion–supramentale (point B)
(2000, 200)	angle
3. ANB: 2 ± 1° (Steiner, 1953)	Point A-point B-nasion angle
4. Facial angle: $87 \pm 3^{\circ}$	Frankfort horizontal plane-facial
Ricketts, 1960)	plane (nasion to pogonion) angle
5. Facial convexity: $2 \pm 2 \text{ mm}$	Point A to facial plane
Ricketts, 1960)	
5. Face height ratio: $61 \pm 2\%$	Posterior face height (sella-gonion) to
Siriwat and Jarabak, 1985)	anterior face height (nasion-menton)
7. Lower lip to the E plane:	
$-2 \pm -2 \text{ mm}$ (Ricketts, 1960)	

All cephalometric measurements were undertaken twice by the same author (ABMG), with an interval of 2 weeks. For each variable, the average value of the two measurements was used.

All data analyses were carried out using the software PASW (SPSS) version 17 (PAWS/SPSS; IBM Company, Armonk, New York, USA). As the panel of judges contained a higher number of females than males (30 females and 4 males), the intraclass correlation coefficient (ICC), two-way random effects, was calculated to assess inter-rater reliability in scoring the 89 photographs. To evaluate reproducibility of the measuring system, six duplicate slides were used in order to standardize the judges in terms of scoring.

Results

The cephalometric analysis performed on the 11 subjects to obtain the different variables previously indicated showed that all measurements were inside the norm for the total sample (Table 2). When divided by gender, Wilcoxon's *W* non-parametric test showed significant differences between the male and female photographs for the following measurement: SNB (W = 45.0; Z = -2.13; P < 0.05), higher in males than females; ANB (W = 3.0; Z = -2.14; P < 0.05), higher in females than in males; facial convexity (W = 3.5; Z = -2.01; P < 0.05), higher in females than in males; and face height ratio (W = 45.0; Z = -2.12; P < 0.05), higher in males than in females. No significant gender differences were found for SNA, facial angle, or lower lip to E plane (Table 3).

The ICC was 0.962 (lower bound 0.95; upper bound 0.973, with 95% confidence) indicating a high level of agreement among the 34 judges, when scoring each of the photographs. Spearman's rho correlations ranged from 0.46 to 0.75 (P = 0.001) for the five pairs of repeated photographs, indicating a moderate level of reproducibility for the scores of the same slide. Nevertheless, there was no correlation between scores for one of the duplicated presentations (rho = 0.05, P > 0.05).

Discussion

A system to evaluate facial aesthetics should be simple, applicable to the everyday clinic, and provide quantitative data (Moyers, 1992). The use of lateral radiographs, profile silhouettes, or photographs has been reported in the literature (Czarnecki *et al.*, 1993; Spyropoulos and Halazonetis, 2001), but the most complex system is probably simultaneous visualization of a frontal and profile photograph and of the smile (Kiekens *et al.*, 2005). Howells and Shaw (1985) demonstrated a high correlation between scores assigned to individuals classified *in vivo* and photographs of the same individuals; they also demonstrated that evaluation of facial aesthetics using photographs and a panel of judges is valid and reproducible.

Differences have been found in the evaluation of facial aesthetics, depending on panel composition, with regard to age, gender, and demographic origin (Howells and Shaw, 1985; Newton and Minhas, 2005; Kiekens *et al.*, 2007). As differences in judging facial attractiveness between orthodontists and non-orthodontists (Knight and Keith, 2005; Maple *et al.*, 2005) have also been found, the raters, who were not dental personnel, in the present study were selected to be similar to the subjects on the basis of age, gender, and demographic origin. Since the aims of the study were not only to judge facial aesthetics but also to determine the skeletal features of the subjects considered the most attractive, the sample included a wide range of faces with different features (Kiekens *et al.*, 2008).

When analysing the results of the research without dividing the model by gender, the variables indicating skeletal Class (ANB, facial angle, and facial convexity) had values within

 Table 2
 Mean and standard deviation (SD) of the cephalometric measurements of the 11 subjects with the highest final facial aesthetic score.

Gender		SNA (°)	SNB (°)	ANB (°)	Facial angle (°)	Facial convexity (mm)	Face height ratio (%)	Lower lip to E plane (mm)
Male, $n = 2$	Mean	83.50	83.00	0.50	91.00	-1.75	79.46	-4.50
	SD	0.70	0.00	0.70	0.00	1.06	1.89	0.70
Female, $n = 9$	Mean	78.50	75.27	3.22	89.50	2.05	65.99	-1.66
	SD	3.42	2.85	1.34	3.58	1.89	5.26	2.01
Total, <i>n</i> = 11	Mean	79.40	76.68	2.72	89.77	1.36	68.44	-2.18
	SD	3.67	4.03	1.64	3.26	2.31	7.22	2.14

Table 3 Wilconxon's non-parametric test for the cephalometric measurements of males and females.

	SNA (°)	SNB (°)	ANB (°)	Facial angle (°)	Facial convexity (mm)	Face height ratio (%)	Lower lip to E plane (mm)
<i>W</i> of Wilcoxon	46.50	45.00	3.00	49.00	3.50	45.00	4.50
Z	-1.77	-2.13	-2.14	-1.19	-2.01	-2.12	-1.78
Asymptotic significance (two-tailed)	0.07	0.03	0.03	0.23	0.04	0.03	0.07
Exact significance [2 × (significance one-tailed)]	0.07	0.03	0.03	0.32	0.03	0.03	0.07

Grouping variable: gender of slides.

the norm, i.e. the faces considered more attractive had a skeletal Class I occlusion. This is in agreement with previous research where it has been suggested that a Class I occlusion is the more attractive profile (Tulloch *et al.*, 1993; Kitay *et al.*, 1999; Knight and Keith, 2005; Maple *et al.*, 2005; Tatarunaite *et al.*, 2005; Chan *et al.*, 2008).

Although SNB angle in the sample was within the norm, the value was slightly diminished and was similar to that obtained by Johnston *et al.* (2005a) as representative of facial attractiveness. It indicates that a convex profile, with a prominent maxilla relative to the mandible, is a feature of facial attractiveness, which is in agreement with the results of Sforza *et al.* (2007). Because diminution of SNB angle causes an increase in ANB angle, this variation in angles should be minimal for a face to be attractive. Knight and Keith (2005) showed that an ANB angle deviated more than 5 degrees from the norm causes a diminution in facial attractiveness. Johnston *et al.* (2005a) also showed that a Class III outline is considered more attractive than a Class II, with a similar degree of skeletal discrepancy.

When analysing face height ratios, it was observed that even though the value was within the norm, it was slightly increased when compared with the norm determined by Siriwat and Jarabak (1985). This indicates that not only is this proportion related to facial attractiveness (Lundström et al., 1987; Johnston et al., 2005b) but also that the most attractive faces are shorter. Lundström et al. (1987) found that a horizontal growth pattern corresponded to a higher facial attractiveness. Johnston et al. (2005b) showed that the most attractive faces corresponded with a neutral face height ratio, but they also showed, in agreement with Michiels and Sather (1994), that a diminished inferior face height is more acceptable when it comes to judging facial attractiveness. This indicates that treatment that increases face height should be avoided (Padrós, 2000; Johnston et al., 2005b).

The position of the lips, as determined by maxillomandibular protrusion or retrusion, by dental protrusion or retrusion and/ or by lip thickness (Sarver and Jacobson, 2007), was correct for the total sample; therefore, although nowadays the tendency is for protrusive lips to be considered as more attractive, and variations in occlusion and in the thickness of the upper and lower lip vermilion have a significant influence on facial attractiveness perception (Nguyen and Turley, 1998; Kiekens *et al.*, 2006; Scott *et al.*, 2006; McNamara *et al.*, 2008), a correct relationship with Ricketts line should be maintained.

When the sample was divided by gender, it was observed that for the variables that indicate skeletal Class, ANB angle and facial convexity had higher values in females; this indicates a tendency to a Class II in females, with more convex profiles. This result is in accordance with the findings of Matoula and Pancherz (2006), who reported that ANB angle was higher in attractive females, and Foster (1973), who found that in males a straight profile was considered attractive, whereas in females, convex profiles are considered attractive. According to previous studies, more convex faces have a younger appearance than more concave faces (Nguyen and Turley, 1998; Peck and Peck, 1995) and statistically significant differences were found when comparing facial attractiveness and estimated age, with those perceived as younger being considered more attractive (Tatarunaite *et al.*, 2005). In the same way, Czarnecki *et al.* (1993) reported that straight profiles with a prominent chin are more suited to males than females.

Height ratio was significantly higher in males considered as attractive, indicating a more brachyfacial type. For females, 44 per cent displayed a neutral type and the remaining 55.6 per cent a brachyfacial type. This result is in keeping with the findings of Lundström *et al.* (1987), who showed that the most attractive female profiles were those with neutral growth, followed by brachyfacial; while for males, the most attractive profiles were brachyfacial.

Matoula and Pancherz (2006) did not find differences between attractive and unattractive females with regard to face height. Knight and Keith (2005) showed that a decrease in females and an increase in males in height ratio were associated with less attractive faces. This result is contrary to the present findings for males; this could be because in the study of Knight and Keith (2005), the relationship between height ratio and facial attractiveness was not significant and to the fact that the present sample was small (11 individuals).

Significant differences between males and females were not found for the parameter lower lip to E plane. This result is contrary to the findings of Czarnecki *et al.* (1993) and Padrós (2000) who reported that labial protrusion was better tolerated in females than in males. Matoula and Pancherz (2006) also found an increased distance of the lower lip to the aesthetic line in unattractive females.

Conclusions

- 1. The faces considered more attractive in this study fulfilled the cephalometric and facial norms commonly used for diagnosis and treatment planning; more attractive individuals have a skeletal Class I, mesobrachyfacial type, and the lips inside Ricketts' aesthetic line.
- 2. Attractive females showed a tendency to a Class II, with the mandible slightly retrusive to the maxilla; therefore, they present more convex profiles than males. By contrast, attractive males presented a straighter profile with a prominent chin.
- 3. The facial pattern is more horizontal in attractive males than in attractive females.
- 4. None of the faces considered attractive had a moderate or severe Class II or Class III skeletal malocclusion.
- 5. None of the faces considered attractive had a dolichofacial pattern.

References

- Birkeland K, Bøe O E, Wisth P J 2000 Relationship between occlusion and satisfaction with dental appearance in orthodontically treated and untreated groups. A longitudinal study. European Journal of Orthodontics 22: 509–518
- Chan E K M, Soh J, Petocz P, Darendeliler M A 2008 Esthetic evaluation of Asian-Chinese profiles from a white perspective. American Journal of Orthodontics and Dentofacial Orthopedics 133: 532–538
- Cunningham M R, Roberts A R, Barbee A P, Druen P B, Wu C H 1995 Their ideas of beauty are, on the whole, the same as ours: consistency and variability in the cross-cultural perception of female physical attractiveness. Journal of Personality and Social Psychology 68: 261–279
- Czarnecki S T, Nanda R S, Currier G F 1993 Perceptions of a balanced facial profile. American Journal of Orthodontics and Dentofacial Orthopedics 104: 180–187
- Dipboye R L, Arvey R D, Terpstra D E 1977 Sex and physical attractiveness of raters and applicants as determinants of resume evaluation. Journal of Applied Psychology 4: 288–294
- Foster E 1973 Profile preferences among diversified groups. Angle Orthodontist 43: 34–40
- Howells D J, Shaw W C 1985 The validity and reliability of ratings of dental and facial attractiveness for epidemiologic use. American Journal of Orthodontics 88: 402–408
- Johnston C, Hunt O, Burden D, Stevenson M, Hepper P 2005a The influence of mandibular prominence on facial attractiveness. European Journal of Orthodontics 27: 129–133
- Johnston D J, Hunt O, Johnston C D, Burden D J, Stevenson M, Hepper P 2005b The influence of lower face vertical proportion on facial attractiveness. European Journal of Orthodontics 27: 349–354
- Kerosuo H, Hausen H, Laine T, Shaw W C 1995 The influence of incisal maloclusion on the social attractiveness of young adults in Finland. European Journal of Orthodontics 17: 505–512
- Kiekens R M A, Maltha J C, van 't Hof M A, Kuijpers-Jagtman A M 2005 A measuring system for facial aesthetic in Caucasian adolescents: reproducibility and validity. European Journal of Orthodontics 27: 579–584
- Kiekens R M A, Maltha J C, van 't Hof M A, Kuijpers-Jagtman A M 2006 Objective measures as indicators for facial esthetics in white adolescents. Angle Orthodontist 76: 551–556
- Kiekens R M A, Van 't Hof M A, Straatman H, Kuijpers-Jagtman A M, Maltha J C 2007 Influence of panel composition on aesthetic evaluation of adolescent faces. European Journal of Orthodontics 29: 95–99
- Kiekens R M A, Kuijpers-Jagtman A M, van 't Hof M A, van 't Hof B E, Straatman H, Maltha J C 2008 Facial esthetics in adolescents and its relationship to ideal ratios and angles. American Journal of Orthodontics and Dentofacial Orthopedics 133: 188.e1–188.e8
- Kitay D, BeGole E, Evans C A, Giddon D B 1999 Computer-animated comparison of self-perception with actual profiles of orthodontic and nonorthodontic subjects. International Journal of Adult Orthodontics and Orthognathic Surgery 14: 125–134
- Knight H, Keith O 2005 Ranking facial attractiveness. European Journal of Orthodontics 27: 340–348
- Langlois J H, Ritter J M, Roggman L A, Vaughn L S 1991 Facial diversity and infants preferences for attractive faces. Developmental Psychology 27: 79–84
- Langlois J H, Roggman L A 1990 Attractive faces are only average. Psychological Science 1: 115–121
- Langlois J L, Roggman L A, Musselman L 1994 What is average and what is not average about attractive faces? Psychological Science 5: 214–220
- Lundström A, Woodside D G, Popovich F 1987 Panel assessments of facial profile related to mandibular growth direction. European Journal of Orthodontics 9: 271–278
- Maple J R, Vig K W L, Beck F M, Larsen P E, Shanker S 2005 A comparison of providers' and consumers' perceptions of facial-profile attractiveness. American Journal of Orthodontics and Dentofacial Orthopedics 128: 690–696

- Matoula S, Pancherz H 2006 Skeletofacial morphology of attractive and nonattractive faces. Angle Orthodontist 76: 204–210
- McKoy-White J, Evans C A, Viana G, Anderson N K, Giddon D B 2006 Facial profile preferences of black women before and after orthodontic treatment. American Journal of Orthodontics and Dentofacial Orthopedics 129: 17–23
- McNamara L, McNamara J A Jr, Ackerman M B, Baccetti T 2008 Hardand soft-tissue contributions to the esthetics of the posed smile in growing patients seeking orthodontic treatment. American Journal of Orthodontics and Dentofacial Orthopedics 133: 491–499
- Michiels G, Sather A H 1994 Determinants of facial attractiveness in a sample of white women. International Journal of Adult Orthodontics and Orthognathic Surgery 9: 95–103
- Miner R M, Anderson N K, Evans C A, Giddon D B 2007 The perception of children's computer-imaged facial profiles by patients, mothers and clinicians. Angle Orthodontist 77: 1034–1039
- Moyers R E 1992 Evolution of concepts of the face. In: McNamara J A (ed.). Esthetics and the treatment of facial form. Monograph No. 28, Craniofacial Growth Series, Center for Human Growth and Development, University of Michigan, Ann Arbour, pp. 27–39
- Newton J T, Minhas G 2005 Exposure to 'ideal' facial images reduces facial satisfaction: an experimental study. Community Dentistry and Oral Epidemiology 33: 410–418
- Nguyen D D, Turley P K 1998 Changes in the Caucasian male facial profile as depicted in fashion magazines during the twentieth century. American Journal of Orthodontics and Dentofacial Orthopedics 114: 208–217
- O'Doherty J, Winston J, Critchley H, Perrett D, Burt D M, Dolan R J 2003 Beauty in a smile: the role of medial orbitofrontal cortex in facial atractiveness. Neuropsychologia 41: 147–155
- Ong E, Brown R, Richmond S 2006 Peer assessment of dental attractiveness. American Journal of Orthodontics and Dentofacial Orthopedics 130: 163–169
- Padrós E 2000 Patrón actual de estética facial. Ortodoncia clínica 3: 72-85
- Peck H, Peck S 1970 A concept of facial aesthetics. Angle Orthodontist 40: 284–318
- Peck S, Peck L 1995 Selected aspects of the art and science of facial esthetics. Seminars in Orthodontics 1: 105–126
- Ricketts R 1960 The influence of orthodontic treatment on facial growth and development. Angle Orthodontist 30: 103–133
- Sarver D, Jacobson R S 2007 The aesthetic dentofacial analysis. Clinics in Plastic Surgery 34: 369–394
- Scott C R, Goonewardene M S, Murray K 2006 Influence of lips on the perception of malocclusion. American Journal of Orthodontics and Dentofacial Orthopedics 130: 152–162
- Sforza C, Laino A, D'Alessio R, Dellavia C, Grandi G, Ferrario V F 2007 Threedimensional facial morphometry of attractive children and normal children in the deciduous and early mixed dentition. Angle Orthodontist 77: 1025–1033
- Shaw W C, Rees G, Dawe M, Charles C R 1985 The influence of dentofacial appearance on the social attractiveness of young adults. American Journal of Orthodontics 87: 21–26
- Shell T L, Woods M G 2004 Facial aesthetics and the divine proportion: a comparison of surgical and non-surgical Class II treatment. Australian Orthodontic Journal 20: 51–63
- Siriwat P P, Jarabak J R 1985 Maloclusion and facial morphology Is there a relationship? An epidemiologic study. Angle Orthodontist 55: 127–138
- Spyropoulos M N, Halazonetis D J 2001 Significance of the soft tissue profile on facial esthetics. American Journal of Orthodontics and Dentofacial Orthopedics 119: 464–471
- Steiner C C 1953 Cephalometrics for you and me. American Journal of Orthodontics 39: 729–755
- Tatarunaite E, Playle R, Hood K, Shaw W, Richmond S 2005 Facial attractiveness: a longitudinal study. American Journal of Orthodontics and Dentofacial Orthopedics 127: 676–682
- Tulloch C, Phillips C, Dann I V C 1993 Cephalometric measures as indicators of facial attractiveness. International Journal of Adult Orthodontics and Orthognathic Surgery 8: 171–179

Copyright of European Journal of Orthodontics is the property of Oxford University Press / UK and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.