# Association between root resorption incident to orthodontic treatment and treatment factors

# Masahide Motokawa, Tomoko Sasamoto, Masato Kaku, Toshitsugu Kawata, Yayoi Matsuda, Akiko Terao and Kazuo Tanne

Department of Orthodontics and Craniofacial Developmental Biology, Graduate School of Biomedical Sciences, Hiroshima University, Japan

*Correspondence to:* Dr Masahide Motokawa, Department of Orthodontics and Craniofacial Developmental Biology, Graduate School of Biomedical Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8553, Japan. E-mail: motumo@hiroshima-u.ac.jp

SUMMARY The purpose of this study was to clarify the prevalence and degree of root resorption induced by orthodontic treatment in association with treatment factors. The files of 243 patients (72 males and 171 females) aged 9–51 years were randomly selected from subjects treated with multi-bracket appliances. The severity of root resorption was classified into five categories on radiographs taken before and after treatment. The subjects were divided into extraction (n = 113 patients, 2805 teeth) and non-extraction (n = 130 patients, 3616 teeth) groups and surgical (n = 56 patients, 1503 teeth) and non-surgical treatment (n = 187 patients, 4918 teeth) groups. These subjects were also divided into two or three groups based on the duration of multiloop edgewise archwire (MEAW) treatment, elastic use, and total treatment time: 0 month (T1; n = 184 patients, 4831 teeth), range 1–6 months (T2; n = 37 patients, 3914 teeth) more than 6 months ( $T_3$ ; n = 22 patients, 596 teeth); range 0–6 months (n = 114 patients, 3016 teeth) more than 30 months (n = 95 patients, 2508 teeth). The prevalence of overall and severe root resorption evaluated by the number of subjects and teeth was compared with a chi-square test. A Student's *t*-test for unpaired data was used to determine any statistically significant differences.

The prevalence of severe root resorption based on the number of teeth was significantly higher in the group with extractions (P < 0.01). Longer use of a MEAW appliance and elastics also produced a significantly higher prevalence of root resorption (P < 0.05). On the other hand, the prevalence of severe root resorption was not significantly different between the subjects treated with or without surgery, but there was a significant increase when treatment time was prolonged (P < 0.05). A significant difference was found in the amount of root movement of the upper central incisors and the distance from their root apices to the cortical bone surface (P < 0.05). These are regarded as essential factors in the onset of root resorption. These results indicate that orthodontic treatment with extractions, long-term use of a MEAW appliance and elastics, treatment time, and distance of tooth movement are risk factors for severe root resorption.

# Introduction

Apical root resorption during orthodontic treatment is an unfavourable sequelae, which is difficult to predict and repair. There is an individual difference in the prevalence, indicating the influence of host factors. For the prevention of severe root resorption, the progress of root resorption should be periodically assessed on radiographs taken during orthodontic treatment. In addition, in subjects with severe root resorption, treatment should be interrupted and the treatment plan changed.

Root resorption associated with orthodontic treatment has been recognized since the report of Ketcham (1927). Later, Massler and Malone (1954) found root resorption in 86.4 per cent of orthodontic patients. Risks for root resorption such as systemic factors (Linge and Linge, 1983; Goldie and King, 1984), treatment mechanics (Malmgren *et al.*, 1982; Linge and Linge, 1983), treatment period (Linge and Linge, 1983; Levander and Malmgren, 1988; Mirabella and Årtun, 1995), age (Oppenheim, 1942; Newman, 1975; Linge and Linge, 1983), root shape (Goldie and King, 1984; Levander and Malmgren, 1988; Mirabella and Årtun, 1995), density of alveolar bone (Oppenheim, 1942; Goldin, 1989; Kaley and Phillips, 1991), and oral habits (Odenrick and Brattström, 1985) have been investigated.

However, the above studies evaluated risk factors for root resorption in terms of the number of subjects, which may lead to an incorrect interpretation because patients with the same degree of root resorption have a different number of teeth with resorption. Furthermore, no studies have evaluated orthodontic treatment factors relevant to severe root resorption in all types of teeth. Thus, the purpose of the present study was to clarify the prevalence and degree of root resorption induced by orthodontic treatment with special reference to treatment factors.

#### Subjects and methods

The study was approved by the Ethics Committee of Hiroshima University (Epidemiologic study-142; 1 June 2009).

The files of 243 patients (72 males and 171 females) treated with multi-bracket appliances ( $0.018 \times 0.025$  inch bracket slots) were randomly selected from those attending the Orthodontic Clinic at Hiroshima University Hospital. Treatment in all patients resulted in an excellent occlusion with optimal tooth alignment and root parallelism. Those with incomplete records and poor quality radiographs or with incomplete root formation and missing or extracted teeth were excluded. Thus, the subjects included in the present study were aged 9–51 years before treatment [mean 18.9, standard deviation (SD) 6.7] and were treated for 0.4–5 years (mean 2.4, SD 0.8). A total of 243 patients and 6421 teeth from the incisors to second molars in both arches were evaluated.

The subjects were divided into extraction (n = 113) patients, 2805 teeth) and non-extraction groups (n = 130) patients, 3616 teeth) and surgical (n = 56) patients, 1503 teeth) and non-surgical treatment groups (n = 187) patients, 4918 teeth). These subjects were also divided into two or three groups based on the duration of multiloop edgewise archwire (MEAW) treatment, elastic use, and total treatment time: 0 month (T1; n = 184 patients, 4831 teeth), range 1–6 months (T2; n = 37 patients, 596 teeth); range 0–6 months (n = 114) patients, 3016 teeth) and more than 6 months (n = 129) patients, 3405 teeth); range 1–30 months (n = 148) patients, 3913 teeth) and more than 30 months (n = 95) patients, 2508 teeth).

The radiographs were obtained using the paralleling long cone technique before and after treatment. The periapical radiographs were scanned with a GT-X970 scanner (Seiko-Epson Corp., Nagano, Japan) with a resolution of 675 dpi at a scale of 1:1. The images were analysed with Photoshop software (version 7.0; Adobe Systems, San Jose, California, USA) at 300 per cent enlargement, without image quality loss. The degree of root resorption was evaluated according to five categories defined by Malmgren *et al.* (1982): 0 = no root resorption, 1 = irregular root contour, 2 = apical root resorption less than 2 mm of the original root length, 3 = apical root resorption from 2 mm to one-third of the original root length, and 4 = root resorption exceeding one-third of the original root length.

Duplicate determination of root resorption categories after treatment showed full agreement for 1870 of 2150 teeth (87 per cent) and a disagreement of one category for 280 teeth (13 per cent). Prior to the examination, duplicate determination of 2150 teeth was performed by two authors (MM and MK). Furthermore, all subjects and teeth with root resorption (n = 190 patients, 1450 teeth) were divided

into two groups with mild (categories 1 and 2) and severe (categories 3 and 4) resorption. For both groups, the following orthodontic treatment factors were examined: orthodontic treatment with or without extractions and surgery, the duration of MEAW treatment and elastic use, and treatment time.

To compare the movement of maxillary central incisors with a normal root (category 0, normal group) and incisors with severe root resorption (categories 3 and 4, severe group), 20 subjects were randomly selected from the 243 patients for each group. The subjects satisfied the following criteria: no differences in inclination and rotation of the right and left maxillary incisors and no tooth with an abnormal pre-treatment root shape. The normal group consisted of 5 males and 15 females aged 11–47 years before treatment (mean 21.4, SD 9.3), treated for 1.8–3.5 years (mean 2.4, SD 0.4), and the group with severe root resorption consisted of 8 males and 12 females aged 13–33 years before treatment (mean 19.7, SD 5.7) who were treated for 1.8–3.8 years (mean 2.8, SD 0.5).

On the pre-treatment lateral cephalogram, the trabecular pattern of the maxilla and the outlines of the pterygomaxillary fissure and key ridges were traced. The incisal edge (U1) and the apex (U1') of the maxillary incisors were also traced. The palatal plane (PP) was drawn as a line connecting anterior and posterior nasal spines (ANS and PNS), and point A' was as an intersection of a vertical line from point A to PP. The pre-treatment cephalometric tracing was superimposed on the corresponding post-treatment cephalogram according to the 'best anatomic fit', defined as the judge's best estimate of the optimal fit of the hard palate and anterior maxillary images when primary emphasis is given to concordance in the region of the PP and point A'. The post-treatment U1' was copied from the pre- to the post-treatment tracing using incisors 'template'. The distances and angles obtained by connecting the related landmarks were measured at the levels of 0.1 mm and 0.5 degrees, respectively, as suggested by Mirabella and Årtun (1995). The amount of movement of the central incisors was evaluated as shown in Figure 1.

The reproducibility of the measurements was assessed by statistically analysing the difference between double measurements taken with an interval of at least 1 week for 40 cephalograms (20 pre- and 20 post-treatment) selected at random. The measurement error was calculated according to the following equation:

$$Sx = \sqrt{\frac{\sum D^2}{2N}},$$

where *D* is the difference between duplicate measurements and *N* is the number of double measurements. The errors were 0.21 mm for Dx, 0.31 mm for Dy, 0.56 mm for Dz, 0.46 mm for Dx', 0.19 mm for Dy', 0.18 mm for Dz', 0.42 degrees for  $\angle A$ , and 0.46 degrees for  $\angle B$ .



Figure 1 Measurements of maxillary central incisor movement.

The prevalence of overall and severe root resorption, evaluated by the number of subjects and teeth, was compared using a chi-square test (Statview; Abacus Concepts, Inc., Berkeley, California, USA). The mean distances and angles obtained by measuring the movement of the central incisors were calculated for the two subgroups of patients and teeth with normal and severe root resorption. A Student's *t*-test for unpaired data was used to determine any significant differences. The level of significance was defined as P < 0.05.

#### Results

No significant differences in the prevalence of root resorption were observed in terms of the percentages evaluated by the number of patients and teeth in the groups with or without extractions (Table 1). For severe root resorption, no significant differences were found in the number of patients, but the prevalence of severe root resorption evaluated by the number of teeth was significantly greater in the extraction than in the non-extraction group (number of teeth; P = 0.0004; Table 2).

With respect to the prevalence of root resorption evaluated by the number of patients and teeth, no significant differences were found between the two patient groups with or without surgery (Table 1) or in the prevalence of severe root resorption between the two groups in terms of the numbers of patients or teeth (Table 2). **Table 1** The prevalence of root resorption between subjects with or without extractions and surgery or no surgery, the duration of multiloop edgewise archwire treatment [0 month (T1), range 1–6 months (T2), and more than 6 months (T3)], elastic use (range 0–6 months or more than 6 months), and treatment time (range 1–30 months or more than 30 months) for the number of patients and teeth.

|   | Rate resor | of root<br>ption (%)                     | P value |
|---|------------|--|---------|
| The number of patients                    |            |  |         |
| Extraction group $(n = 113)$              | 81.4       | ٦  | 0.28    |
| Non-extraction group $(n = 130)$          | 75.4       |  | 0.20    |
| Surgical treatment group $(n = 56)$       | 82.1       | -<br>T                                   | 0.37    |
| Non-surgical treatment group $(n = 187)$  | 77.0       |  |         |
| T1 (n = 184)                              | 78.3       | 7 9                                      | a; 0.69 |
| T2(n = 37)                                | 75.7       | $\begin{bmatrix} a \\ b \end{bmatrix} c$ | b: 0.28 |
| T3(n = 22)                                | 81.8       |  | c; 0.51 |
| Range 0–6 months $(n = 114)$              | 74.6       | ٦  | 0.27    |
| More than 6 months $(n = 129)$            | 81.4       |  |         |
| Range 1–30 months $(n = 148)$             | 70.3       | ٦  | 0.00**  |
| More than 30 months $(n = 95)$            | 90.5       |  |         |
| The number of teeth                       |            |  |         |
| Extraction group $(n = 2805)$             | 26.6       | 7  | 0.22    |
| Non-extraction group $(n = 3616)$         | 19.5       |  |         |
| Surgical treatment group $(n = 1503)$     | 23.8       | ٦  | 0.79    |
| Non-surgical treatment group $(n = 4918)$ | 22.2       |  |         |
| T1 $(n = 4831)$                           | 22.0       | a  | a; 0.94 |
| T2 $(n = 994)$                            | 21.5       | b c                                      | b; 0.23 |
| T3 $(n = 596)$                            | 28.7       |  | c; 0.27 |
| Range 0–6 months ( $n = 3016$ )           | 21.3       | 7  | 0.67    |
| More than 6 months $(n = 3405)$           | 23.7       |  |         |
| Range 1–30 months ( $n = 3913$ )          | 19.3       | 7  | 0.16    |
| More than 30 months ( $n = 2508$ )        | 27.7       |  |         |
|   |            | -  |         |

a = T1 versus T2; b = T2 versus T3; c = T1 versus T3. \*\*P < 0.01.

For the relationship between the duration of MEAW treatment and the prevalence of root resorption, no significant differences were observed in the number of patients or teeth with root resorption (Table 1). There were also no significant differences in the number of patients with severe root resorption, but the prevalence evaluated by the number of teeth was significantly higher in the T3 than in the T1 and T2 groups (number of teeth; T1 versus T3, P = 0.00003, T2 versus T3, P = 0.04; Table 2), indicating a significantly greater prevalence of severe root resorption associated with long-term use of MEAW appliances.

For the relationship between the period of elastic use and root resorption, no significant differences in prevalence were observed between the two groups in terms of the numbers of patients and teeth (Table 1). For severe root resorption, no significant difference was found in the number of patients, but the prevalence of severe root resorption evaluated by the number of teeth was significantly higher in the group that used elastics for more than 6 months (number of teeth; P = 0.003; Table 2).

A statistically significant difference was observed for the relationship between treatment time and prevalence of root resorption when evaluated by the number of patients (P = 0.0002) but not by the number of teeth (Table 1). With

**Table 2** The prevalence of mild and severe root resorption between subjects with or without extractions, and surgery or no surgery, the duration of multiloop edgewise archwire treatment [0 month (T1), range 1–6 months (T2), or more than 6 months (T3)], elastic use (range 0–6 months or more than 6 months), and treatment time (range 1–30 months or more than 30 months) for the number of patients and teeth.

|   | Mild resorption |      | Severe resorption |      |   | P value   |  |
|---|-----------------|------|-------------------|------|---|-----------|--|
|   | N               | %    | N                 | %    |   |           |  |
| The number of patients                    |                 |      |                   |      |   |           |  |
| Extraction group $(n = 113)$              | 66              | 71.7 | 26                | 28.3 | ٦ | 0.91      |  |
| Non-extraction group $(n = 130)$          | 71              | 72.4 | 27                | 27.6 |   |           |  |
| Surgical treatment group $(n = 56)$       | 32              | 69.6 | 14                | 30.4 | ٦ | 0.66      |  |
| Non-surgical treatment group $(n = 187)$  | 105             | 72.9 | 39                | 27.1 |   |           |  |
| T1 $(n = 184)$                            | 108             | 75.0 | 36                | 25.0 | ] | a;0.43    |  |
| T2(n = 37)                                | 19              | 67.9 | 9                 | 32.1 |   | b; 0.40   |  |
| T3(n = 22)                                | 10              | 55.6 | 8                 | 44.4 |   | c;0.08    |  |
| Range 0–6 months $(n = 114)$              | 63              | 74.1 | 22                | 25.9 | ٦ | 0.58      |  |
| More than 6 months $(n = 129)$            | 74              | 70.5 | 31                | 29.5 |   |           |  |
| Range 1–30 months $(n = 148)$             | 81              | 77.9 | 23                | 22.1 | ٦ | 0.04*     |  |
| More than 30 months $(n = 95)$            | 56              | 65.1 | 30                | 34.9 |   |           |  |
| The number of teeth                       |                 |      |                   |      |   |           |  |
| Extraction group $(n = 2805)$             | 643             | 86.2 | 103               | 13.8 | ٦ | 0.00**    |  |
| Non-extraction group $(n = 3616)$         | 648             | 92.0 | 56                | 8.0  |   |           |  |
| Surgical treatment group $(n = 1503)$     | 316             | 88.5 | 41                | 11.5 | ٦ | 0.72      |  |
| Non-surgical treatment group $(n = 4918)$ | 975             | 89.2 | 118               | 10.8 |   |           |  |
| T1 (n = 4831)                             | 966             | 90.7 | 99                | 9.3  | ] | a: 0.20   |  |
| T2(n = 994)                               | 188             | 87.9 | 26                | 12.1 |   | b: 0.04*  |  |
| T3 $(n = 596)$                            | 139             | 80.3 | 34                | 19.7 |   | c: 0.00** |  |
| Range 0–6 months ( $n = 3016$ )           | 589             | 91.7 | 53                | 8.3  | 7 | 0.00**    |  |
| More than 6 months $(n = 3405)$           | 704             | 86.9 | 106               | 13.1 |   |           |  |
| Range 1–30 months ( $n = 3913$ )          | 692             | 91.5 | 64                | 8.5  | 7 | 0.00**    |  |
| More than 30 months $(n = 2508)$          | 599             | 86.3 | 95                | 13.7 |   |           |  |

a = T1 versus T2; b = T2 versus T3; c = T1 versus T3.

\**P* < 0.05; \*\**P* < 0.01.

respect to severe root resorption, the prevalence obtained by the number of patients and teeth was significantly higher in the group treated for more than 30 months (number of patients; P = 0.05, number of teeth; P = 0.001; Table 2).

Displacements of maxillary central incisors, Dx, Dx', Dy', Dz', and changes in tooth inclination were significantly larger in the group with severe root resorption than in the group without resorption (Dx, P = 0.0007; Dx', P = 0.04; Dy', P = 0.04; Dz', P = 0.009; changes in tooth inclination, P = 0.003). Significant differences were found in the vertical and horizontal displacements of U1, the vertical displacement of U1', the change in tooth inclination during treatment, and the distance between U1' and the cortical bone surface (Table 3).

# Discussion

Apical root resorption that occurs during orthodontic treatment is difficult to predict and repair. Massler and Malone (1954) found root resorption in 86.4 per cent of orthodontic patients. The incidence of root resorption was 78.2 per cent in the present study. However, it is difficult to compare these results because the methods for evaluating the degree of root resorption were different.

In previous studies, treatment-related factors for root resorption were listed as the amount and direction of tooth movement (Stuteville, 1938; Phillips, 1955; Hollender et al., 1980), the type of treatment technique (Kaley and Phillips, 1991; Beck and Harris, 1994; Parker and Harris, 1998), history of extractions (McFadden et al., 1989; Lee et al., 1999; Sameshima and Sinclair, 2001), treatment time (Phillips, 1955; Linge and Linge, 1983; McFadden et al., 1989; Sameshima and Sinclair, 2001), type of wire used (Mirabella and Årtun, 1995; Sameshima and Sinclair, 2001), the period of elastic use (Linge and Linge, 1983; Mirabella and Årtun, 1995; Sameshima and Sinclair, 2001), the amount and direction of orthodontic force (Kurol et al., 1996; Owman-Moll et al., 1995, 1996), and the distance between the apex and cortical bone surface (Mirabella and Årtun, 1995; Kaley and Phillips, 1991). In the present study, the relationship between root resorption and factors such as orthodontic treatment with extractions, surgical treatment, the MEAW technique and elastic use, treatment time, and the amount of movement of upper incisors were investigated.

Lee *et al.* (1999) and McFadden *et al.* (1989) observed a relationship between root resorption and orthodontic treatment with extractions. Sameshima and Sinclair (2001)

**Table 3** The amount of movement of the maxillary central incisors with normal (category 0) and severe (categories 3 and 4) root resorption, which was evaluated as Dx: the horizontal movement of U1, Dx': the horizontal movement of U1', Dy: the vertical movement of U1, Dy': the vertical movement of U1', Dz: the distance between U1' and the cortical bone surface at pretreatment, Dz': the distance between U1' and the cortical bone surface at post-treatment,  $\angle A$ : the angles U1–U1/PP at pretreatment,  $\angle B$ : the angles U1–U1/PP at post-treatment,  $\angle A-\angle B$ : change in tooth inclination during treatment.

|                 | Group             | Mean  | SD   | P value |
|-----------------|-------------------|-------|------|---------|
| Dx (mm)         | Normal $(n = 20)$ | 1.6   | 1.2  | 0.00**  |
|                 | Severe $(n = 20)$ | 3.4   | 2.0  |         |
| Dx' (mm)        | Normal $(n = 20)$ | 1.4   | 0.9  | 0.04*   |
|                 | Severe $(n = 20)$ | 2.2   | 1.4  |         |
| Dy (mm)         | Normal $(n = 20)$ | 1.1   | 1.1  | 0.89    |
|                 | Severe $(n = 20)$ | 1.1   | 0.6  |         |
| Dy' (mm)        | Normal $(n = 20)$ | 1.0   | 0.7  | 0.04*   |
|                 | Severe $(n = 20)$ | 1.6   | 1.1  |         |
| Dz (mm)         | Normal $(n = 20)$ | 4.8   | 1.4  | 0.23    |
|                 | Severe $(n = 20)$ | 5.5   | 1.9  |         |
| Dz' (mm)        | Normal $(n = 20)$ | 4.6   | 1.8  | 0.00**  |
|                 | Severe $(n = 20)$ | 6.5   | 2.5  |         |
| ∠A (°)          | Normal $(n = 20)$ | 114.1 | 8.5  | 0.54    |
|                 | Severe $(n = 20)$ | 115.9 | 10.0 |         |
| ∠B (°)          | Normal $(n = 20)$ | 114.3 | 8.5  | 0.30    |
|                 | Severe $(n = 20)$ | 117.4 | 10.4 |         |
| Change in tooth | Normal $(n = 20)$ | 4.2   | 2.8  | 0.00**  |
| inclination (°) | Severe $(n = 20)$ | 9.1   | 6.5  |         |

\**P* < 0.05; \*\**P* < 0.01.

reported that patients with four first premolar extractions had greater resorption than those without extractions. In the present study, no significant difference in the prevalence of root resorption was found between patients with or without extractions. However, the prevalence of severe root resorption was higher in the extraction group than in the non-extraction group when evaluated by the number of teeth but not by the number of patients. Parker and Harris (1998) indicated that the amount of tooth movement was a highly predictive parameter, explaining up to 90 per cent of the variation in root resorption. They also identified apical vertical movements and incisor proclination as strong predictors of external apical root resorption. From the present results, it was considered that an increase in tooth movement in the extraction group might cause severe root resorption.

Kaley and Phillips (1991) reported that a Le Fort I osteotomy is a risk factor for apical root resorption. Periods of ischaemia and hyperaemia after Le Fort I osteotomy (Ramsay *et al.*, 1991), which may result in long-term pulpal changes (Ellingsen and Årtun, 1993), may be a reason for root resorption after maxillary osteotomy. On the other hand, this was contradicted by Mirabella and Årtun (1995) who did not find any association between Le Fort I osteotomy and root resorption. In the present study, no significant differences were found in the prevalence of root

resorption between patients with or without surgical orthodontic treatment. A similar finding was noted for the prevalence of severe root resorption; however, a tendency towards the higher prevalence of overall and severe root resorption was observed in patients who had undergone surgical orthodontic treatment. Linge and Linge (1991) reported that functional soft tissue factors such as the lips and tongue were significantly associated with apical root resorption. Thus, from these findings, it is considered that an increase in chronic tongue pressure and circumoral muscle tonus after surgical treatment might increase the prevalence of root resorption.

There have been no reports on the relationship between the period of MEAW use and the prevalence of root resorption. In the present study, no significant differences were observed in the number of patients and teeth. However, the prevalence of severe root resorption in terms of the number of teeth was significantly greater in the T3 group than in T1 and T2 groups, whereas no significant difference was found between T1 and T2. From these results, it is demonstrated that orthodontic treatment with the MEAW is not a risk factor for severe root resorption, but the use of the MEAW for more than 6 months causes the most severe root resorption. It is considered that long-term 'jiggling' forces to the anterior teeth might increase root resorption, due to less cooperation of patients in using elastics to eliminate the force applied to the teeth.

Various studies have been conducted on the relationship between elastics and root resorption. Mirabella and Årtun (1995) reported that the duration of Class II elastics was associated with canine root resorption. Linge and Linge (1983, 1991) reported that increased apical root resorption was associated with the use of Class II elastics, contrary to the findings of Sameshima and Sinclair (2001). In the present study, no significant difference was found in the prevalence of root resorption between the two groups where elastics were used for less or more than 6 months. For severe root resorption, the prevalence was significantly higher in the group that used elastics for more than 6 months. Thus, it is hypothesized that long-term jiggling force caused by intermittent use of elastics might increase the prevalence of root resorption.

There have been many reports concerning the relationship between root resorption and treatment time. Linge and Linge (1983), Mirabella and Årtun (1995), Lee *et al.* (1999), Phillips (1955), and Morse (1971) reported a negative conclusion; however, Sameshima and Sinclair (2001), DeShields (1969), and McFadden *et al.* (1989) found that long-term orthodontic treatment increased the prevalence of root resorption. From the present results, a significant difference was observed in terms of the number of patients, and a tendency to a higher prevalence of overall root resorption was also recognized for the number of teeth. Furthermore, the prevalence of severe root resorption evaluated by the number of patients and teeth was significantly higher in the group treated for more than 30 months. It was also shown that an increase in overall root resorption leads to an increase in severe root resorption because of continuous stimulation to the root. It is therefore considered that accumulation of surface root resorption might lead to the onset of severe root resorption.

There have been conflicting conclusions concerning the association between root resorption and the distance the teeth are moved. Some studies reported a relationship between root resorption and distance (Stuteville, 1938; Hollender et al., 1980), whereas Phillips (1955) and Dermaut and De Munck (1986) did not find any association. Displacements of Dx, Dx', Dy', Dz', and changes in tooth inclination were significantly greater in the group with severe root resorption than in the controls. A significant difference was observed in the vertical and horizontal displacement of U1, the vertical movement of U1', the change in tooth inclination, and the distance between U1' and the cortical bone surface during treatment. Furthermore, displacement of the maxillary central incisors in the severe group was classified as follows: 1. lingual inclination moved the apex close to the labial cortical bone, 2. buccal inclination moved the apex close to the lingual cortical bone, 3. lingual root torque moved the apex close to the lingual cortical bone, 4. lingual bodily movement moved the apex close to the lingual cortical bone. Anatomically, the cortical bone of the alveolus is fourth on the buccal side. In addition, there are the cortical bones of the incisive canal and alveolar bone on the lingual side. It was therefore hypothesized that a maxillary central incisor near the cortical bone of the alveolus and incisive canal might cause severe root resorption. Moreover, the type of tooth movement such as tipping, bodily movement, and torque might be a risk factor.

Other studies evaluated risk factors for root resorption in terms of the number of patients or restricted teeth, which may lead to an incorrect interpretation because patients with the same degree of root resorption had various numbers of root-resorbed teeth and this may not apply to teeth with different root shapes. In this investigation, it was shown that the prevalence of severe root resorption in the number of teeth resulted in a clearer tendency than that number of patients.

#### Conclusions

Long-term orthodontic treatment increases the prevalence of root resorption, extractions, the period of the MEAW and elastic use, treatment time, and distance of tooth movement are regarded as risk factors for severe root resorption. Among these, long-term use of the MEAW produced the highest prevalence of severe root resorption. Moreover, an increase in overall root resorption leads to an increase in severe root resorption during orthodontic treatment. These should be considered as the predictive factors for overall and severe root resorption.

# Funding

Grant-in-Aid for Young Scientists (B) (No. 20791576) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

### Acknowledgements

We are grateful to Drs Ohtani and Inbushi for their helpful advice.

#### References

- Beck B W, Harris E F 1994 Apical root resorption in orthodontically treated subjects: analysis of edgewise and light wire mechanics. American Journal of Orthodontics and Dentofacial Orthopedics 105: 350–361
- Dermaut L R, De Munck A 1986 Apical root resorption of upper incisors caused by intrusive tooth movement: a radiographic study. American Journal of Orthodontics and Dentofacial Orthopedics 90: 321–326
- DeShields R W 1969 A study of root resorption in treated Class II division 1 malocclusion. Angle Orthodontist 39: 231–245
- Ellingsen R H, Årtun J 1993 Pulpal response to orthognathic surgery: a long-term radiographic study. American Journal of Orthodontics and Dentofacial Orthopedics 103: 338–343
- Goldie R S, King G J 1984 Root resorption and tooth movement in orthodontically treated, calcium-deficient, and lactating rats. American Journal of Orthodontics 85: 424–430
- Goldin B 1989 Labial root torque: effect on the maxilla and incisor root apex. American Journal of Orthodontics and Dentofacial Orthopedics 95: 208–219
- Hollender L, Rönnerman A, Thilander B 1980 Root resorption, marginal bone support and clinical crown length in orthodontically treated patients. European Journal of Orthodontics 2: 197–205
- Kaley J, Phillips C 1991 Factors related to root resorption in edgewise practice. Angle Orthodontist 61: 125–132
- Ketcham A H 1927 A preliminary report of an investigation of apical root resorption of permanent teeth. International Journal of Orthodontics 13: 97–99
- Kurol J, Owman-Moll P, Lundgren D 1996 Time-related root resorption after application of a controlled continuous orthodontic force. American Journal of Orthodontics and Dentofacial Orthopedics 110: 303–310
- Lee R Y, Årtun J, Alonzo T A 1999 Are dental anomalies risk factors for apical root resorption in orthodontic patients? American Journal of Orthodontics and Dentofacial Orthopedics 116: 187–195
- Levander E, Malmgren O 1988 Evaluation of the risk of root resorption during orthodontic treatment: a study of upper incisors. European Journal of Orthodontics 10: 30–38
- Linge B O, Linge L 1983 Apical root resorption in upper anterior teeth. European Journal of Orthodontics 5: 173–183
- Linge L, Linge B O 1991 Patient characteristics and treatment variables associated with apical root resorption during orthodontic treatment. American Journal of Orthodontics and Dentofacial Orthopedics 99: 35–43
- Malmgren O, Goldson L, Hill C, Orwin A, Petrini L, Lundberg M 1982 Root resorption after orthodontic treatment of traumatized teeth. American Journal of Orthodontics 82: 487–491
- Massler M, Malone A J 1954 Root resorption in human permanent teeth: a roentgenographic study. American Journal of Orthodontics 40: 619–633

- McFadden W M, Engstrom C, Engstrom H, Anholm J M 1989 A study of the relationship between incisor intrusion and root shortening. American Journal of Orthodontics and Dentofacial Orthopedics 96: 390–396
- Mirabella A D, Årtun J 1995 Risk factors for apical root resorption of maxillary anterior teeth in adult orthodontic patients. American Journal of Orthodontics and Dentofacial Orthopedics 108: 48–55
- Morse P H 1971 Resorption of upper incisors following orthodontic treatment. Dental Practitioner and Dental Record 22: 21–35
- Newman W G 1975 Possible etiologic factors in external root resorption. American Journal of Orthodontics 67: 522–529
- Odenrick L, Brattström V 1985 Nailbiting: frequency and association with root resorption during orthodontic teatment. British Journal of Orthodontics 12: 78–81
- Oppenheim A 1942 Human tissue response to orthodontic intervention of short and long duration. American Journal of Orthodontics 28: 263–301
- Owman-Moll P, Kurol J, Lundgren D 1995 Continuous versus interrupted continuous orthodontic force related to early tooth movement and root resorption. Angle Orthodontist 65: 395–401

- Owman-Moll P, Kurol J, Lundgren D 1996 The effects of a four-fold increased orthodontic force magnitude on tooth movement and root resorptions. An intra-individual study in adolescents. European Journal of Orthodontics 18: 287–294
- Parker R J, Harris E F 1998 Directions of orthodontic tooth movements associated with external apical root resorption of the maxillary central incisor. American Journal of Orthodontics and Dentofacial Orthopedics 114: 677–683
- Phillips R W 1955 Apical root resorption under orthodontic therapy. Angle Orthodontist 25: 1–22
- Ramsay D, Årtun J, Bloomquist D 1991 Orthognathic surgery and pulpal blood flow: a pilot study using laser Doppler flowmetry. Journal of Oral and Maxillofacial Surgery 49: 2–8
- Sameshima G T, Sinclair P M 2001 Predicting and preventing root resorption: Part II. Treatment factors. American Journal of Orthodontics and Dentofacial Orthopedics 119: 511–515
- Stuteville O H 1938 Injuries caused by orthodontic forces and the ultimate results of these injuries. American Journal of Orthodontics and Oral Surgery 24: 103–116

Copyright of European Journal of Orthodontics is the property of Oxford University Press / UK and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.