# Design and implementation of a hybrid genetic algorithm and artificial neural network system for predicting the sizes of unerupted canines and premolars

## S. Moghimi\*, M. Talebi\*\* and I. Parisay\*\*\*

\*Department of Electrical Engineering, Ferdowsi University of Mashhad, \*\*Department of Pedodontics, Mashhad University of Medical Sciences and \*\*\*Department of Pedodontics, Shahid Sadooghi University of Medical Sciences, Yazd, Iran

*Correspondence to:* Maryam Talebi, Department of Pedodontics, Dental School, Mashhad University of Medical Sciences, Mashhad, Iran. E-mail: talebim@mums.ac.ir

SUMMARY The aim of this study was to develop a novel hybrid genetic algorithm and artificial neural network (GA–ANN) system for predicting the sizes of unerupted canines and premolars during the mixed dentition period.

This study was performed on 106 untreated subjects (52 girls, 54 boys, aged 13–15 years). Data were obtained from dental cast measurements. A hybrid GA–ANN algorithm was developed to find the best reference teeth and the most accurate mapping function. Based on a regression analysis, the strongest correlation was observed between the sum of the mesiodistal widths of the mandibular canines and premolars and the mesiodistal widths of the mandibular first molars and incisors (r = 0.697). In the maxilla, the highest correlation was observed between the sum of the mandibular first molars and maxillary central incisors (0.742). The hybrid GA–ANN algorithm selected the mandibular first molars and incisors and the maxillary central incisors as the reference teeth for predicting the sum of the mesiodistal widths of the canines and premolars. The prediction error rates and maximum rates of over/underestimation using the hybrid GA–ANN algorithm were smaller than those using linear regression analyses.

### Introduction

Predicting the size of unerupted teeth during the mixed dentition period is important in managing the developing occlusion of growing children. Accurate prediction can help determine whether the available space in the posterior segments is sufficient to allow the permanent teeth to erupt freely and in good alignment in their respective arches (Kirschen *et al.*, 2000). Three prediction methods are typically used: direct measurement of the unerupted tooth size on radiographs (Staley *et al.*, 1984; De Paula *et al.*, 1995), calculations from prediction equations and tables (Tanaka and Johnston, 1974; Ferguson *et al.*, 1978; Moyers, 1988), and a combination of radiographic measurements and prediction tables (Hixon and Oldfather, 1958; Staley *et al.*, 1984; Bishara *et al.*, 1989).

The Hixon and Oldfather approach is considered to be the most accurate (Gardner, 1979; Irwin *et al.*, 1995), but it is complex and difficult to implement. The Tanaka and Johnston prediction equations and table are also widely used (Al-Khadra, 1993; Schirmer and Wiltshire, 1997). However, they were developed for white North American children and their use is questionable in other populations since tooth sizes vary significantly between and within different ethnicities (Bishara *et al.*, 1989; Al-Khadra, 1993; Schirmer and Wiltshire, 1997). Moyer's regression scheme has achieved widespread clinical acceptance because of its simplicity and ease of application. Several simple linear regression equations have also been proposed for populations of different ethnic origins (Al-Khadra, 1993; Yuen *et al.*, 1998; Lee-Chan *et al.*, 1998; Jaroontham and Godfrey, 2000; Diagne *et al.*, 2003), with mixed dentition analyses varying between different racial and population groups.

The increasing use of information systems in health care and the considerable growth of medical databases require traditional data analyses to be adjusted to new more efficient computational models. Methods based on artificial intelligent algorithms have found numerous applications in different biomedical sciences. Genetic algorithms (GAs) and artificial neural networks (ANNs) have been widely used for predicting and interpreting biological activities, separately or as hybrid structures (Niwa, 2004; Zheng and Thomson, 2005; Fernandez and Caballero, 2006). In particular, ANNs are widely used to analyze medical data (Meersman *et al.*, 1996; Lo *et al.*, 1998; Ngan and Hu, 1999) and have also been used for dental caries prediction (Devito *et al.*, 2008).

The aim of this study was to develop a hybrid GA–ANN algorithm to identify the best reference teeth for accurately predicting the size of unerupted canines and premolars.

#### Subjects and methods

#### Data preparation

This cross-sectional study was carried out on 106 subjects (52 girls, 54 boys, aged 13–15 years) selected from a group of 1400 students in the seven regional directories of Mashhad, Iran. Informed consent was obtained prior to the investigation. Inclusion criteria were: complete permanent dentition with no caries, proximal restorations, attrition, or dental anomalies; all teeth fully erupted to the occlusal plane; no previous or ongoing orthodontic treatment; and no transverse discrepancies, such as a crossbite or scissor bite.

Polysilicone impressions (Coltene speedex, Polysiloxane condensation type silicone elastomer, Whaledent Inc., Mahwah, New Jersey, USA) were obtained from each subject and were poured in type IV plaster (Zhermack Elite Rock Gypsums, Elite Rock Class IV Die Stone, Zhermack SpA., Badia Polesine, Italy) on the same day by a technician. Measurements of the maximum mesiodistal widths of all mandibular and maxillary incisors, canines, premolars, and first molars (upper lateral incisors were not included as predictors because of their form variability) were carried out with an electronic digital vernier calliper (Zhenjiang Richoice Machinery Imp. & Exp. Co., Ltd, Jiangsu, China), with an accuracy of  $\pm 0.02$  mm and repeatability of  $\pm 0.01$ mm. For easier access to the interdental spaces, the measuring tips of the calliper were narrowed. Since the narrowed tips fit the interdental spaces, they provide reliable estimations of the teeth sizes. The measurements were obtained perpendicular to the occlusal plane. All the measurements were carried out by one author (IP) and repeated after 24 hours. Intra-observer reliability was predetermined as 0.2 mm. When the two sets of measurements varied by 0.2 mm or less, the measurements were averaged; otherwise, a new set of measurements was made and the nearest three measurements were averaged.

#### Development of the intelligent hybrid algorithm

The role of GA is to find the best reference input for the prediction process, while the ANN searches for the best mapping function to predict the targets (sizes of premolars and canines) based on the inputs provided by the GA.

The GA is a search algorithm based on survival of the fittest among string structures. Since 10 measurements were made on each dental cast (six on the upper dental cast and four on the lower dental cast), the GA chromosomes consisted of  $10 \times 1$  binary vectors consisting of 0s and 1s. To obtain the best possible results, a fitness function was defined based on the error of the neural network, as follows:

$$F = \|T - O\|_{1} + \|T - O\|_{\infty} + \min(|T - O|) + std(T - O), \quad (1)$$

where T and O represent the target and output vectors, respectively, of the training process. The absolute values of

the maximum and minimum error and standard deviation were used in the fitness function, which forced the algorithm to focus on reducing the error range, maximum, and minimum. A population size of 50 and 200 generations were used to predict the mesiodistal widths of the unerupted maxillary and mandibular canines and premolars. The GA tends to converge on a near-optimal solution through iterative use of three operators (selection, crossover, and mutation; Baese, 2004). Selection determines the survival of each population, based on the value of the fitness function, mutation arbitrarily alters one or more components of a selected chromosome, while crossover allows the search to fan out in diverse directions, and chromosomes of the best population members to be combined to produce the next generation. In the GA, the crossover fraction was set to 0.8.

The proposed algorithm was trained as follows. During each iteration, the GA introduced the reference teeth (by setting the corresponding values in the chromosomal binary vector to 1) into the ANN. The ANN tries to find the best mapping function for relating the reference inputs to the targets. Next, the algorithm checks the stopping criteria. If satisfied, the results are reported; if not, the GA moves to the next generation, searching for better possible candidates among the reference teeth. This process is repeated until the algorithm finds a result that satisfies the stopping criteria or until the number of generations exceeds the predefined value. This process is illustrated in Figure 1.

Due to the dimensions of the input vector, fivefold crossvalidation was used to avoid any bias or error during testing of the prediction efficiency of the proposed algorithm. This would provide an estimate of the generalization error through resampling the training and test samples, which were selected randomly.



Figure 1 Block diagram of the proposed technique.

To reduce the number of measurements used as input vectors, a linear inequality constraint for the GA was defined:

$$A \times V \le c \quad A = \begin{bmatrix} 11 \dots 1 \end{bmatrix}_{1 \times 10} \tag{2}$$

where V is the  $10 \times 1$  input binary vector. By changing the value of c, the maximum number of references (the number of chromosomes entries set to one) that the GA is allowed to select may be defined. For example, by setting c = 4, the GA was forced to use at most four reference values.

A multi layered perceptron neural network was used. The ANN had two hidden layers with eight and four neurons, respectively. The nodes of the first hidden layer had a tansigmoidal transfer function and those of the second hidden and output layers had purely linear transfer functions. The weights and biases of the network were initialized with random values. Batch training, where weights between the neurons were updated only after all the training samples were exposed to the network. The Levenberg–Marquardt method (Wu, 2008) was used for training, which gave the best possible results.

Once the training was completed, the GA was no longer used in the testing process. The reference teeth introduced by the GA as the best input candidates are used by the ANN for predicting the output through utilizing the best mapping function, developed during training.

All processing steps for the development of the hybrid GA–ANN algorithm were performed using matrix laboratory, version 7.8.0.347 (Matlab, MathWorks, Inc., Natick, Massachusetts, USA).

#### Regression analysis

The Kolmogrov–Smirnov test was used to confirm data normality. Pearson product-moment coefficients were used to estimate the correlation between the mesiodistal widths of the canines and premolars in the maxilla and mandible with the reference teeth (maxillary central incisors and first molars, mandibular incisors and first molars). Statistical calculations and analyses, including error variance of the estimates (mean square of error), correlation coefficients (r), coefficients of determination ( $r^2$ ), and analysis of variance (ANOVA) of the regression equations were performed using the Statistical Package for Social Sciences version 15.1 (SPSS Inc., Chicago, Illinois, USA). Statistical significance was established at the 0.05 level.

#### Results

#### Regression analyses

Table 1 shows the linear regression equations derived for predicting the mesiodistal widths of the mandibular canines and premolars using different reference teeth. The linear equations were obtained using the sum of the mesiodistal reference teeth as inputs.

The highest correlation in the mandible was between the sum of the mesiodistal widths of the canines and premolars with the mesiodistal widths of the mandibular first molars and central and lateral incisors (r = 0.697). The prediction errors (estimated as the sum of the mesiodistal widths of the canines and premolars on both sides of a dental arch) are indicated in Figure 2a.

The linear regression equations used to predict the mesiodistal widths of the maxillary canines and premolars are illustrated in Table 2. As in the previous case, the linear equations were obtained using the sum of the mesiodistal widths of the reference teeth as inputs. The highest correlation in the maxilla was between the sum of the mesiodistal widths of the canines and premolars with the mesiodistal widths of the mandibular first molars and maxillary central incisors (r = 0.742). The prediction errors are indicated in Figure 2b.

## Hybrid GA-ANN

The prediction error of the hybrid GA–ANN algorithm is illustrated in Figure 3. The reference teeth selected by the GA were the mandibular first molars, central and lateral incisors, and maxillary central incisors. The same references were used to predict the mesiodistal widths of both the mandibular and maxillary canines and premolars. The under/ overestimation rates of the real sizes of the maxillary canines and premolars were less than those of the mandible.

Table 3 shows comparison of the maximum errors in predicting the sizes of the mandibular and maxillary

 Table 1
 Regression equations and their parameters for predicting the mesiodistal widths of the canines and premolars in the mandible.

Regression	x	Correlation coefficient	Coefficients		MSE	95% CI of MSE	
1 			а	b			
Y = ax + b	Mesiodistal width of the mandibular first molars, central and lateral incisors	0.697	0.644	13.449	1.856	0 ± 2.65	
Y = ax + b	Mesiodistal width of the mandibular first molars and maxillary central incisors	0.658	0.658	16.353	2.0039	$0 \pm 2.76$	
Y = ax + b	Mesiodistal width of the mandibular and maxillary first molars	0.646	0.674	13.066	2.266	$0 \pm 2.93$	
Y = ax + b	Mesiodistal width of the mandibular central and lateral incisors and maxillary central incisors	0.577	0.492	22.609	2.39	$0\pm3.01$	

MSE, mean square of error; CI, confidence interval.

canines and premolars using linear regression analysis and the hybrid GA–ANN. The maximum prediction errors obtained using the hybrid GA–ANN were noticeably smaller than those obtained using linear regression. The prediction accuracy was higher in the maxilla than in the mandible when using the hybrid GA–ANN algorithm.

The maximum over/underestimation errors were plotted as a function of c (Figure 4). These are the errors in predicting the sizes of the maxillary canines and premolars for the same test samples (20 per cent of the total cases, selected randomly). For c = 4, the algorithm selected the maxillary and mandibular central incisors, and the mandibular first molars as reference teeth. When c = 3, the



**Figure 2** Prediction error in estimating the mesiodistal widths of the canines and premolars in the mandible (a) and maxilla (b) using linear regression and the reference teeth with the highest and second highest correlation in (a) Table 1 and (b) Table 2.

 Table 2
 Regression equations and their parameters for predicting the mesiodistal widths of the canines and premolars in the maxilla.

Regression	Correlation coefficient		Coefficients		MSE	95% CI of MSE
			а	b		
Y = ax + b $Y = ax + b$ $Y = ax + b$	Mesiodistal width of the mandibular first molars and maxillary central incisors Mesiodistal width of the mandibular first molars, central and lateral incisors Mesiodistal width of the mandibular central and lateral incisors and maxillary central	0.742 0.690 0.680	0.740 0.620 0.589	14.271 15.730 19.962	1.951 2.072 2.156	$0 \pm 2.72 \\ 0 \pm 2.80 \\ 0 \pm 2.86$
Y = ax + b	Incisors Mesiodistal width of the mandibular and maxillary first molars	0.624	0.634	15.990	2.493	$0\pm3.07$

MSE: mean square of error, CI: confidence interval.

selected references were the maxillary central incisors and mandibular first molars. When c = 2, the selected references were the maxillary central incisors. The absolute value of the rates of under/overestimation increased when the algorithm was forced to select fewer references.

### Discussion

Different racial and ethnic groups present variations in the mesiodistal widths of their permanent teeth (Bishara *et al.*, 1989; Schirmer and Wiltshire, 1997; Niwa, 2004; Zheng and Thomson, 2005; Fernández and Caballero, 2006). Thus, equations defined by data collected from one ethnic group to predict the size of unerupted permanent teeth might not be applicable to another (Al-Khadra, 1993; Yuen *et al.*, 1998). In contrast, the hybrid algorithm adjusts its structure based on the training samples presented to the system. Therefore, the hybrid GA–ANN algorithm can be used to predict the size of unerupted teeth in different ethnic groups, provided that an appropriate training data set is presented to the system.

An ideal prediction method should result in no difference between the predicted and actual widths of the permanent canines and premolars. However, prediction methods are not 100 per cent precise and can over- or underestimate the actual sizes of unerupted teeth (Fisk and Markin, 1979;



**Figure 3** Prediction error in estimating the mesiodistal widths of the canines and premolars in the mandible and maxilla using the hybrid genetic algorithms and artificial neural network.

Moyers, 1988). Overestimation by 1 mm of the actual widths of the permanent canines and premolars on each side of the arch is better than underestimation at preventing crowding and should not seriously affect an extraction decision (De Paula et al., 1995, Bernabé and Flores-Mir, 2005). Using the hybrid GA-ANN algorithm, the prediction error for the maxillary unerupted teeth was between -1 and 0 mm a greater percentage of the time than the prediction error for the mandible. Thus, when using the hybrid GA-ANN algorithm, the prediction accuracy was higher in the maxilla than in the mandible. For both the maxilla and mandible, the prediction error was between -1 and 0 mm a greater percentage of the time when using the GA-ANN algorithm than when regression equations were used. This indicates that the hybrid GA-ANN algorithm provided more accurate results than the most correlated regression equations. In both the mandible and maxilla, the percentage of prediction errors >1.5 mm or <-1.5 mm was noticeably smaller using the hybrid GA-ANN algorithm than when regression equations were used. This is because the fitness function in equation 1 was defined to minimize the maximum absolute error value.

During regression analysis, the sizes of the mandibular first molars and maxillary central incisors were strongly correlated with those of the maxillary canines and premolars (r = 0.742). In the mandible, the highest correlation was observed between the sizes of the mandibular first molar and incisors with the sizes of the canines and premolars (r = 0.697).



**Figure 4** Maximum error in predicting the sizes of the canines and premolars in the maxilla as a function of c.

Table 3	Maximum	errors in	predicting	the sizes	of the	canines and	l premolars	in the	e mandible a	ind maxilla
---------	---------	-----------	------------	-----------	--------	-------------	-------------	--------	--------------	-------------

	Linear regression		Linear regression		Hybrid GA–ANN			
	References: mesioc mandibular first mo central incisors	listal widths of the plars and maxillary	References: mesioo mandibular first mo lateral incisors	listal widths of the olars, central and	References: mesiodistal widths of mandibular first molars, central and lateral incisors, and maxillary central incisors			
	Overestimating	Underestimating	Overestimating	Underestimating	Overestimating	Underestimating		
Mandible Maxilla	-3.30 -3.93	3.97 383	-3.84 -3.19	3.78 3.84	-2.37 -1.53	2.0 1.56		

Genetic algorithm-artificial neural network (GA-ANN).



Figure 5 The main page of the developed graphical user interface.

The GA–ANN selected the mandibular first molars and incisors, and maxillary central incisors as the reference teeth for predicting the sizes of the unerupted mandibular and maxillary canines and premolars.

Reducing the value of c, which limited the number of references that the GA–ANN algorithm could choose, increased the maximum error values. However, the maximum error was not large compared with errors obtained using the linear regression equations. For c = 2, the algorithm identified the mandibular first molar as the most suitable reference tooth for predicting the sizes of the canines and premolars. This agreed well with regression analysis between the maxillary central incisors and canines and premolars was 0.697. This was also the highest correlation observed when using a single reference for prediction.

The data were prepared by manual measurements. For ease of use of the introduced algorithm, a graphical user interface was developed (Figure 5). The system parameters vary depending on whether the prediction was for unerupted teeth in the mandible or maxilla; thus, the user first selects the aim of the prediction (i.e. mandible or maxilla) then uploads a digital photograph of the corresponding cast. The measurement system is calibrated by clicking on a 10 mm line (e.g. a ruler) three times, and the average value of the selected distance is recorded as the calibration basis. The user then clicks three times on the two ends of the predefined teeth in the captured cast image. The recorded measurements are fed into the algorithm software, and the results are provided in both graphical and numerical form. In the calibration phase, the horizontal or vertical distance between the selected points is recorded (based on the positioning of the calibration element), while in the measurement phase the Euclidean distance is calculated since the teeth may have many different directions. As the static occlusal view may not be efficient for finding the mesiodistal width of tilted teeth, digital photographs cannot be useful for predicting the width of the unerupted teeth when measurements need to be carried out on such teeth.

This study has several limitations. Due to the regional nature, the proposed prediction method was only tested in one ethnic group. More generalized studies in different ethnic groups are needed to validate the feasibility of the proposed method.

Multivariable regression could provide more accurate results compared with the single variable regression technique adopted in this study. However, these equations are complex and difficult to memorize, justifying the use of the simple equations adopted (Staley *et al.*, 1984; Nourallah *et al.*, 2002; Legović *et al.*, 2003; Bernabé and Flores-Mir, 2005).

#### Conclusions

The proposed technique is a promising tool for predicting the sizes of unerupted canines and premolars with greater accuracy than linear regression analyses. Most notably, the structure of the technique can be adjusted based on the data collected from different ethnic groups.

#### Acknowledgements

We would like to thank the vice president of the research center of Mashhad University of Medical Science for financial support.

#### References

- Al-Khadra B H 1993 Prediction of the size of unerupted canines and premolars in a Saudi Arab population. American Journal of Orthodontics and Dentofacial Orthopedics 104: 369–372
- Baese A M 2004 Pattern recognition for medical imaging. Elsevier Academic Press, London
- Bernabé E, Flores-Mir C 2005 Are the lower incisors the best predictors for the unerupted canines and premolars sums? An analysis of a Peruvian samples. Angle Orthodontist 75: 198–203
- Bishara S E, Jakobsen J R, Abdallah E M, Fernandez Garcia A 1989 Comparisons of mesiodistal and buccolingual crown dimensions of the permanent teeth in three populations from Egypt, Mexico and the United States. American Journal of Orthodontics and Dentofacial Orthopedics 96: 416–422
- De Paula S, Almeida A O, Lee P C 1995 Prediction of mesiodistal diameter of unerupted lower canines and premolar using 45° cephalometric radiography. American Journal of Orthodontics and Dentofacial Orthopedics 107: 309–313
- Devito K L, Barbosa F D S, Filho W N F 2008 An artificial multilayer perceptron neural network for diagnosis of proximal dental caries. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology 106: 879–884
- Diagne F, Diop-Ba K, Ngom P L, Mbow K 2003 Mixed dentition analysis in a Senegalese population: elaboration of prediction tables. American Journal of Orthodontics and Dentofacial Orthopedics 124: 178–183
- Ferguson F S, Mucko D J, Sonnenberg E M, Shakun M L 1978 The use of regression constants in estimating tooth size in a Negro population. American Journal of Orthodontics 73: 68–72
- Fernández M, Caballero J 2006 Ensembles of Bayesian-regularized genetic neural networks for modeling of acetylcholinesterase inhibition by huprines. Chemical Biology and Drug Design 68: 201–212
- Fisk R O, Markin S 1979 Limitations of the mixed dentition analysis. Ontario Dentist 56: 16–20
- Gardner R B 1979 A comparison of four methods of predicting arch length. American Journal of Orthodontics 75: 387–398

- Hixon E H, Oldfather R E 1958 Estimation of the size of unerupted cuspid and bicuspid teeth. Angle Orthodontist 28: 236–240
- Irwin R D, Herold J S, Richardson A 1995 Mixed dentition analysis: a review of methods and their accuracy. International Journal of Pediatric Dentistry 5: 137–142
- Jaroontham J, Godfrey K 2000 Mixed dentition space analysis in a Thai population. European Journal of Orthodontics 22: 127–134
- Kirschen R H, O'Higgins E A, Lee R T 2000 The Royal London Space Planning: an integration of space analysis and treatment planning: Part I: assessing the space required to the treatment objectives. American Journal of Orthodontics and Dentofacial Orthopedics 118: 448–455
- Lee-Chan S, Jacobson B N, Chwa K H, Jacobson R S 1998 Mixed dentition analysis for Asian Americans. American Journal of Orthodontics and Dentofacial Orthopedics 113: 293–299
- Legović M, Novosel A, Legović A 2003 Regression equations for determining mesiodistal crown diameters of canines and premolars. Angle Orthodontist 73: 314–318
- Lo S B, Lin J, Freedman M T, Mun S K 1998 Application of artificial neural networks to medical image pattern recognition: detection of clustered microcalcifications on mammograms and lung cancer on chest radiographs. Journal of VLSI Signal Processing Systems for Signal, Image, and Video Technology 18: 263–274
- Meersman D, Scheunders P, VanDyck D 1996 Detection of microcalcifications using neural networks. Digital Mammography 96: 287–290
- Moyers R E 1988 Handbook of orthodontics, 4th edn. Year Book Medical Publishers Inc., Chicago

- Ngan S, Hu X 1999 Analysis of functional imaging data using selforganizing mapping with spatial connectivity. Magnetic Resonance in Medicine 41: 939–946
- Niwa T 2004 Prediction of biological targets using probabilistic neural networks and atom-type descriptors. Journal of Medicinal Chemistry 47: 2645–2650
- Nourallah A W, Gesch D, Khordaji M N, Splieth C 2002 New regression equation for predicting the size of unerupted canines and premolars in a contemporary population. Angle Orthodontist 72: 210–220
- Schirmer U R, Wiltshire W A 1997 Orthodontic probability tables for black patients of African descent: mixed dentition analysis. American Journal of Orthodontics and Dentofacial Orthopedics 112: 545–551
- Staley R N, O'Gorman T W, Hoag J F, Shelly T H 1984 Prediction of the widths of unerupted canines and premolars. Journal of the American Dental Association 108: 185–190
- Tanaka M M, Johnston L E 1974 The prediction of the size of unerupted canines and premolars in a contemporary orthodontic population. Journal of the American Dental Association 88: 798–801
- Wu J M 2008 Multilayer Potts perceptrons with Levenberg-Marquardt learning. IEEE Transactions on Neural Networks 19: 2032–2043
- Yuen K K, Tang E L, So L L 1998 Mixed dentition analysis for Hong-Kong Chinese. Angle Orthodontist 68: 21–28
- Zheng R Y, Thomson R 2005 Bio-basis function neural network for prediction of protease cleavage sites in proteins. IEEE Transactions on Neural Networks 16: 263–274

Copyright of European Journal of Orthodontics is the property of Oxford University Press / USA and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.