# Survival of *Enterococcus faecalis* in root canals *ex vivo*

# C. M. Sedgley, S. L. Lennan & O. K. Appelbe

Department of Cariology, Restorative Sciences and Endodontics, School of Dentistry, University of Michigan, Ann Arbor, MI, USA

### Abstract

Sedgley CM, Lennan SL, Appelbe OK. Survival of *Enterococcus faecalis* in root canals *ex vivo*. *International Endodontic Journal*, **38**, 735–742, 2005.

**Aim** The hypotheses tested in this study were that: (i) *Enterococcus faecalis* can survive long-term entombment in root filled teeth without additional nutrients, (ii) initial cell density influences the survival of *E. faecalis* in instrumented root canals and (iii) gelatinase-production capacity influences the survival of *E. faecalis* in root canals.

**Methodology** The root canals of 150 extracted single canal teeth were instrumented to apical size 60 and divided into six groups of 25. Within each group 10 canals were inoculated with either gelatinase-producing *E. faecalis* OG1-S and the other 10 with its gelatinase-defective mutant *E. faecalis* OG1-X. Five canals per group were kept as uninoculated controls. The root canals in groups 1 and 2 were inoculated with  $10^6$  bacteria, incubated for 48 h at 37 °C then filled with gutta-percha and zinc-oxide eugenol sealer. Root canals were inoculated with  $10^6$ ,  $10^5$ ,  $10^4$  and  $10^3$  bacteria in groups 3–6, respectively, and left unfilled. All teeth were sealed coronally with glass–ionomer

cement. After 6- (groups 1, 3–6) and 12-month (group 2) incubation at 37 °C in 100% humidity, root fragments were analysed for presence of *E. faecalis*, using culture, polymerase chain reaction and histological methods.

**Results** Viable *E. faecalis* was recovered from all root filled teeth and from 95-100% of unfilled inoculated teeth. Initial cell density and gelatinase production did not influence the recovery of viable *E. faecalis* (P > 0.05; chi-square test). *Enterococcus faecalis* 16S rRNA gene products were present in all inoculated teeth and absent in all noninoculated controls. Dentinal tubule infection was evident under light microscopy in sections from inoculated teeth after 48-h, 6- and 12-month incubation.

**Conclusions** *Enterococcus faecalis* inoculated into root canals maintained viability for 12-months *ex vivo*. The clinical implications are that viable *E. faecalis* entombed at the time of root filling could provide a long-term nidus for subsequent infection.

**Keywords:** *Enterococcus faecalis, in vitro,* root canal, root filling, survival.

Received 17 January 2005; accepted 16 May 2005

#### Introduction

Studies on primary root canal treatment have indicated that a negative culture at the time of filling is associated with a better prognosis (Byström *et al.* 1987, Sjögren *et al.* 1997). Sjögren *et al.* (1997) reported that the presence of bacteria at filling was associated with the presence of apical periodontitis 5 years after treatment in 32% of the cases. Persistent radiolucent lesions have been associated with bacteria in root filled, asymptomatic teeth with periapical lesions (Nair *et al.* 1990). However, the difficulty in rendering canals bacteria-free *in vivo* has been well-documented, regardless of instrumentation method and irrigation regimen (Byström & Sundqvist 1985, Card *et al.* 2002).

*Enterococcus faecalis* has long been implicated in persistent root canal infections (Engström 1964) and more recently has been identified as the species most

Correspondence: Dr Christine Sedgley, Department of Cariology, Restorative Sciences and Endodontics, School of Dentistry, University of Michigan, 1011 N. University Drive, Ann Arbor, MI 48109-1078, USA (Tel.: +1 734 647 4182; fax: +1 734 936 1597; e-mail: csedgley@umich.edu).

commonly recovered from root canals of teeth with post-treatment disease (Sundqvist *et al.* 1998, Siqueira & Rôças 2004). Three of nine cases from which *E. faecalis* was recovered at the time of retreatment failed to heal after 5 years (Sundqvist *et al.* 1998). The recovery of *E. faecalis* from failed previously filled root canals does not seem to be related to the use of a particular root filling material (Peciuliene *et al.* 2001). Enterococci might be selected in root canals undergoing standard endodontic treatment because of low sensitivity to antimicrobial agents (Dahlén *et al.* 2000) or their ability to inactivate antimicrobial agents (Portenier *et al.* 2002).

Siqueira Jr (2001) speculated that bacteria entombed by the root filling material usually die, but that some entombed species 'probably survive for relatively long periods', deriving nutrients from tissue remnants. The ability of *E. faecalis* to survive 'entombed' in root canals over an extended period has never been measured. This study tested the hypotheses that: (i) *E. faecalis* can survive 'entombment' in filled root canals, (ii) initial cell density influences the survival of *E. faecalis* in instrumented root canals and (iii) gelatinase-production capacity influences the survival of *E. faecalis* in root canals.

# **Materials and methods**

#### Microorganisms

Two isogenic *E. faecalis* strains were used in this study: the gelatinase-producing, streptomycin resistant, *E. faecalis* OG1-S and its reduced gelatinase-producing mutant derivative, *E. faecalis* OG1-X.

*Enterococcus faecalis* strains were stored at -80 °C. When required, each strain was inoculated onto Todd–Hewitt broth (THB) (Difco, Becton Dickinson and Company, Sparks, MD, USA) supplemented with 1.5% agar and incubated aerobically at 37 °C for 24–48 h. One colony-forming unit was transferred to

Table 1	1 Stud	ly design
---------	--------	-----------

5 mL THB and grown in aerobic incubation for 24 h at 37 °C. Bacterial suspensions (50 µL) were transferred to 5 mL fresh THB and grown to late exponential phase at which point cells were harvested. Cultures were centrifuged at 1500 q for 10 min. Deposits were washed twice in phosphate-buffered saline  $(0.1 \text{ mol } L^{-1}, \text{ pH } 7.3)$  and reconstituted in THB to a final concentration of approximately  $1 \times 10^9$  microorganisms per mL, corresponding to OD 0.75 at 660 nm (Spectronic 20D+; Thermo Electron Corporation, Pittsburgh, PA, USA). Serial dilutions (1:10) corresponding to  $10^8$ ,  $10^7$  and  $10^6$  cells mL<sup>-1</sup> were prepared in THB. Calibration between viable counts and suspension optical density was confirmed by spiral plating serial dilutions to THB agar in triplicate and enumerating after 48 h aerobic incubation at 37 °C.

### Selection and instrumentation of teeth

Permanent anterior teeth (n = 152) with single straight roots, clinically intact crowns, no restorations, no cracks and no caries were selected from the collection of extracted teeth stored in 50% glycerine/ 50% ethyl alcohol at the School of Dentistry, University of Michigan, Ann Arbor, MI, USA. Teeth were autoclaved at 121 °C for 15 min at 26 psi and stored at 25 °C in 100% humidity until use. Two teeth were used for histological observations described below. The remaining 150 teeth were divided into six groups (Table 1). Root canals were instrumented to an apical size 60 at working length (WL) as previously described (Sedgley et al. 2004a). Access openings were prepared using tungsten carbide burs in a high-speed handpiece with constant water spray. WL was set at 1 mm short of the apical foramen by visual inspection of a size 15 stainless steel K-file extending beyond the apical foramen. Canals were instrumented using a crowndown method and nickel-titanium (Ni-Ti) rotary and hand instruments (Dentsply; Tulsa Dental Products,

	Filled canals		Unfilled canals			
Incubation period (month): Inoculum (cells per 15 µL):	Group 1 6 10 <sup>6</sup> cells	Group 2 12 10 <sup>6</sup> cells	Group 3 6 10 <sup>6</sup> cells	Group 4 6 10 <sup>5</sup> cells	Group 5 6 10 <sup>4</sup> cells	Group 6 6 10 <sup>3</sup> cells
Experimental						
Enterococcus faecalis OG1-S	10	10	10	10	10	10
Enterococcus faecalis OG1-X	10	10	10	10	10	10
Sterility controls						
No inoculum	5	5	5	5	5	5

736

Copious irrigation with 5.25% sodium hypochlorite was used between instruments. A 0.012 mm diameter round diamond bur in a high-speed handpiece was used to make 0.5 mm deep grooves on the mesial and distal aspects of each tooth, avoiding the apical 2 mm of narrow roots in order to not penetrate into the root canal; the purpose of this step was to facilitate subsequent splitting of the tooth at the completion of the incubation period.

For each tooth in groups 1 and 2, a master guttapercha cone (Greater Taper Gutta-Percha; Analytic Endodontics, Glendora, CA, USA) was custom-fitted to WL and indented at the level of the coronal reference point. Each master gutta-percha cone was transferred to a labelled, empty 0.6 mL polymerase chain reaction (PCR) tube with attached cap (Dot Scientific Inc., Burton, MI, USA). In addition, nonstandardized guttapercha cones (Fine-Fine; Kerr, Romulus, MI, USA), for use as accessory cones, were sealed in PCR tubes. Gutta-percha cones were sterilized by  $\gamma$ -radiation at the Ford Nuclear Reactor Phoenix Memorial Laboratory (University of Michigan).

The smear layer was removed from canal walls by ultrasonic treatment for 4 min each in 17% EDTA and 5.25% NaOCl (Haapasalo & Ørstavik 1987) after which canals were copiously rinsed with distilled water and dried with paper points. Root apices were coated with nail varnish to seal the apical foramen. Teeth were inserted with crown exposed into individual compartments in polypropylene boxes and autoclaved at 121 °C for 15 min at 26 psi, then aseptically stored in 100% humidity at 25 °C until use.

# Inoculation and filling of root canals (groups 1 and 2)

All following procedures were performed under strictly aseptic conditions. Immediately before inoculation with bacteria, canals were dried with sterile paper points (Kerr). The root canals of teeth in groups 1 and 2 were inoculated with  $10^6$  cells in 15 µL THB and aerobically incubated at 37 °C for 48 h to allow penetration of *E. faecalis* into root canal dentine. Immediately prior to filling, canals were dried with paper points (Kerr) placed to WL. Canals were filled to WL by an endodontist using cold lateral compaction with guttapercha and previously unopened sealer (Roth-801;

Roth International Ltd, Chicago, IL, USA). The sealer was mixed according to the manufacturer's instructions on a sterile slab using a sterile spatula. The sealer was introduced into the canal to WL using a size 60 stainless steel K-file that was rotated counterclockwise as it was withdrawn. The apical half of each prefitted custom gutta-percha cone was coated lightly with sealer and placed to the WL. Cold lateral compaction was performed using a sterile D11T hand spreader (Kerr) and nonstandardized gutta-percha cones. Excess coronal gutta-percha was seared off at the level of the cemento-enamel junction using the System B heat source (SybronEndo, Orange, CA, USA) at 300 °C.

#### Inoculation of root canals (groups 3-6)

Each root canal was inoculated with 15  $\mu$ L of microbial suspension. Canals in groups 3–6 were inoculated with  $10^6$ ,  $10^5$ ,  $10^4$  and  $10^3$  cells, respectively (Table 1).

#### Restoration of access cavities

Access cavities were sealed with glass–ionomer cement (Ketac-Fil Plus; 3M-ESPE, AG, Seefeld, Germany) immediately following filling for groups 1 and 2, and after inoculation for teeth in groups 3–6. This material was selected based on previous reports of its sealability over long periods of time (Barthel *et al.* 1999). *In vitro* agar diffusion and broth dilution studies showed that Ketac-Fil Plus did not have bactericidal activity against *E. faecalis* OG1-S and *E. faecalis* OG1-X over concentrations ranging from  $10^1$  to  $10^9$  cells mL<sup>-1</sup> (data not shown).

#### Incubation of teeth

Each tooth was transferred to an empty labelled screwtop 2 mL sterile cryogenic vial (Corning Costar Corporation, Cambridge, MA, USA) and the cap replaced loosely in order to allow maintenance of humidity. Vials were placed in individual compartments in sterile polypropylene boxes containing sterile water at their base. Boxes were sealed hermetically and placed in an incubator at 37 °C under aerobic conditions.

# Splitting teeth and assessment of viability of microorganism in teeth

At the end of each incubation period, each tooth was aseptically transferred from its individual vial to the base of a sterile foil-covered Teflon beaker (Nalgene Labware, Rochester, NY, USA). Teeth were aseptically split into fragments using sterile long-handled endcutter pliers (Channellock Inc., Meadville, PA, USA). Care was taken not to introduce any contaminants by keeping the beaker covered during splitting. All inoculated and uninoculated sterility control teeth were split using this method. Tooth fragments were aseptically transferred for distribution among (i) two 10 mL test tubes containing 5 mL THB broth, (ii) two 10 mL test tubes containing 4 mL THB broth supplemented with 1 mL human serum (Sigma-Aldrich, St Louis, MO, USA) and (iii) bile esculin agar (BEA: Difco). The contents of each tube were vortexed very vigorously to enable displacement of cells from tooth structure into suspension. Pairs of test tubes containing tooth fragments in THB alone and in THB/serum were incubated at 37 °C for up to 72 h under aerobic and anaerobic conditions (candle jar). Growth in broth was monitored by measuring optical density at 660 nm. BEA agar was incubated aerobically at 37 °C. After 72 h each tooth was scored as positive or negative for evidence of viable E. faecalis as determined by growth under any condition. Confirmation of E. faecalis OG1-S and OG1-X was made by culturing a loopful of suspension from each positive broth culture onto THB agar supplemented with 500  $\mu$ g mL<sup>-1</sup> streptomycin, BEA and 3% gelatin agar which were aerobically incubated at 37 °C for up to 72 h.

# Presence of *E. faecalis* by 16S rRNA gene using polymerase chain reaction

For each tooth sample in groups 1 and 2, the presence of *E. faecalis* in 1.5 mL vortexed suspensions of tooth fragments in THB was assessed by 16S rRNA gene product detection using end-point PCR and by investigators blinded to the culture results. These steps were not to confirm viability but to determine if the method had allowed displacement of cellular material into the suspension and to check sterility controls.

Methods have been previously described for total DNA isolation, development of universal bacteria and *E. faecalis*-specific 16S rRNA gene primer pairs and analyses of PCR products (Sedgley *et al.* 2005a). PCR amplifications were prepared in a 30  $\mu$ L final reaction volume with the following: 100 ng total DNA template; 6 pmol of each primer; 100  $\mu$ mol L<sup>-1</sup> dNTPs (Invitrogen, Carlsbad, CA, USA); 1.5 U HotStarTaq DNA polymerase (Qiagen, Valencia, CA, USA); 3  $\mu$ L 10X HotStarTaq DNA polymerase PCR buffer (Qiagen). PCR conditions were: 15 min initial enzyme activation/DNA

denaturation step at 95 °C followed by 40 consecutive cycles at 94 °C for 20 s, 66 °C for 45 s and 72 °C for 15 s using an Eppendorf Mastercycler (Brinkmann Instruments Inc., Westbury, NY, USA).

#### Histology

The development of tubule infection after 48-h, 6- and 12-month incubation was evaluated by light microscopy. For the 48-h evaluation, two instrumented teeth were inoculated with  $10^6$  cells in 15 µL THB of either *E. faecalis* OG1-S or *E. faecalis* OG1-X and incubated as previously described then split. For groups 1–6, tooth fragments were obtained from randomly selected teeth. Fragments were immersed in 10% neutral buffered formalin for 24 h and demineralized in 0.5 mol L<sup>-1</sup> ethylene diamine tetra-acetic acid (pH 7.3) paraffin-embedded, sectioned and Brown and Brenn stained. Images of sections were captured with the Olympus BX-51 microscope and DP-70 digital camera and software (Olympus, Melville, NY, USA).

## Data analysis

Each tooth was scored as either positive or negative for viable *E. faecalis* under any of the growth conditions. Chi-square tests were used to compare the number of teeth positive for growth after (i) 6- and 12-month filling (groups 1 and 2), (ii) 6-month incubation with either  $10^6$ ,  $10^5$ ,  $10^4$ , or  $10^3$  bacteria (groups 3–6) and (iii) inoculation with the gelatinase-producing *E. faecalis* OG1-S and the reduced gelatinase-producing *E. faecalis* OG1-X. Significance was set at P < 0.05.

# Results

After 6 (group 1) and 12 months (group 2) incubation viable *E. faecalis* was recovered from all filled experimental teeth under all growth conditions (P > 0.05) and never from any noninoculated sterility control teeth. PCR results showed that *E. faecalis* 16S rRNA gene products were absent in all noninoculated controls and were present in all inoculated teeth. Growth of all recovered *E. faecalis* OG1-S and *E. faecalis* OG1-X was confirmed on selective media.

Initial cell count did not influence recovery (P > 0.05). Viable *E. faecalis* was recovered from 20, 19, 19 and 20 teeth in groups 3, 4, 5 and 6 respectively under at least one growth condition (Table 2). There was no difference in the number of

	Aerobic Anaero			robic		
<i>Enterococcus</i> <i>faecalis</i> strain	BEA	тнв	THB- serum	тнв	THB- serum	Overall group positive
Group 3						
OG1-S	10	10	10	10	10	10
OG1-X	9	10	9	9	9	10
Group 4						
OG1-S	10	10	10	9	10	10
OG1-X	7	9	9	9	9	9
Group 5						
OG1-S	9	9	9	9	9	9
OG1-X	7	7	10	7	8	10
Group 6						
OG1-S	10	10	10	10	10	10
OG1-X	9	10	10	10	10	10
Total	71	75	77	73	75	78

*Enterococcus faecalis* OG1-S, gelatinase producing; OG1-X, reduced gelatinase producing mutant derivative of OG1-S; BEA, bile esculin agar; THB, Todd–Hewitt broth. Canals were inoculated with  $10^6$  cells in group 3,  $10^5$  cells in group 4,  $10^4$  cells in group 5 and  $10^3$  cells in group 6.

teeth with viable *E. faecalis* OG1-S compared to *E. faecalis* OG1-X (P > 0.05) indicating that the capacity for gelatinase production did not influence the recovery of viable *E. faecalis*.

Dentinal tubule infection was evident under light microscopy in sections from inoculated teeth after 48-h, 6- and 12-month incubation. Representative images obtained after 48-h and 12-month incubation are shown in Fig. 1. Bacteria were seen in the dentinal tubules on the root canal side only and not from the cementum side.

#### Discussion

The present in vitro investigations showed that E. faecalis introduced into root canals maintained viability for 12 months without additional nutrients. Others have reported that E. faecalis has the capacity to survive under various environmental stresses, for example, when exposed to salt, bile salts, acid, heat (Flahaut et al. 1996), glucose starvation (Capiaux et al. 2000), NaOCl (Laplace et al. 1997) and starvation in tap water (Hartke et al. 1998). In contrast, there is less known about the survival of E. faecalis over an extended period. E. faecalis was capable of surviving more than 90 days on commonly used hospital fabrics and plastics (Neely & Maley 2000) and for up to 4 months in water (Figdor et al. 2003). Other in vitro studies have shown that E. faecalis is capable of entering and recovering from the viable but nonculturable (VBNC) state, a survival strategy adopted by bacteria when exposed to environmental stress (Lleò et al. 2001). VBNC E. faecalis displayed cell wall alterations that might provide protection under unfavourable environmental conditions (Signoretto et al. 2000) and maintained adhesive properties to cultured human cells (Pruzzo et al. 2002).

Sundqvist *et al.* (1998) speculated that *E. faecalis* 'can enter the canal, survive the antibacterial treat-

International Endodontic Journal, 38, 735-742, 2005



**Figure 1** Infection of dentinal tubules by *Enterococcus faecalis* OG1-S after 48-h incubation (a and inset in b) and after root canal filling and 12-month incubation (c and inset in d). Brown and Brenn stain.

ment and then persist after obturation'. However, there are few data on long-term survival of *E. faecalis* in association with endodontic treatment. *E. faecalis* was able to survive in infected bovine dentine in water and in calcium hydroxide for at least 10 days (Ørstavik & Haapasalo 1990). The ability of *E. faecalis* to survive in the root canal *in vivo* is restricted to the classic monkey study by Fabricius *et al.* (1982) in which it was shown that viable *E. faecalis* was recovered from all nine (unobturated) root canals 6 months after inoculation with approximately  $10^5$  cells. In the present study, recovery of viable cells after 6 months was evident after an initial inoculum of  $10^3$  cells.

Anaerobic or aerobic conditions did not influence recovery of the facultatively anaerobic *E. faecalis* species in agreement with previous data (Figdor *et al.* 2003). In addition to oxygen conditions, other factors that did not influence recovery of *E. faecalis* were the gelatinase capacity of the *E. faecalis* strain and initial cell density. Gelatinases are extracellular zinc endopeptidases capable of hydrolyzing gelatin, collagen and other peptides that were produced by 70% of *E. faecalis* strains isolated from root canals (Sedgley *et al.* 2005b). Expression of GelE might contribute to increased dissemination of *E. faecalis* in high-density environments (Waters *et al.* 2003).

Recovery of E. faecalis from tooth fragments was also not influenced by the addition of human serum to growth media after splitting teeth. The addition of human serum was investigated because fluid resembling serum in the periapical tissues could potentially supply nutrients to support microorganisms left behind in or near dentine; starved E. faecalis cells recovered better in media supplemented with human serum (Figdor et al. 2003). Serum enhanced E. faecalis biofilm formation on gutta-percha (Takemura et al. 2004). In the present study, the addition of 20% human serum to THB was not essential for recovery. However, there was a high survival rate of bacteria over all conditions. Thus, whether serum enhances survival of starved E. faecalis would need to be tested under more challenging conditions than those in the present study, including longer incubation periods, less dense initial cell inocula, thermal obturation methods and the use of different sealers. The influence of other products potentially found in the periapical region on survival of starved E. faecalis could also be investigated.

In this study, bacterial suspensions were inoculated directly into the root canal rather than immersing root samples into a liquid culture and allowing growth into dentinal tubules from both root canal and cementum sides (Haapasalo & Ørstavik 1987). It is important to note that, using the present method, bacteria could have entered lateral canals and grown and survived without nutrients on the root surface, although this was not observed histologically in 48-h, 6- and 12-month specimens. From a clinical perspective, recesses and diverticula inaccessible to biomechanical instrumentation can provide a niche for microorganisms, mostly as biofilms (Nair et al. 2005). The incubation period of 48 h in this study was intentionally shorter than the 2-week period used to obtain a wellinfected dentine sample for studying the bactericidal action of endodontic materials (Haapasalo & Ørstavik 1987). Penetration of E. faecalis to a depth of 300-400 µm in bovine dentinal tubules was evident after 1 day (Haapasalo & Ørstavik 1987), similarly observed after 48 h in this study (Fig. 1). As opposed to testing the viability of large numbers of bacteria after exposure to potential antimicrobial agents, this study sought to determine if any cell would recover after a period of starvation and no attempts were made to quantify the number of surviving bacteria. In vivo, the number of cells required to initiate and sustain the periapical infection remains to be determined, but would be also dependent on the location of the cells and the host response.

The survival of E. faecalis after 'entombment' by filling reported here supports data from clinical studies (Sundqvist et al. 1998, Siqueira & Rôças 2004). However, with reference to the use of a zinc-oxide eugenol-based sealer (Roth), the present data contrast with those obtained using a dentine block model infected with E. faecalis (Saleh et al. 2004). Saleh et al. (2004) reported that Grossman's sealer (also a zincoxide eugenol-based sealer) killed all E. faecalis within 7 days in samples of circumpulpal dentine collected from the inner 300  $\mu$ m of the canal; deeper samples were not obtained. In contrast, in this study, sampling was not limited to any part of the tooth. In the event of leakage alongside the root canal filling material, logically the initial source of bacteria would be closest to the root canal. It has been observed that invasion of bacteria into the root canals varies in terms of depth of penetration and not all tubules are equally invaded (Peters et al. 2001, Haapasalo et al. 2003). These observations were also borne out in light microscopy examination of infected sections that showed some dentinal tubules heavily invaded while adjacent tubules were empty (Fig. 1). The primary purpose of histological examination was to confirm the presence of bacteria in dentinal tubules at the three time periods.

740

There was no evidence of an increase in bacterial numbers over time. Interestingly, bacteria seemed to stain more weakly after the starvation periods. Whether these observations can be attributed to cell wall changes as a result of starvation (Signoretto *et al.* 2000) could be addressed in investigations using more sophisticated microscopic methods.

The original source of E. faecalis recovered from previously filled root canals undergoing retreatment is uncertain. Clinical studies have shown that E. faecalis usually make up a small proportion of the flora in the untreated root canal (Byström & Sundqvist 1985, Sjögren et al. 1997). One source of E. faecalis infecting the filled root canal might be the oral cavity (Rams et al. 1992, Sedgley et al. in press), potentially via coronal leakage after root canal filling. Alternatively, E. faecalis strains might be introduced into the root canal as contaminants during endodontic treatment procedures and bacteria left in dentinal tubules might be a reservoir from which canal reinfection can occur (Oguntebi 1994). Peters et al. (1995) stated that a sound obturation technique immediately after the cleaning, shaping and disinfection phases allows the remaining bacteria in the tubules to be either inactivated or prevented from repopulating the voids in the (former) canal space. They speculated that in the vast majority of cases, those bacteria appear not to jeopardize the successful outcome of root canal treatment. The present in vitro data indicate that E. faecalis has the capacity to recover from a prolonged starvation state in root filled teeth. This capacity could provide E. faecalis with an ecological advantage over other less hardy species that might be remaining in the filled root canal and thereby explain its recovery from intact previously treated root filled teeth (Sjögren et al. 1997). The clinical implications are that viable E. faecalis entombed at the time of root filling could provide a long-term nidus for subsequent infection if the opportunity arises.

#### Acknowledgements

The authors gratefully acknowledge the Endodontic Research Fund, University of Michigan for financial support and the Michigan Memorial Phoenix Project for gamma radiation sterilization of endodontic materials. The authors thank John Westman and Chris Edwards from the University of Michigan for their expert technical assistance with histology.

#### References

- Barthel CR, Strobach A, Briedigkeit H, Gobel UB, Roulet JF (1999) Leakage in roots coronally sealed with different temporary fillings. *Journal of Endodontics* 25, 731–4.
- Byström A, Sundqvist G (1985) The antibacterial action of sodium hypochlorite and EDTA in 60 cases of endodontic therapy. *International Endodontic Journal* **18**, 35–40.
- Byström A, Happonen R-P, Sjögren U, Sundqvist G (1987) Healing of periapical lesions of pulpless teeth after endodontic treatment with controlled asepsis. *Endodontics and Dental Traumatology* **3**, 58–63.
- Capiaux H, Giard JC, Lemarinier S, Auffray Y (2000) Characterization and analysis of a new gene involved in glucose starvation response in *Enterococcus faecalis*. *International Journal of Food Microbiology* **55**, 99–102.
- Card SJ, Sigurdsson A, Ørstavik D, Trope M (2002) The effectiveness of increased apical enlargement in reducing intracanal bacteria. *Journal of Endodontics* 28, 779–83.
- Dahlén G, Samuelsson W, Molander A, Reit C (2000) Identification and antimicrobial susceptibility of enterococci isolated from the root canal. *Oral Microbiology and Immunology* 15, 309–12.
- Engström B (1964) The significance of enterococci in root canal treatment. *Odontologisk Revy* **15**, 87–106.
- Fabricius L, Dahlén G, Ohman AE, Möller AJR (1982) Predominant indigenous oral bacteria isolated from infected root canals after varied times of closure. *Scandinavian Journal* of Dental Research **90**, 134–44.
- Figdor D, Davies JK, Sundqvist G (2003) Starvation survival, growth and recovery of *Enterococcus faecalis* in human serum. Oral Microbiology and Immunology **18**, 234–9.
- Flahaut S, Benachour A, Giard JC, Boutibonnes P, Auffray Y (1996) Defense against lethal treatments and *de novo* protein synthesis induced by NaCl in *Enterococcus faecalis* ATCC 19433. Archives of Microbiology 165, 317–24.
- Haapasalo M, Ørstavik D (1987) In vitro infection and disinfection of dentinal tubules. Journal of Dental Research 66, 1375–9.
- Haapasalo M, Udnaes T, Endal U (2003) Persistent, recurrent, and acquired infection of the root canal system posttreatment. *Endodontic Topics* **6**, 29–56.
- Hartke A, Giard JC, Laplace JM, Auffray Y (1998) Survival of Enterococcus faecalis in an oligotrophic microcosm: changes in morphology, development of general stress resistance, and analysis of protein synthesis. Applied Environmental Microbiology 64, 4238–45.
- Laplace JM, Thuault M, Hartke A, Boutibonnes P, Auffray Y (1997) Sodium hypochlorite stress in *Enterococcus faecalis*: influence of antecedent growth conditions and induced proteins. *Current Microbiology* **34**, 284–9.
- Lleò MM, Bonato B, Tafi MC, Signoretto C, Boaretti M, Canepari P (2001) Resuscitation rate in different enterococcal species in the viable but non-culturable state. *Applied Microbiology* **91**, 1095–102.

- Nair PNR, Sjögren U, Krey G, Kahnberg KE, Sundqvist G (1990) Intraradicular bacteria and fungi in root-filled, asymptomatic human teeth with therapy-resistant periapical lesions: a long-term light and electron microscopic follow-up study. *Journal of Endodontics* 16, 580–8.
- Nair PN, Henry S, Cano V, Vera J (2005) Microbial status of apical root canal system of human mandibular first molars with primary apical periodontitis after "one-visit" endodontic treatment. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics **99**, 231–52.
- Neely AN, Maley MP (2000) Survival of enterococci and staphylococci on hospital fabrics and plastic. *Journal of Clinical Microbiology* **38**, 724–6.
- Oguntebi BR (1994) Dentine tubule infection and endodontic therapy implications. *International Endodontic Journal* **27**, 218–22.
- Ørstavik D, Haapasalo M (1990) Disinfection by endodontic irrigants and dressings of experimentally infected dentinal tubules. *Endodontics and Dental Traumatology* **6**, 142–9.
- Peciuliene V, Reynaud AH, Balciuniene I, Haapasalo M (2001) Isolation of yeasts and enteric bacteria in root-filled teeth with chronic apical periodontitis. *International Endodontic Journal* **34**, 429–34.
- Peters LB, Wesselink PR, Moorer WR (1995) The fate and the role of bacteria left in root dentinal tubules. *International Endodontic Journal* **28**, 95–9.
- Peters LB, Wesselink PR, Buijs JF, van Winkelhoff AJ (2001) Viable bacteria in root dentinal tubules of teeth with apical periodontitis. *Journal of Endodontics* **27**, 76–81.
- Portenier I, Haapasalo H, Ørstavik D, Yamauchi M, Haapasalo M (2002) Inactivation of the antibacterial activity of iodine potassium iodide and chlorhexidine digluconate against *Enterococcus faecalis* by dentin, dentin matrix, type-I collagen, and heat-killed microbial whole cells. *Journal of Endodontics* 28, 634–7.
- Pruzzo C, Tarsi R, Lleo MM et al. (2002) In vitro adhesion to human cells by viable but nonculturable Enterococcus faecalis. Current Microbiology 45, 105–10.
- Rams TE, Feik D, Young V, Hammond BF, Slots J (1992) Enterococci in human periodontitis. *Oral Microbiology and Immunology* **7**, 249–52.
- Saleh IM, Ruyter IE, Haapasalo M, Ørstavik D (2004) Survival of *Enterococcus faecalis* in infected dentinal tubules after root canal filling with different root canal sealers *in vitro*. *International Endodontic Journal* 37, 193–8.

- Sedgley C, Applegate B, Nagel A, Hall D (2004a) Real-time imaging and quantification of bioluminescent bacteria in root canals *in vitro*. *Journal of Endodontics* **30**, 893–8.
- Sedgley CM, Lennan SL, Clewell DB (2004b) Prevalence, phenotype and genotype of oral enterococci. Oral Microbiology and Immunology 19, 95–101.
- Sedgley CM, Nagel A, Shelburne C, Clewell DB, Appelbe OK, Molander A (2005a) Quantitative real-time PCR detection of oral Enterococcus faecalis. Archives of Oral Biology 50, 575–83.
- Sedgley CM, Molander A, Flannagan SE et al. (2005b) Virulence, phenotype and genotype characteristics of endodontic Enterococcus spp. Oral Microbiology and Immunology 20, 10–9.
- Sedgley CM, Buck G, Appelbe OK. Detection of *Enterococcus faecalis* at multiple oral sites in endodontic patients using culture and PCR. *Journal of Endodontics* (in press).
- Signoretto C, del Mar Lleo M, Tafi MC, Canepari P (2000) Cell wall chemical composition of *Enterococcus faecalis* in the viable but nonculturable state. *Applied Environmental Microbiology* **66**, 1953–9.
- Siqueira JF Jr, Rôças IN (2004) Polymerase chain reactionbased analysis of microorganisms associated with failed endodontic treatment. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics 97, 85–94.
- Siqueira JF Jr (2001) Aetiology of root canal treatment failure: why well-treated teeth can fail. *International Endodontic Journal* **34**, 1–10.
- Sjögren U, Figdor D, Persson S, Sundqvist G (1997) Influence of infection at the time of root filling on the outcome of endodontic treatment of teeth with apical periodontitis. *International Endodontic Journal* **30**, 297–306.
- Sundqvist G, Figdor D, Persson S, Sjögren U (1998) Microbiologic analysis of teeth with failed endodontic treatment and the outcome of conservative re-treatment. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics 85, 86–93.
- Takemura N, Noiri Y, Ehara A, Kawahara T, Noguchi N, Ebisu S (2004) Single species biofilm-forming ability of root canal isolates on gutta-percha points. *European Journal of Oral Sciences* **112**, 523–9.
- Waters CM, Antiporta MH, Murray BE, Dunny GM (2003) Role of the *Enterococcus faecalis* GelE protease in determination of cellular chain length, supernatant pheromone levels, and degradation of fibrin and misfolded surface proteins. *Journal of Bacteriology* 185, 3613–23.

742

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.