

Keyword Index

- 3d reconstruction, Garrè's osteomyelitis managed by root canal treatment of a mandibular second molar: incorporation of computed tomography with 3D reconstruction in the diagnosis and monitoring of the disease, 255
- adhesion, Growth at high pH increases *Enterococcus faecalis* adhesion to collagen, 389
- adhesion, *Ex vivo* study of the adhesion of an epoxy-based sealer to human dentine submitted to irradiation with Er : YAG and Nd : YAG lasers, 866
- adhesive bonding, Cytotoxicity of substances leached or dissolved from pulp capping materials, 505
- adoption of technology, The adoption of new endodontic technology amongst Danish general dental practitioners, 52
- aging, Reduction of osteocalcin expression in aged human dental pulp, 817
- albumin, Growth at high pH increases *Enterococcus faecalis* adhesion to collagen, 389
- ameloblastoma, Unicystic ameloblastoma: a possible pitfall in periapical diagnosis, 334
- anatomic endodontic technology, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- anatomy, The isthmuses of the mesial root of mandibular molars: a micro-computed tomographic study, 558
- antibacterial, An *in vitro* evaluation of the ability of ozone to kill a strain of *Enterococcus faecalis*, 22
- antibiotic prophylaxis, The role of preoperative prophylactic antibiotic administration in periapical endodontic surgery: a randomized, prospective double-blind placebo-controlled study, 877
- antimicrobial activity, Short-term antibacterial activity of root canal sealers towards *Enterococcus faecalis*, 483
- antimicrobial activity, *In vitro* antimicrobial activity of Fill Canal, Sealapex, Mineral Trioxide Aggregate, Portland cement and EndoRez, 539
- apex locator, *Ex vivo* comparison of two electronic apex locators with different scales and frequencies, 855
- apexification, The effect of the renewal of calcium hydroxide paste on the apexification and periapical healing of teeth with incomplete root formation, 436
- apical extrusion, Apical extrusion of intracanal bacteria following use of two engine-driven instrumentation techniques, 871
- apical file size determination, Influence of cervical preflaring on apical file size determination, 430
- apical leakage, The sealing ability of an epoxy resin root canal sealer after Nd:YAG laser irradiation of the root canal, 302
- apical periodontitis, Frequency and distribution of root filled teeth and apical periodontitis in a Greek population, 105
- apical periodontitis, Prevalence of apical periodontitis and the quality of endodontic treatment in an adult Belarusian population, 238
- apical periodontitis, Calculus-like deposit on the apical external root surface of teeth with post-treatment apical periodontitis: report of two cases, 262
- apical periodontitis, A comparative analysis of periapical health based on historic and current data, 277
- apical periodontitis, High prevalence of apical periodontitis amongst type 2 diabetic patients, 564
- apical surgery, SEM analysis of the integrity of resected root apices of cadaver and extracted teeth after ultrasonic root-end preparation at different intensities, 310
- apicectomy, Periapical resurgery versus periapical surgery: a 5-year longitudinal comparison, 320
- apicectomy, Determining the apical terminus of root-end resected teeth using three modern apex locators: a comparative *ex vivo* study, 827
- apoptosis, Apoptosis in epithelial cells of apical radicular cysts, 465
- arachidonic acid, Heat shock induces the synthesis of the inflammatory mediator leukotriene B₄ in human pulp cells, 882
- asians, Difference in root canal length between Asians and Caucasians, 149
- atomic force microscopy, Effects of 2% chlorhexidine and 5.25% sodium hypochlorite on gutta-percha cones studied by atomic force microscopy, 425
- automated root canal preparation, Comparative study of six rotary nickel–titanium systems and hand instrumentation for root canal preparation, 743
- automated root canal preparation, Comparison of root canal preparation using RaCe and ProTaper rotary Ni-Ti instruments, 8
- automatic circuit breaker, Reduction of excessive electrical stimulus during electric pulp testing, 544
- bacteria, Growth at high pH increases *Enterococcus faecalis* adhesion to collagen, 389
- bacteria, Apical extrusion of intracanal bacteria following use of two engine-driven instrumentation techniques, 871
- bacteria, Influence of irrigant needle depth in removing bioluminescent bacteria inoculated into

- instrumented root canals using real-time imaging *in vitro*, 97
- bacterial morphology, Microscopic observation of bacteria: review highlighting the use of environmental SEM, 775
- biocompatibility, Effect of ProRoot® MTA mixed with chlorhexidine on apoptosis and cell cycle of fibroblasts and macrophages *in vitro*, 137
- biocompatibility, The chemical constitution and biocompatibility of accelerated Portland cement for endodontic use, 834
- biocompatibility, Effect of mineral trioxide aggregate on cytokine production by peritoneal macrophages, 896
- bioluminescence, Influence of irrigant needle depth in removing bioluminescent bacteria inoculated into instrumented root canals using real-time imaging *in vitro*, 97
- bleaching agents, Effect of bleaching agents on bonding to pulp chamber dentine, 211
- bleaching gel, *In vitro* peroxide penetration into the pulp chamber from newer bleaching products, 516
- bonding resins, Penetration of bonding resins into fibre-reinforced composite posts: a confocal microscopic study, 46
- bonding, Effect of bleaching agents on bonding to pulp chamber dentine, 211
- bridges, Fate of vital pulps beneath a metal-ceramic crown or a bridge retainer, 521
- cadavers, SEM analysis of the integrity of resected root apices of cadaver and extracted teeth after ultrasonic root-end preparation at different intensities, 310
- calcitonin gene-related peptide receptor, Calcitonin gene-related peptide receptor expression in healthy and inflamed human pulp tissue, 712
- calcium hydroxide plus points, Calcium release and pH-characteristics of calcium hydroxide plus points, 683
- calcium hydroxide points, Calcium release and pH-characteristics of calcium hydroxide plus points, 683
- calcium hydroxide, Dental pulp capping: effect of Emdogain Gel on experimentally exposed human pulps, 186
- calcium hydroxide, Growth at high pH increases *Enterococcus faecalis* adhesion to collagen, 389
- calcium hydroxide, The effect of the renewal of calcium hydroxide paste on the apexification and periapical healing of teeth with incomplete root formation, 436
- calcium hydroxide, Cytotoxicity of substances leached or dissolved from pulp capping materials, 505
- calcium hydroxide, Recovery of *Enterococcus faecalis* after single- or multiple-visit root canal treatments carried out in infected teeth *ex vivo*, 697
- calcium hydroxide, Calcium hydroxide dressings using different preparation and application modes: density and dissolution by simulated tissue pressure, 889
- calcium release, Calcium release and pH-characteristics of calcium hydroxide plus points, 683
- calculus, Calculus-like deposit on the apical external root surface of teeth with post-treatment apical periodontitis: report of two cases, 262
- canal aberration, Comparative study of six rotary nickel-titanium systems and hand instrumentation for root canal preparation, 743
- canal anatomy, Difference in root canal length between Asians and Caucasians, 149
- canal cleaning, The effect of high-frequency electrical pulses on organic tissue in root canals, 531
- canal curvature, Influence of different types of automated devices on the shaping ability of rotary nickel-titanium FlexMaster instruments, 627
- canal detection, Detection of the second mesiobuccal canal in mesiobuccal roots of maxillary molar teeth *ex vivo*, 124
- canal fillings, Intracanal temperature rise evaluation during the usage of the System B: replication of intracanal anatomy, 218
- canal isthmus, Frequency and type of canal isthmuses in first molars detected by endoscopic inspection during periradicular surgery, 160
- canal length, Difference in root canal length between Asians and Caucasians, 149
- canal shape, Root canal preparation with Endo-Eze AET: changes in root canal shape assessed by micro-computed tomography, 456
- canal shape, Calcium hydroxide dressings using different preparation and application modes: density and dissolution by simulated tissue pressure, 889
- canal straightening, Influence of different types of automated devices on the shaping ability of rotary nickel-titanium FlexMaster instruments, 627
- canal transportation, Canal preparation with Hero 642 rotary Ni-Ti instruments compared with stainless steel hand K-file assessed using computed tomography, 402
- canal transportation, Influence of different types of automated devices on the shaping ability of rotary nickel-titanium FlexMaster instruments, 627
- capillary flow porometry, Longitudinal study on microleakage of three root-end filling materials by the fluid transport method and by capillary flow porometry, 129
- capillary rheometer, A study of the rheological properties of endodontic sealers, 499
- capsaicin, The effect of capsaicin on substance P expression in pulp tissue inflammation, 30
- carbamide peroxide, *In vitro* peroxide penetration into the pulp chamber from newer bleaching products, 516
- Carisolv™, Carisolv™: an alternative to NaOCl in immature root canals?, 448

- casein zymography, Examination of the signal transduction pathways leading to upregulation of tissue type plasminogen activator by *Porphyromonas endodontalis* in human pulp cells, 860
- Caucasians, Difference in root canal length between Asians and Caucasians, 149
- cell culture, Cytotoxicity of substances leached or dissolved from pulp capping materials, 505
- cell death, Effect of ProRoot® MTA mixed with chlorhexidine on apoptosis and cell cycle of fibroblasts and macrophages *in vitro*, 137
- chelating agent, The effect of application time of EDTA and NaOCl on intracanal smear layer removal: an SEM analysis, 285
- chemical constitution, The chemical constitution and biocompatibility of accelerated Portland cement for endodontic use, 834
- chlorhexidine gel, Recovery of *Enterococcus faecalis* after single- or multiple-visit root canal treatments carried out in infected teeth *ex vivo*, 697
- chlorhexidine, Effects of 2% chlorhexidine and 5.25% sodium hypochlorite on gutta-percha cones studied by atomic force microscopy, 425
- chloroform, Effect of customization of master gutta-percha cone on apical control of root filling using different techniques: an *ex vivo* study, 658
- cleaning efficacy, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- collagen, Growth at high pH increases *Enterococcus faecalis* adhesion to collagen, 389
- composite crown, *Ex vivo* fracture resistance of direct resin composite complete crowns with and without posts on maxillary premolars, 230
- computed tomography, Garre's osteomyelitis managed by root canal treatment of a mandibular second molar: incorporation of computed tomography with 3D reconstruction in the diagnosis and monitoring of the disease, 255
- computed tomography, Canal preparation with Hero 642 rotary Ni-Ti instruments compared with stainless steel hand K-file assessed using computed tomography, 402
- condensing osteitis, A comparative analysis of periapical health based on historic and current data, 277
- conditioning dentine, Endodontic complications after plastic restorations in general practice, 409
- configuration of root canal systems, Detection of the second mesiobuccal canal in mesiobuccal roots of maxillary molar teeth *ex vivo*, 124
- confocal microscopy, Penetration of bonding resins into fibre-reinforced composite posts: a confocal microscopic study, 46
- controlled torque, Shaping ability of two rotary instruments in simulated canals: stainless steel ENDOflash and nickel-titanium HERO Shaper, 637
- coronal flaring, Influence of cervical preflaring on apical file size determination, 430
- coronal leakage, The sealing ability of an epoxy resin root canal sealer after Nd:YAG laser irradiation of the root canal, 302
- cross-sectional, A 24-year follow-up of root filled teeth and periapical health amongst middle aged and elderly women in Göteborg, Sweden, 246
- crown-down, Anatomic Endodontic Technology (AET) – a crown-down root canal preparation technique: basic concepts, operative procedure and instruments, 575
- crowns, Fate of vital pulps beneath a metal-ceramic crown or a bridge retainer, 521
- cryogenic, Effect of cryogenic treatment on nickel-titanium endodontic instruments, 364
- curved canals, A comparative study of Endoflare–Hero Shaper and Mtwo NiTi instruments in the preparation of curved root canals, 610
- curved root canals, Influence of different types of automated devices on the shaping ability of rotary nickel-titanium FlexMaster instruments, 627
- customization, Effect of customization of master gutta-percha cone on apical control of root filling using different techniques: an *ex vivo* study, 658
- cutting efficiency, Effect of cryogenic treatment on nickel-titanium endodontic instruments, 364
- cyclic loading, Physical and mechanical characterization and the influence of cyclic loading on the behaviour of nickel-titanium wires employed in the manufacture of rotary endodontic instruments, 795
- cytokines, Effect of mineral trioxide aggregate on cytokine production by peritoneal macrophages, 896
- cytotoxicity, Effect of ProRoot® MTA mixed with chlorhexidine on apoptosis and cell cycle of fibroblasts and macrophages *in vitro*, 137
- cytotoxicity, Cytotoxicity of MTA and Portland cement on human ECV 304 endothelial cells, 604
- decision-making, An assessment of endodontic re-treatment decision-making in an educational setting, 470
- dental anthropology, A comparative analysis of periapical health based on historic and current data, 277
- dental pulp capping, Cytotoxicity of substances leached or dissolved from pulp capping materials, 505
- dental pulp exposure, Dental pulp capping: effect of Emdogain Gel on experimentally exposed human pulps, 186
- dental pulp, Copper–zinc superoxide dismutase activity in healthy and inflamed human dental pulp, 195
- dental pulp, Cytotoxicity of substances leached or dissolved from pulp capping materials, 505
- dental pulp, Reduction of osteocalcin expression in aged human dental pulp, 817

- dental radiograph, Methods of experimental induction of periapical inflammation. Microbiological and radiographic evaluation, 477
- dental radiography, Periapical status and quality of endodontic treatment in an adult Irish population, 81
- dental, A laboratory comparison of three imaging systems for image quality and radiation exposure characteristics, 645
- dentin, Microhardness changes in dentine after neonatal capsaicin application, 570
- dentine debris, The efficacy of ultrasonic irrigation to remove artificially placed dentine debris from human root canals prepared using instruments of varying taper, 764
- dentine permeability, Influence of the pulpal components on human dentine permeability *in vitro*, 152
- dentine, Effect of bleaching agents on bonding to pulp chamber dentine, 211
- dentinogenesis, Dental pulp capping: effect of Emdogain Gel on experimentally exposed human pulps, 186
- dentistry – treatment, The use of an intra-oral injection of ketorolac in the treatment of irreversible pulpitis, 789
- diabetes mellitus, High prevalence of apical periodontitis amongst type 2 diabetic patients, 564
- diagnosis, Unicystic ameloblastoma: a possible pitfall in periapical diagnosis, 334
- diagnosis, Three-dimensional, non-destructive visualization of vertical root fractures using flat panel volume detector computer tomography: an *ex vivo* *in vitro* case report, 904
- digital radiography, An *ex vivo* comparison of conventional and digital radiography for perceived image quality of root fillings, 397
- digital, A laboratory comparison of three imaging systems for image quality and radiation exposure characteristics, 645
- dissolution, Calcium hydroxide dressings using different preparation and application modes: density and dissolution by simulated tissue pressure, 889
- DNA replication, Effect of ProRoot® MTA mixed with chlorhexidine on apoptosis and cell cycle of fibroblasts and macrophages *in vitro*, 137
- dye leakage, The sealing ability of an epoxy resin root canal sealer after Nd:YAG laser irradiation of the root canal, 302
- EDTA, Effect of liquid and paste-type lubricants on torque values during simulated rotary root canal instrumentation, 223
- EDTA, The effect of application time of EDTA and NaOCl on intracanal smear layer removal: an SEM analysis, 285
- EDTA, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- EDTA, Assessment of a gel-type chelating preparation containing 1-hydroxyethylidene-1, 1-bisphosphonate, 810
- electric pulp test, Reduction of excessive electrical stimulus during electric pulp testing, 544
- electrical conductivity, Physico-chemical properties of MTA and a novel experimental cement, 443
- electromyography, Reduction of excessive electrical stimulus during electric pulp testing, 544
- electronic apex locator, *Ex vivo* evaluation of the capacity of the Tri Auto ZX to locate the apical foramen during root canal retreatment, 718
- electronic apex locators, A comparison between two electronic apex locators: an *in vivo* investigation, 36
- electronic apex locators, The adoption of new endodontic technology amongst Danish general dental practitioners, 52
- electronic apex locators, Determining the apical terminus of root-end resected teeth using three modern apex locators: a comparative *ex vivo* study, 827
- electronic odontology, *Ex vivo* evaluation of the capacity of the Tri Auto ZX to locate the apical foramen during root canal retreatment, 718
- endodontic epidemiology, Frequency and distribution of root filled teeth and apical periodontitis in a Greek population, 105
- endodontic failure, Calculus-like deposit on the apical external root surface of teeth with post-treatment apical periodontitis: report of two cases, 262
- endodontic instruments, Anatomic Endodontic Technology (AET) – a crown-down root canal preparation technique: basic concepts, operative procedure and instruments, 575
- endodontic instruments, Physical and mechanical characterization and the influence of cyclic loading on the behaviour of nickel-titanium wires employed in the manufacture of rotary endodontic instruments, 795
- endodontic obturation, Presence of pores and vacuoles in set endodontic sealers, 690
- endodontic re-treatment, An assessment of endodontic re-treatment decision-making in an educational setting, 470
- endodontic retreatment, *Ex vivo* evaluation of the capacity of the Tri Auto ZX to locate the apical foramen during root canal retreatment, 718
- endodontic sealers, A study of the rheological properties of endodontic sealers, 499
- endodontic sealers, *In vitro* antimicrobial activity of Fill Canal, Sealapex, Mineral Trioxide Aggregate, Portland cement and EndoRez, 539
- endodontic sealers, Presence of pores and vacuoles in set endodontic sealers, 690
- endodontic spreader loads, A comparison of spreader penetration depth and load required during lateral condensation in teeth prepared using various root canal preparation techniques, 510

- endodontic therapy, Effect of endodontic procedures on enterococci, enteric bacteria and yeasts in primary endodontic infections, 372
- endodontic treatment, Prevalence of persistent pain after endodontic treatment and factors affecting its occurrence in cases with complete radiographic healing, 169
- endodontic treatment, An *ex vivo* comparison of conventional and digital radiography for perceived image quality of root fillings, 397
- endodontic, A 24-year follow-up of root filled teeth and periapical health amongst middle aged and elderly women in Göteborg, Sweden, 246
- endodontically treated teeth, Frequency and distribution of root filled teeth and apical periodontitis in a Greek population, 105
- endodontically treated, *Ex vivo* fracture resistance of direct resin composite complete crowns with and without posts on maxillary premolars, 230
- endodontics, Effect of ProRoot® MTA mixed with chlorhexidine on apoptosis and cell cycle of fibroblasts and macrophages *in vitro*, 137
- endodontics, Difference in root canal length between Asians and Caucasians, 149
- endodontics, Outcome of root canal treatment using Thermafil and cold lateral condensation filling techniques, 179
- endodontics, Periapical resurgery versus periapical surgery: a 5-year longitudinal comparison, 320
- endodontics, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- endodontics, Methods of experimental induction of periapical inflammation. Microbiological and radiographic evaluation, 477
- endodontics, The isthmuses of the mesial root of mandibular molars: a micro-computed tomographic study, 558
- endodontics, High prevalence of apical periodontitis amongst type 2 diabetic patients, 564
- endodontics, A laboratory comparison of three imaging systems for image quality and radiation exposure characteristics, 645
- endodontics, Effect of *E. faecalis* on the release of serine proteases elastase and cathepsin G, and collagenase-2 (MMP-8) by human polymorphonuclear leukocytes (PMNs), 667
- endodontics, Neurological complications following extrusion of sodium hypochlorite solution during root canal treatment, 843
- endodontics, *Ex vivo* comparison of two electronic apex locators with different scales and frequencies, 855
- endodontology, Periapical status and quality of endodontic treatment in an adult Irish population, 81
- Endo-Eze aet, Root canal preparation with Endo-Eze AET: changes in root canal shape assessed by micro-computed tomography, 456
- Endorez, *In vitro* antimicrobial activity of Fill Canal, Sealapex, Mineral Trioxide Aggregate, Portland cement and EndoRez, 539
- endoscopy, Frequency and type of canal isthmuses in first molars detected by endoscopic inspection during periradicular surgery, 160
- energy-dispersive x-ray spectroscopy, Effect of cryogenic treatment on nickel-titanium endodontic instruments, 364
- engine-driven techniques, Apical extrusion of intracanal bacteria following use of two engine-driven instrumentation techniques, 871
- Enterococcus faecalis*, An *in vitro* evaluation of the ability of ozone to kill a strain of *Enterococcus faecalis*, 22
- enteric bacteria, Effect of endodontic procedures on enterococci, enteric bacteria and yeasts in primary endodontic infections, 372
- enterococci, Effect of endodontic procedures on enterococci, enteric bacteria and yeasts in primary endodontic infections, 372
- enterococci, Effect of *E. faecalis* on the release of serine proteases elastase and cathepsin G, and collagenase-2 (MMP-8) by human polymorphonuclear leukocytes (PMNs), 667
- Enterococcus faecalis*, Recovery of *Enterococcus faecalis* after single- or multiple-visit root canal treatments carried out in infected teeth *ex vivo*, 697
- Enterococcus faecalis*, Survival of *Enterococcus faecalis* in root canals *ex vivo*, 735
- Enterococcus faecalis*, Short-term antibacterial activity of root canal sealers towards *Enterococcus faecalis*, 483
- epidemiology, A 24-year follow-up of root filled teeth and periapical health amongst middle aged and elderly women in Göteborg, Sweden, 246
- epidemiology, Periapical status and quality of endodontic treatment in an adult Irish population, 81
- evidence-based dentistry, Effectiveness of single- versus multiple-visit endodontic treatment of teeth with apical periodontitis: a systematic review and meta-analysis, 347
- excessive stimulus time, Reduction of excessive electrical stimulus during electric pulp testing, 544
- expansion, A temporary filling material may cause cuspal deflection, infractions and fractures in endodontically treated teeth, 653
- extraradicular infection, Calculus-like deposit on the apical external root surface of teeth with post-treatment apical periodontitis: report of two cases, 262
- failure mode, *Ex vivo* fracture resistance of direct resin composite complete crowns with and without posts on maxillary premolars, 230
- fatigue, Physical and mechanical characterization and the influence of cyclic loading on the behaviour of nickel-titanium wires employed in the manufacture of rotary endodontic instruments, 795

- fatigue, Defects in ProTaper S1 instruments after clinical use: fractographic examination, 802
- fibre posts, Penetration of bonding resins into fibre-reinforced composite posts: a confocal microscopic study, 46
- fibre reinforced, *Ex vivo* fracture resistance of direct resin composite complete crowns with and without posts on maxillary premolars, 230
- field emission-scanning electron microscope, Shear bond strength of Resilon to a methacrylate-based root canal sealer, 753
- fill, Calcium hydroxide dressings using different preparation and application modes: density and dissolution by simulated tissue pressure, 889
- filter membrane, Short-term antibacterial activity of root canal sealers towards *Enterococcus faecalis*, 483
- filter paper, Short-term antibacterial activity of root canal sealers towards *Enterococcus faecalis*, 483
- fixed prosthodontics, Fate of vital pulps beneath a metal-ceramic crown or a bridge retainer, 521
- flat panel volume detector computer tomography, Three-dimensional, non-destructive visualization of vertical root fractures using flat panel volume detector computer tomography: an *ex vivo in vitro* case report, 904
- flexural fatigue, Defects in ProTaper S1 instruments after clinical use: longitudinal examination, 550
- flowable composite resin, The effect of fibre insertion on fracture resistance of root filled molar teeth with MOD preparations restored with composite, 73
- fluid movement, Periapical biomechanics and the role of cyclic biting force in apical retrograde fluid movement, 597
- fluid transport method, Longitudinal study on microleakage of three root-end filling materials by the fluid transport method and by capillary flow porometry, 129
- fluid transport model, The sealing ability of an epoxy resin root canal sealer after Nd:YAG laser irradiation of the root canal, 302
- fluorescent probes, Penetration of bonding resins into fibre-reinforced composite posts: a confocal microscopic study, 46
- fracture resistance, *Ex vivo* fracture resistance of direct resin composite complete crowns with and without posts on maxillary premolars, 230
- fracture resistance, The effect of fibre insertion on fracture resistance of root filled molar teeth with MOD preparations restored with composite, 73
- fracture, Defects in ProTaper S1 instruments after clinical use: longitudinal examination, 550
- fracture, Defects in ProTaper S1 instruments after clinical use: fractographic examination, 802
- fractured instrument, Probability of removing fractured instruments from root canals, 112
- Fuji IX, Longitudinal study on microleakage of three root-end filling materials by the fluid transport method and by capillary flow porometry, 129
- Garré's osteomyelitis, Garré's osteomyelitis managed by root canal treatment of a mandibular second molar: incorporation of computed tomography with 3D reconstruction in the diagnosis and monitoring of the disease, 255
- general practice, Endodontic complications after plastic restorations in general practice, 409
- Glyde, Assessment of a gel-type chelating preparation containing 1-hydroxyethylidene-1, 1-bisphosphonate, 810
- granulation tissue, Collagenase-3 (MMP-13) is expressed in periapical lesions: an immunohistochemical study, 297
- gutta-percha cone, Effects of 2% chlorhexidine and 5.25% sodium hypochlorite on gutta-percha cones studied by atomic force microscopy, 425
- gutta-percha, Effectiveness of different techniques for removing gutta-percha during retreatment, 2
- gutta-percha, Intracanal temperature rise evaluation during the usage of the System B: replication of intracanal anatomy, 218
- gutta-percha, Calcium release and pH-characteristics of calcium hydroxide plus points, 683
- gutta-percha, An evaluation of .06 tapered gutta-percha cones for filling of .06 taper prepared curved root canals, 87
- hardness, Microhardness changes in dentine after neonatal capsaicin application, 570
- heat shock, Heat shock induces the synthesis of the inflammatory mediator leukotriene B₄ in human pulp cells, 882
- HEBP, Assessment of a gel-type chelating preparation containing 1-hydroxyethylidene-1, 1-bisphosphonate, 810
- high-frequency electrical pulses, The effect of high-frequency electrical pulses on organic tissue in root canals, 531
- histology, Dental pulp capping: effect of Emdogain Gel on experimentally exposed human pulps, 186
- human polymorphonuclear leukocyte, Effect of *E. faecalis* on the release of serine proteases elastase and cathepsin G, and collagenase-2 (MMP-8) by human polymorphonuclear leukocytes (PMNs), 667
- human pulp, Calcitonin gene-related peptide receptor expression in healthy and inflamed human pulp tissue, 712
- human, Reduction of osteocalcin expression in aged human dental pulp, 817
- hydrogen peroxide peroxide penetration, *In vitro* peroxide penetration into the pulp chamber from newer bleaching products, 516
- hypochlorite, Neurological complications following extrusion of sodium hypochlorite solution during root canal treatment, 843

- iatrogenic errors, Technical quality of root canal treatment and detection of iatrogenic errors in an undergraduate dental clinic, 725
- in vitro*, Survival of *Enterococcus faecalis* in root canals *ex vivo*, 735
- image enhancement, An *ex vivo* comparison of conventional and digital radiography for perceived image quality of root fillings, 397
- imaging techniques, The imaging of root canal obturation using micro-CT, 617
- immunohistochemistry, Dental pulp capping: effect of Emdogain Gel on experimentally exposed human pulps, 186
- immunohistochemistry, Upregulation of tissue-type plasminogen activator in inflamed human dental pulps, 328
- immunohistochemistry, Apoptosis in epithelial cells of apical radicular cysts, 465
- impedance, A comparison between two electronic apex locators: an *in vivo* investigation, 36
- inflammation, Copper-zinc superoxide dismutase activity in healthy and inflamed human dental pulp, 195
- inflammation, Upregulation of tissue-type plasminogen activator in inflamed human dental pulps, 328
- innate immunity, Effect of mineral trioxide aggregate on cytokine production by peritoneal macrophages, 896
- instrument fracture, Effect of liquid and paste-type lubricants on torque values during simulated rotary root canal instrumentation, 223
- instrument removal, Probability of removing fractured instruments from root canals, 112
- instrument type, Influence of cervical preflaring on apical file size determination, 430
- intermediate restorative material, Behaviour of bone marrow osteoblast-like cells on mineral trioxide aggregate: morphology and expression of type I collagen and bone-related protein mRNAs, 203
- inter-observer variation, An assessment of endodontic re-treatment decision-making in an educational setting, 470
- intracanal medication, Calcium release and pH-characteristics of calcium hydroxide plus points, 683
- intraosseous, Pressures generated *in vitro* during Stabident intraosseous injections, 291
- IRM, Longitudinal study on microleakage of three root-end filling materials by the fluid transport method and by capillary flow porometry, 129
- irrigant needle depth, Influence of irrigant needle depth in removing bioluminescent bacteria inoculated into instrumented root canals using real-time imaging *in vitro*, 97
- irrigant, CarisolTM: an alternative to NaOCl in immature root canals?, 448
- irrigants, The effect of high-frequency electrical pulses on organic tissue in root canals, 531
- irrigating solutions, The effect of application time of EDTA and NaOCl on intracanal smear layer removal: an SEM analysis, 285
- irrigation, The efficacy of ultrasonic irrigation to remove artificially placed dentine debris from human root canals prepared using instruments of varying taper, 764
- irrigation, Influence of irrigant needle depth in removing bioluminescent bacteria inoculated into instrumented root canals using real-time imaging *in vitro*, 97
- isthmus, The isthmuses of the mesial root of mandibular molars: a micro-computed tomographic study, 558
- ketorolac, The use of an intra-oral injection of ketorolac in the treatment of irreversible pulpitis, 789
- K-Flexofiles, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- laser and irradiation, *Ex vivo* study of the adhesion of an epoxy-based sealer to human dentine submitted to irradiation with Er : YAG and Nd : YAG lasers, 866
- lateral condensation, Outcome of root canal treatment using Therafil and cold lateral condensation filling techniques, 179
- lateral condensation, A comparison of spreader penetration depth and load required during lateral condensation in teeth prepared using various root canal preparation techniques, 510
- leukotriene, Heat shock induces the synthesis of the inflammatory mediator leukotriene B₄ in human pulp cells, 882
- local anaesthetic, Pressures generated *in vitro* during Stabident intraosseous injections, 291
- logistic regression, A comparative analysis of periapical health based on historic and current data, 277
- longitudinal, A 24-year follow-up of root filled teeth and periapical health amongst middle aged and elderly women in Göteborg, Sweden, 246
- long-time follow-up, Garré's osteomyelitis managed by root canal treatment of a mandibular second molar: incorporation of computed tomography with 3D reconstruction in the diagnosis and monitoring of the disease, 255
- lubricant, Effect of liquid and paste-type lubricants on torque values during simulated rotary root canal instrumentation, 223
- macrophages, Effect of mineral trioxide aggregate on cytokine production by peritoneal macrophages, 896
- matrix concept, Perforation repair with mineral trioxide aggregate: a modified matrix concept, 59

- maxillary molars, Detection of the second mesiobuccal canal in mesiobuccal roots of maxillary molar teeth *ex vivo*, 124
- mechanical properties, Physical and mechanical characterization and the influence of cyclic loading on the behaviour of nickel-titanium wires employed in the manufacture of rotary endodontic instruments, 795
- mesio-lingual canal, Detection of the second mesiobuccal canal in mesiobuccal roots of maxillary molar teeth *ex vivo*, 124
- methacrylate sealer, Shear bond strength of Resilon to a methacrylate-based root canal sealer, 753
- micro-computed tomography, Root canal preparation with Endo-Eze AET: changes in root canal shape assessed by micro-computed tomography, 456
- micro-computed tomography, The isthmuses of the mesial root of mandibular molars: a micro-computed tomographic study, 558
- micro-computed tomography, The imaging of root canal obturation using micro-CT, 617
- microhardness, Effect of cryogenic treatment on nickel-titanium endodontic instruments, 364
- microorganism, Methods of experimental induction of periapical inflammation. Microbiological and radiographic evaluation, 477
- microscopic techniques, Microscopic observation of bacteria: review highlighting the use of environmental SEM, 775
- mineral trioxide aggregate, Behaviour of bone marrow osteoblast-like cells on mineral trioxide aggregate: morphology and expression of type I collagen and bone-related protein mRNAs, 203
- mineral trioxide aggregate, Physico-chemical properties of MTA and a novel experimental cement, 443
- mineral trioxide aggregate, *In vitro* antimicrobial activity of Fill Canal, Sealapex, Mineral Trioxide Aggregate, Portland cement and EndoRez, 539
- mineral trioxide aggregate, Perforation repair with mineral trioxide aggregate: a modified matrix concept, 59
- mineral trioxide aggregate, Cytotoxicity of MTA and Portland cement on human ECV 304 endothelial cells, 604
- mineral trioxide aggregate, The chemical constitution and biocompatibility of accelerated Portland cement for endodontic use, 834
- mineral trioxide aggregate, Effect of mineral trioxide aggregate on cytokine production by peritoneal macrophages, 896
- MMP-13, Collagenase-3 (MMP-13) is expressed in periapical lesions: an immunohistochemical study, 297
- MMP-8, Effect of *E. faecalis* on the release of serine proteases elastase and cathepsin G, and collagenase-2 (MMP-8) by human polymorphonuclear leukocytes (PMNs), 667
- molar, Frequency and type of canal isthmuses in first molars detected by endoscopic inspection during periradicular surgery, 160
- MTA, Longitudinal study on microleakage of three root-end filling materials by the fluid transport method and by capillary flow porometry, 129
- NaOCl, Assessment of a gel-type chelating preparation containing 1-hydroxyethylidene-1, 1-bisphosphonate, 810
- naproxen sodium, Influence of the pulpal components on human dentine permeability *in vitro*, 152
- native state, Microscopic observation of bacteria: review highlighting the use of environmental SEM, 775
- natural teeth, The efficacy of ultrasonic irrigation to remove artificially placed dentine debris from human root canals prepared using instruments of varying taper, 764
- Nd:YAG laser, The sealing ability of an epoxy resin root canal sealer after Nd:YAG laser irradiation of the root canal, 302
- nerve damage, Neurological complications following extrusion of sodium hypochlorite solution during root canal treatment, 843
- neurogenic inflammation, Calcitonin gene-related peptide receptor expression in healthy and inflamed human pulp tissue, 712
- neuronal, Microhardness changes in dentine after neonatal capsaicin application, 570
- Ni-Ti instruments, Comparative study of six rotary nickel-titanium systems and hand instrumentation for root canal preparation, 743
- nickel-titanium, Canal preparation with Hero 642 rotary Ni-Ti instruments compared with stainless steel hand K-file assessed using computed tomography, 402
- nickel-titanium, Defects in ProTaper S1 instruments after clinical use: longitudinal examination, 550
- nickel-titanium, Effectiveness of different techniques for removing gutta-percha during retreatment, 2
- nickel-titanium, Analysis of forces developed during mechanical preparation of extracted teeth using RaCe rotary instruments and ProFiles, 17
- nickel-titanium, Effect of cryogenic treatment on nickel-titanium endodontic instruments, 364
- nickel-titanium, Influence of different types of automated devices on the shaping ability of rotary nickel-titanium FlexMaster instruments, 627
- nickel-titanium, Physical and mechanical characterization and the influence of cyclic loading on the behaviour of nickel-titanium wires employed in the manufacture of rotary endodontic instruments, 795
- nickel-titanium, Defects in ProTaper S1 instruments after clinical use: fractographic examination, 802
- NiTi instruments, The adoption of new endodontic technology amongst Danish general dental practitioners, 52

- NiTi instruments, Comparison of root canal preparation using RaCe and ProTaper rotary Ni-Ti instruments, 8
- NiTi, A comparative study of Endoflare–Hero Shaper and Mtwo NiTi instruments in the preparation of curved root canals, 610
- nonsteroidal anti-inflammatory drugs, The use of an intra-oral injection of ketorolac in the treatment of irreversible pulpitis, 789
- obturation, Comparison of cold lateral compaction and continuous wave of obturation techniques following manual or rotary instrumentation, 381
- obturation, Effect of customization of master gutta-percha cone on apical control of root filling using different techniques: an *ex vivo* study, 658
- odontoblast, Microhardness changes in dentine after neonatal capsaicin application, 570
- odontoblastic layer, Influence of the pulpal components on human dentine permeability *in vitro*, 152
- odontogenic tumours, Unicystic ameloblastoma: a possible pitfall in periapical diagnosis, 334
- odontometry, Determining the apical terminus of root-end resected teeth using three modern apex locators: a comparative *ex vivo* study, 827
- one-step endodontics, Effectiveness of single- versus multiple-visit endodontic treatment of teeth with apical periodontitis: a systematic review and meta-analysis, 347
- open apices, The effect of the renewal of calcium hydroxide paste on the apexification and periapical healing of teeth with incomplete root formation, 436
- operating microscope, Probability of removing fractured instruments from root canals, 112
- oral health, A 24-year follow-up of root filled teeth and periapical health amongst middle aged and elderly women in Göteborg, Sweden, 246
- osteoblast-like cells, Behaviour of bone marrow osteoblast-like cells on mineral trioxide aggregate: morphology and expression of type I collagen and bone-related protein mRNAs, 203
- osteocalcin, Reduction of osteocalcin expression in aged human dental pulp, 817
- oval canals, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- oxygen scavenging, Copper–zinc superoxide dismutase activity in healthy and inflamed human dental pulp, 195
- ozone, An *in vitro* evaluation of the ability of ozone to kill a strain of *Enterococcus faecalis*, 22
- paint-on whiteners, *In vitro* peroxide penetration into the pulp chamber from newer bleaching products, 516
- partial least squares regression, A comparative analysis of periapical health based on historic and current data, 277
- pathogenesis, Apoptosis in epithelial cells of apical radicular cysts, 465
- perforation repair, Perforation repair with mineral trioxide aggregate: a modified matrix concept, 59
- periapical diseases, Unicystic ameloblastoma: a possible pitfall in periapical diagnosis, 334
- periapical healing, The effect of the renewal of calcium hydroxide paste on the apexification and periapical healing of teeth with incomplete root formation, 436
- periapical inflammation, Methods of experimental induction of periapical inflammation. Microbiological and radiographic evaluation, 477
- periapical lesions, Collagenase-3 (MMP-13) is expressed in periapical lesions: an immunohistochemical study, 297
- periapical region, Periapical biomechanics and the role of cyclic biting force in apical retrograde fluid movement, 597
- periapical status, Prevalence of apical periodontitis and the quality of endodontic treatment in an adult Belarusian population, 238
- periapical surgery, The role of preoperative prophylactic antibiotic administration in periapical endodontic surgery: a randomized, prospective double-blind placebo-controlled study, 877
- periodontal ligament, Effect of ProRoot® MTA mixed with chlorhexidine on apoptosis and cell cycle of fibroblasts and macrophages *in vitro*, 137
- periradicular surgery, Frequency and type of canal isthmuses in first molars detected by endoscopic inspection during periradicular surgery, 160
- persistent pain, Prevalence of persistent pain after endodontic treatment and factors affecting its occurrence in cases with complete radiographic healing, 169
- pH, Physico-chemical properties of MTA and a novel experimental cement, 443
- phosphonate, Assessment of a gel-type chelating preparation containing 1-hydroxyethylidene-1, 1-bisphosphonate, 810
- photostimulable storage phosphor system, An *ex vivo* comparison of conventional and digital radiography for perceived image quality of root fillings, 397
- placebo, The role of preoperative prophylactic antibiotic administration in periapical endodontic surgery: a randomized, prospective double-blind placebo-controlled study, 877
- plastic restorations, Endodontic complications after plastic restorations in general practice, 409
- polycaprolactone, Shear bond strength of Resilon to a methacrylate-based root canal sealer, 753
- polyethylene ribbon fibre, The effect of fibre insertion on fracture resistance of root filled molar teeth with MOD preparations restored with composite, 73
- polymers, Penetration of bonding resins into fibre-reinforced composite posts: a confocal microscopic study, 46

- Porphyromonas endodontalis*, Examination of the signal transduction pathways leading to upregulation of tissue type plasminogen activator by *Porphyromonas endodontalis* in human pulp cells, 860
- porosity, Presence of pores and vacuoles in set endodontic sealers, 690
- Portland cement, Physico-chemical properties of MTA and a novel experimental cement, 443
- Portland cement, *In vitro* antimicrobial activity of Fill Canal, Sealapex, Mineral Trioxide Aggregate, Portland cement and EndoRez, 539
- Portland cement, Cytotoxicity of MTA and Portland cement on human ECV 304 endothelial cells, 604
- post removal, Temperature rise of the post and on the root surface during ultrasonic post removal, 705
- post, *Ex vivo* fracture resistance of direct resin composite complete crowns with and without posts on maxillary premolars, 230
- pressure, Pressures generated *in vitro* during Stabident intraosseous injections, 291
- ProFile, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- ProFile, Comparison of cold lateral compaction and continuous wave of obturation techniques following manual or rotary instrumentation, 381
- ProTaper, Comparison of root canal preparation using RaCe and ProTaper rotary Ni-Ti instruments, 8
- prototype cements, The chemical constitution and biocompatibility of accelerated Portland cement for endodontic use, 834
- pulp breakdown, Endodontic complications after plastic restorations in general practice, 409
- pulp cells, Examination of the signal transduction pathways leading to upregulation of tissue type plasminogen activator by *Porphyromonas endodontalis* in human pulp cells, 860
- pulp cells, Heat shock induces the synthesis of the inflammatory mediator leukotriene B₄ in human pulp cells, 882
- pulp chamber, Effect of bleaching agents on bonding to pulp chamber dentine, 211
- pulp, Upregulation of tissue-type plasminogen activator in inflamed human dental pulps, 328
- pulp, Microhardness changes in dentine after neonatal capsaicin application, 570
- pulpal components, Influence of the pulpal components on human dentine permeability *in vitro*, 152
- pulpal necrosis, Fate of vital pulps beneath a metal-ceramic crown or a bridge retainer, 521
- pulpitis – irreversible, The use of an intra-oral injection of ketorolac in the treatment of irreversible pulpitis, 789
- pulpitis – treatment, The use of an intra-oral injection of ketorolac in the treatment of irreversible pulpitis, 789
- pulpitis, The effect of capsaicin on substance P expression in pulp tissue inflammation, 30
- pulpless teeth, Continued root formation of a pulpless permanent incisor following root canal treatment: a case report, 489
- quality of endodontic treatment, Prevalence of apical periodontitis and the quality of endodontic treatment in an adult Belarusian population, 238
- quantitative rt-pcr, Reduction of osteocalcin expression in aged human dental pulp, 817
- RaCe, Comparison of root canal preparation using RaCe and ProTaper rotary Ni-Ti instruments, 8
- radicular cyst, Apoptosis in epithelial cells of apical radicular cysts, 465
- radiodensity, Calcium hydroxide dressings using different preparation and application modes: density and dissolution by simulated tissue pressure, 889
- radiography, A laboratory comparison of three imaging systems for image quality and radiation exposure characteristics, 645
- radioimmunoanalysis, The effect of capsaicin on substance P expression in pulp tissue inflammation, 30
- radioreceptor assay, Calcitonin gene-related peptide receptor expression in healthy and inflamed human pulp tissue, 712
- randomized controlled trial, Dental pulp capping: effect of Emdogain Gel on experimentally exposed human pulps, 186
- randomized controlled trial, The role of preoperative prophylactic antibiotic administration in periapical endodontic surgery: a randomized, prospective double-blind placebo-controlled study, 877
- rat, Behaviour of bone marrow osteoblast-like cells on mineral trioxide aggregate: morphology and expression of type I collagen and bone-related protein mRNAs, 203
- RC-Prep, Assessment of a gel-type chelating preparation containing 1-hydroxyethylidene-1, 1-bisphosphonate, 810
- release of ions, Physico-chemical properties of MTA and a novel experimental cement, 443
- replication, Intracanal temperature rise evaluation during the usage of the System B: replication of intracanal anatomy, 218
- resilon, Shear bond strength of Resilon to a methacrylate-based root canal sealer, 753
- resin blocks, Influence of different types of automated devices on the shaping ability of rotary nickel-titanium FlexMaster instruments, 627
- rests of malassez, Collagenase-3 (MMP-13) is expressed in periapical lesions: an immunohistochemical study, 297
- retreatment, Determining the apical terminus of root-end resected teeth using three modern apex locators: a comparative *ex vivo* study, 827

- retrofill, Longitudinal study on microleakage of three root-end filling materials by the fluid transport method and by capillary flow porometry, 129
- rheology, A study of the rheological properties of endodontic sealers, 499
- rhodamine b, Penetration of bonding resins into fibre-reinforced composite posts: a confocal microscopic study, 46
- root canal anatomy, Anatomic Endodontic Technology (AET) – a crown-down root canal preparation technique: basic concepts, operative procedure and instruments, 575
- root canal anatomy, Three-dimensional, non-destructive visualization of vertical root fractures using flat panel volume detector computer tomography: an *ex vivo in vitro* case report, 904
- root canal diameter, Comparative study of six rotary nickel–titanium systems and hand instrumentation for root canal preparation, 743
- root canal filling, The imaging of root canal obturation using micro-CT, 617
- root canal filling, An evaluation of .06 tapered gutta-percha cones for filling of .06 taper prepared curved root canals, 87
- root canal infection, Microscopic observation of bacteria: review highlighting the use of environmental SEM, 775
- root canal instrumentation, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- root canal instrumentation, Comparison of cold lateral compaction and continuous wave of obturation techniques following manual or rotary instrumentation, 381
- root canal length, A comparison between two electronic apex locators: an *in vivo* investigation, 36
- root canal obturation, The imaging of root canal obturation using micro-CT, 617
- root canal obturation, Presence of pores and vacuoles in set endodontic sealers, 690
- root canal preparation techniques, A comparison of spreader penetration depth and load required during lateral condensation in teeth prepared using various root canal preparation techniques, 510
- root canal preparation, Influence of different types of automated devices on the shaping ability of rotary nickel–titanium FlexMaster instruments, 627
- root canal preparation, Shaping ability of two rotary instruments in simulated canals: stainless steel ENDOfash and nickel–titanium HERO Shaper, 637
- root canal retreatment, Probability of removing fractured instruments from root canals, 112
- root canal retreatment, Effectiveness of different techniques for removing gutta-percha during retreatment, 2
- root canal sealer, *Ex vivo* study of the adhesion of an epoxy-based sealer to human dentine submitted to irradiation with Er : YAG and Nd : YAG lasers, 866
- root canal sealers, Short-term antibacterial activity of root canal sealers towards *Enterococcus faecalis*, 483
- root canal therapy, A laboratory comparison of three imaging systems for image quality and radiation exposure characteristics, 645
- root canal treatment, Garrè's osteomyelitis managed by root canal treatment of a mandibular second molar: incorporation of computed tomography with 3D reconstruction in the diagnosis and monitoring of the disease, 255
- root canal treatment, Continued root formation of a pulpless permanent incisor following root canal treatment: a case report, 489
- root canal treatment, A comparative study of Endo-flare–Hero Shaper and Mtwo NiTi instruments in the preparation of curved root canals, 610
- root canal treatment, A temporary filling material may cause cusp deflection, infractions and fractures in endodontically treated teeth, 653
- root canal treatment, Recovery of *Enterococcus faecalis* after single- or multiple-visit root canal treatments carried out in infected teeth *ex vivo*, 697
- root canal treatment, Technical quality of root canal treatment and detection of iatrogenic errors in an undergraduate dental clinic, 725
- root canal treatment, Determining the apical terminus of root-end resected teeth using three modern apex locators: a comparative *ex vivo* study, 827
- root canal treatment, Neurological complications following extrusion of sodium hypochlorite solution during root canal treatment, 843
- root canal, Survival of *Enterococcus faecalis* in root canals *ex vivo*, 735
- root canal, Influence of irrigant needle depth in removing bioluminescent bacteria inoculated into instrumented root canals using real-time imaging *in vitro*, 97
- root filling, Outcome of root canal treatment using Thermafil and cold lateral condensation filling techniques, 179
- root filling, Effect of customization of master gutta-percha cone on apical control of root filling using different techniques: an *ex vivo* study, 658
- root filling, Survival of *Enterococcus faecalis* in root canals *ex vivo*, 735
- root formation, Continued root formation of a pulpless permanent incisor following root canal treatment: a case report, 489
- root morphology, Root morphology of mandibular premolars in 40 patients with Turner syndrome, 822
- root perforation, Perforation repair with mineral trioxide aggregate: a modified matrix concept, 59
- root, Temperature rise of the post and on the root surface during ultrasonic post removal, 705

- root-end preparation, SEM analysis of the integrity of resected root apices of cadaver and extracted teeth after ultrasonic root-end preparation at different intensities, 310
- root-end resection, Frequency and type of canal isthmuses in first molars detected by endoscopic inspection during periradicular surgery, 160
- rotary instrument, Defects in ProTaper S1 instruments after clinical use: longitudinal examination, 550
- rotary instrument, Defects in ProTaper S1 instruments after clinical use: fractographic examination, 802
- rotary instrumentation, Effectiveness of different techniques for removing gutta-percha during retreatment, 2
- rotary instrumentation, Canal preparation with Hero 642 rotary Ni-Ti instruments compared with stainless steel hand K-file assessed using computed tomography, 402
- rotary instruments, Analysis of forces developed during mechanical preparation of extracted teeth using RaCe rotary instruments and ProFiles, 17
- rotary instruments, Influence of different types of automated devices on the shaping ability of rotary nickel-titanium FlexMaster instruments, 627
- rotary nickel-titanium file, Shaping ability of two rotary instruments in simulated canals: stainless steel ENDOflash and nickel-titanium HERO Shaper, 637
- rubber dam, The adoption of new endodontic technology amongst Danish general dental practitioners, 52
- sample preparation, Microscopic observation of bacteria: review highlighting the use of environmental SEMO, 775
- scanning electron microscopy, SEM analysis of the integrity of resected root apices of cadaver and extracted teeth after ultrasonic root-end preparation at different intensities, 310
- scanning electron microscopy, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- Sealapex, *In vitro* antimicrobial activity of Fill Canal, Sealapex, Mineral Trioxide Aggregate, Portland cement and EndoRez, 539
- SEM, CarisolvTM: an alternative to NaOCl in immature root canals?, 448
- separated instrument, Probability of removing fractured instruments from root canals, 112
- serine proteases, Effect of *E. faecalis* on the release of serine proteases elastase and cathepsin G, and collagenase-2 (MMP-8) by human polymorphonuclear leukocytes (PMNs), 667
- shaping, Anatomic Endodontic Technology (AET) – a crown-down root canal preparation technique: basic concepts, operative procedure and instruments, 575
- shear bond strength, Shear bond strength of Resilon to a methacrylate-based root canal sealer, 753
- shear, Defects in ProTaper S1 instruments after clinical use: fractographic examination, 802
- signal transduction pathway, Examination of the signal transduction pathways leading to upregulation of tissue type plasminogen activator by *Porphyromonas endodontalis* in human pulp cells, 860
- simulated canal, Shaping ability of two rotary instruments in simulated canals: stainless steel ENDOflash and nickel-titanium HERO Shaper, 637
- single cone, An evaluation of .06 tapered gutta-percha cones for filling of .06 taper prepared curved root canals, 87
- single visit, Recovery of *Enterococcus faecalis* after single- or multiple-visit root canal treatments carried out in infected teeth *ex vivo*, 697
- smear layer, The effect of application time of EDTA and NaOCl on intracanal smear layer removal: an SEM analysis, 285
- smooth wire, A comparison between a smooth wire and a K-file in removing artificially placed dentine debris from root canals in resin blocks during ultrasonic irrigation, 593
- sodium hypochlorite, The effect of application time of EDTA and NaOCl on intracanal smear layer removal: an SEM analysis, 285
- sodium hypochlorite, Effectiveness in cleaning oval-shaped root canals using Anatomic Endodontic Technology, ProFile and manual instrumentation: a scanning electron microscopic study, 356
- sodium hypochlorite, Effects of 2% chlorhexidine and 5.25% sodium hypochlorite on gutta-percha cones studied by atomic force microscopy, 425
- stabident, Pressures generated *in vitro* during Stabident intraosseous injections, 291
- stainless steel file, Shaping ability of two rotary instruments in simulated canals: stainless steel ENDOflash and nickel-titanium HERO Shaper, 637
- stress, Growth at high pH increases *Enterococcus faecalis* adhesion to collagen, 389
- stress, Periapical biomechanics and the role of cyclic biting force in apical retrograde fluid movement, 597
- substance p, The effect of capsaicin on substance P expression in pulp tissue inflammation, 30
- superelastic wires, Physical and mechanical characterization and the influence of cyclic loading on the behaviour of nickel-titanium wires employed in the manufacture of rotary endodontic instruments, 795
- superoxide dismutase, Copper–zinc superoxide dismutase activity in healthy and inflamed human dental pulp, 195
- surgical endodontics, Periapical resurgery versus periapical surgery: a 5-year longitudinal comparison, 320

- surgical endodontics, The role of preoperative prophylactic antibiotic administration in periapical endodontic surgery: a randomized, prospective double-blind placebo-controlled study, 877
- survival analysis, Fate of vital pulps beneath a metal-ceramic crown or a bridge retainer, 521
- survival, Survival of *Enterococcus faecalis* in root canals *ex vivo*, 735
- system b, Comparison of cold lateral compaction and continuous wave of obturation techniques following manual or rotary instrumentation, 381
- technical quality, Technical quality of root canal treatment and detection of iatrogenic errors in an undergraduate dental clinic, 725
- temperature, Intracanal temperature rise evaluation during the usage of the System B: replication of intracanal anatomy, 218
- temperature, Temperature rise of the post and on the root surface during ultrasonic post removal, 705
- temporary filling, A temporary filling material may cause cusp deflection, inflections and fractures in endodontically treated teeth, 653
- temporization, A temporary filling material may cause cusp deflection, inflections and fractures in endodontically treated teeth, 653
- therapeutic efficacy, Effectiveness of single- versus multiple-visit endodontic treatment of teeth with apical periodontitis: a systematic review and meta-analysis, 347
- thermafil, Outcome of root canal treatment using ThermoFil and cold lateral condensation filling techniques, 179
- thermoplasticized, Intracanal temperature rise evaluation during the usage of the System B: replication of intracanal anatomy, 218
- three-dimensional observation, Three-dimensional, non-destructive visualization of vertical root fractures using flat panel volume detector computer tomography: an *ex vivo in vitro* case report, 904
- tissue type plasminogen activator, Examination of the signal transduction pathways leading to upregulation of tissue type plasminogen activator by *Porphyromonas endodontalis* in human pulp cells, 860
- tissue-type plasminogen activator, Upregulation of tissue-type plasminogen activator in inflamed human dental pulps, 328
- tooth fracture, A temporary filling material may cause cusp deflection, inflections and fractures in endodontically treated teeth, 653
- torque value, Effect of liquid and paste-type lubricants on torque values during simulated rotary root canal instrumentation, 223
- torque, Analysis of forces developed during mechanical preparation of extracted teeth using RaCe rotary instruments and ProFiles, 17
- transportation, Root canal preparation with Endo-Eze AET: changes in root canal shape assessed by micro-computed tomography, 456
- treatment effectiveness, Effectiveness of single- versus multiple-visit endodontic treatment of teeth with apical periodontitis: a systematic review and meta-analysis, 347
- treatment sessions, The adoption of new endodontic technology amongst Danish general dental practitioners, 52
- Turner syndrome, Root morphology of mandibular premolars in 40 patients with Turner syndrome, 822
- ultrasonic irrigation, A comparison between a smooth wire and a K-file in removing artificially placed dentine debris from root canals in resin blocks during ultrasonic irrigation, 593
- ultrasonic preparation, SEM analysis of the integrity of resected root apices of cadaver and extracted teeth after ultrasonic root-end preparation at different intensities, 310
- ultrasonics, Probability of removing fractured instruments from root canals, 112
- ultrasound, Periapical resurgery versus periapical surgery: a 5-year longitudinal comparison, 320
- ultrasound, Temperature rise of the post and on the root surface during ultrasonic post removal, 705
- undergraduate dental clinic, Technical quality of root canal treatment and detection of iatrogenic errors in an undergraduate dental clinic, 725
- vacuolar structure, Presence of pores and vacuoles in set endodontic sealers, 690
- vacuoles, Presence of pores and vacuoles in set endodontic sealers, 690
- vertical load, Analysis of forces developed during mechanical preparation of extracted teeth using RaCe rotary instruments and ProFiles, 17
- vertical root fractures, Three-dimensional, non-destructive visualization of vertical root fractures using flat panel volume detector computer tomography: an *ex vivo in vitro* case report, 904
- warm gutta-percha, The adoption of new endodontic technology amongst Danish general dental practitioners, 52
- whitening strip, *In vitro* peroxide penetration into the pulp chamber from newer bleaching products, 516
- working length, Comparative study of six rotary nickel-titanium systems and hand instrumentation for root canal preparation, 743
- working length, Determining the apical terminus of root-end resected teeth using three modern apex locators: a comparative *ex vivo* study, 827

- working length, *Ex vivo* comparison of two electronic apex locators with different scales and frequencies, 855
- working safety, Comparative study of six rotary nickel-titanium systems and hand instrumentation for root canal preparation, 743
- working time, Comparative study of six rotary nickel-titanium systems and hand instrumentation for root canal preparation, 743
- Xchromosome, Root morphology of mandibular premolars in 40 patients with Turner syndrome, 822
- X-ray diffraction, Effect of cryogenic treatment on nickel-titanium endodontic instruments, 364
- yeasts, Effect of endodontic procedures on enterococci, enteric bacteria and yeasts in primary endodontic infections, 372

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.