doi:10.1111/j.1365-2591.2011.01929.x

# The effects of hyperosmosis or high pH on a dual-species biofilm of *Enterococcus faecalis* and *Pseudomonas aeruginosa*: an *in vitro* study

## S. V. van der Waal<sup>1</sup>, L. W. M. van der Sluis<sup>2</sup>, A. R. Özok<sup>1</sup>, R. A. M. Exterkate<sup>1</sup>, J. van Marle<sup>3</sup>, P. R. Wesselink<sup>1</sup> & J. J. de Soet<sup>1</sup>

<sup>1</sup>Department of Cariology, Endodontology, Pedodontology and Oral Microbiology, Academic Centre for Dentistry Amsterdam, Amsterdam, The Netherlands; <sup>2</sup>Department of Conservative Dentistry and Endodontology, Faculty of Dentistry, Paul Sabatier University, Toulouse, France; and <sup>3</sup>Department of Cell Biology and Histology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

#### Abstract

van der Waal SV, van der Sluis LWM, Özok AR, Exterkate RAM, van Marle J, Wesselink PR, de Soet JJ. The effects of hyperosmosis or high pH on a dual-species biofilm of *Enterococcus faecalis* and *Pseudomonas aeruginosa*: an *in vitro* study. *International Endodontic Journal*, **44**, 1110– 1117, 2011.

**Aim** To investigate the effect of hyperosmotic hyperosmosis or alkaline stress on a dual-species biofilm of *Enterococcus faecalis* and *Pseudomonas aeruginosa*.

**Methodology** Biofilms were grown on glass cover slips suspended in bacterial inoculate for 96 h, after which the cover slips with attached biofilms were immersed in brain heart infusion broth (BHI-broth) with 6 mol  $L^{-1}$  sodium chloride (NaCl) representing the hyperosmotic group or Ca(OH)<sub>2</sub>, pH 12.1, representing the alkaline group. Two per cent sodium hypochlorite and BHI- broth served as positive and negative controls, respectively. After treatment, the biofilms were washed, harvested and plated on bloodagar plates after serial dilution. The bactericidal effect was assessed by determining the colony-forming units (CFU). The effect on the biofilm mass was imaged with confocal laser scanning microscopy (CLSM).

**Results** Hyperosmosis reduced the CFU of both species significantly after 72 h (P < 0.0001). After 168 h, *P. aeruginosa* was eradicated and the *E. faecalis* reduction was more than 99%. High pH could not induce a significant bacterial reduction. CLSM revealed dense flocculation of the biofilms incubated in alkaline broth. **Conclusion** Hyperosmosis effectively reduced a dual-species biofilm of *E. faecalis* and *P. aeruginosa*, whilst high pH had limited bactericidal effect in this model.

**Keywords:** biofilm, calcium hydroxide, disinfection, *Enterococcus faecalis*, hyperosmotic stress, *Pseudomonas aeruginosa*.

Received 28 April 2010; accepted 5 July 2011

#### Introduction

1110

Incomplete disinfection of the root canal system is considered to be a major cause of post-treatment disease (Seltzer *et al.* 1967, Lin *et al.* 1992). Hence, elimination of infection is one of the main goals of root canal treatment. However, after contemporary nonsurgical root canal treatment, studies confirm the presence of persisting microorganisms (Nair *et al.* 2005, Ricucci & Siqueira 2010) likely due to the complex architecture of the root canal system (Vertucci 1984, Wu *et al.* 2000, Fan *et al.* 2004). Furthermore, microorganisms live in a biofilm configuration, which renders them less susceptible to endodontic treatment procedures (Portenier *et al.* 2005). A biofilm is a microbial (bacterial, fungal, algal) community, enveloped by extracellular polysaccharides (EPS) that these microbial cells

Correspondance: Suzette V. van der Waal, Department of Cariology, Endodontology, Pedodontology and Oral Microbiology, Gustav Mahlerlaan 3004, 1081 LA Amsterdam, The Netherlands (e-mail: s.vd.waal@acta.nl).

produce, which adheres to the interface of a liquid or a surface. Biofilms might contain 95-99% EPS and 1-5% microorganisms (Körstgens *et al.* 2001).

Osmosis is a physical phenomenon whereby solvent (water) moves, without input of energy, across a semipermeable membrane (permeable to the solvent, but not the solute) separating two solutions of different concentrations, until a certain equilibrium is reached. Hyperosmosis reduces the water activity of microorganisms and their environment. Hyperosmotic metabolic stress inhibits growth and eventually kills microorganisms (Wijnker *et al.* 2006, Bautista-Gallego *et al.* 2008).

Calcium hydroxide  $(Ca(OH)_2)$  is used as an intracanal dressing to prevent re-growth of microorganisms during the inter-appointment period between two treatment sessions and for additional reduction in the microbial load after chemo-mechanical cleaning of the root canal system. The antimicrobial action is based on alkaline stress, because of its pH of approximately 12 (Siqueira & Lopes 1999). However, whilst some studies report bactericidal action of Ca(OH)<sub>2</sub> (Byström *et al.* 1985, Sjögren *et al.* 1991, Gubler *et al.* 2008), others report that its antibacterial effect is insufficient (Haapasalo & Ørstavik 1987, Peters *et al.* 2002).

The aim of this study was to evaluate *in vitro* the effect of hyperosmosis or high pH on a dual-species biofilm of *Enterococcus faecalis* and *Pseudomonas aeru-ginosa*. Viability of the biofilms was assessed by conventional plate counting, whilst the effects on the EPS of the biofilms were demonstrated with confocal laser scanning microscopy (CLSM) imaging.

#### **Materials and methods**

The biofilm model used here was described by Deng *et al.* (2009). In brief, round glass coverslips attached to a custom-made stainless steel lid were suspended in a medium inoculated with bacteria. In this model, biofilm attachment and growth is an active process against gravity.

#### Microbiology

Cells of *E. faecalis* V583 and *P. aeruginosa* HG 4950, both clinical isolates, were grown and maintained as pure cultures on blood agar plates. *E. faecalis* was cultured under anaerobic conditions (80% N<sub>2</sub>, 10% H<sub>2</sub> and 10% CO<sub>2</sub>) whilst *P. aeruginosa* was cultured in air, both at 37 °C. Growth media for overnight cultures and biofilms were BHI (Oxoid, Basingstoke, UK) and BHI broth containing 0.5 BHI, 50 mmol L<sup>-1</sup> 1,4piperazinediethanesulfonic acid, buffer (Sigma, St. Louis, MO, USA), 5% saccharose and 1 mmol L<sup>-1</sup> Ca<sup>2+</sup> during biofilm growth. In the experimental phase, 6 mol L<sup>-1</sup> sodium chloride (NaCl; Merck, Darmstadt, Germany) was added to the BHI broth (pH 6.9) to raise the osmotic value. Addition of NaCl did not alter the pH. Ca(OH)<sub>2</sub> of 0.02 mol L<sup>-1</sup> (to reach saturation) was added to BHI broth for the alkaline treatment (pH 12.1). Two per cent sodium hypochlorite (NaOCl; Merck) was used for the positive control and 2% sodium thiosulphate (STS) in peptone water [1% peptone (Oxoid) in water] for neutralizing the NaOCl. Phosphate-buffered saline (PBS; Oxoid) was used to wash the negative controls, as a neutralizing agent for NaCl and Ca(OH)<sub>2</sub> and for serial dilution.

#### Biofilm growth for viability assessment - CFU

Fresh cultures of each microorganism were grown overnight at 37 °C in air. Numbers of colony-forming units (CFU) were estimated by optical density measurement at 600 nm based on previously performed plate counts. One hundred microlitres with approximately  $2 \times 10^9$  *E. faecalis* and 100 µL with approximately  $2 \times 10^7$  *P. aeruginosa* were inoculated in wells of a 24-well multiwell plate (Greiner Bio-one, Frickenhausen, Germany) containing 1.5 mL BHI broth. Round glass coverslips (Thermo Scientific, Braunschweig, Germany), diameter 12 mm, mounted on a custom-made stainless steel lid, were suspended in the BHI broth.

In this experiment, 96-h biofilms were used because after 96 h, the biomass in a biofilm increases whilst the quantity of living cells is more or less established (Wilson *et al.* 1996). The broth was refreshed daily. Inoculation of bacteria and the biofilm growth took place at 37 °C in air. Biofilm growth was confirmed after swabbing a coverslip and streaking the biofilm onto a glass microscope slide. After heat fixation of the glass slides, the samples were gram-stained. Biofilm growth was confirmed after examination of the glass slides with a reflected-light microscope (1000× magnification; Zeiss Axiolab, Axiolab A, Carl Zeiss B.V., Sliedrecht, The Netherlands).

### Biofilm growth for biofilm matrix assessment – CLSM

The procedure is the same as described above, except that the biofilms used for CLSM imaging were grown for 24 h.

#### **Experiments: CFU**

After 96 h of growth, the coverslips with the attached biofilms were immersed in hyperosmotic or alkaline broth and incubated for 4, 24, 48, 72 and 168 h. The negative controls were kept in BHI broth. Where applicable, the incubation fluids were refreshed daily. Immersion in 2% NaOCl for 10 min served as a positive control. After the experiments, the coverslips were removed from the lid and immersed in 20 mL of neutralizing agents: 10 min in PBS for the NaCl groups, Ca(OH)<sub>2</sub> groups and negative controls and 30 min in STS for the positive controls. Biofilms were harvested using a sterile cotton swab and resuspended in 1 mL of PBS by vigorously moving the cotton swab up and down fifty times in a 1.5mL test tube (Standard Micro Test Tube 3810; Eppendorf AG, Hamburg, Germany). Serial dilutions of the suspension were plated and incubated for 48 h in the absence of oxygen. Each group consisted of three specimens, and each experiment was performed in triplicate.

#### **Experiments: CLSM imaging**

Twenty-four-hour biofilms were incubated for 4 h in hyperosmotic or alkaline broth after which the biofilms were washed twice in 2 mL PBS to remove nonadherent cells. Biofilms were stained in duplicate for 15 min with Calcofluor White 50 mmol  $L^{-1}$  (CW) in a dark chamber at room temperature. CW is a polysaccharidebinding dye, used to stain the extracellular matrix of biofilms formed by bacteria (Stewart *et al.* 1994, Neut *et al.* 2010). Of each sample, three standardized sites were chosen for imaging.

For imaging, a Leica DM-IRB (Leica Microsystems CMS GmbH, Mannheim, Germany) inverted microscope/SP2-AOBS confocal system was used. Images (512\*512 pixels) were acquired with a  $40\times/NA$  1.25 HCX PL APO oil objective. The final pixel size was 0.776 µm. The pinhole size was set at 1 airy unit, resulting in a Z resolution of approximately 0.5 µm. Image stacks were acquired with a stepsize of 0.3 µm. Excitation/detection of CW was achieved with 388-nm excitation and with 440-nm detection. Using gain and offset of the photo-multiplying tube, the images were adapted to the full (8 bit) dynamic range of the system. Presence or absence of EPS was observed by CLSM.

#### Statistical analysis

1112

Colony-forming units counts were logarithmically transformed. Data were analysed by one-way analysis

of variance (ANOVA) to compare the experimental groups with the negative controls. The significance level was set at  $\alpha = 0.05$ . The analyses were carried out with the statistical package SPSS version 12 (SPSS Inc. Chicago, IL, USA).

#### Results

#### Viability of the biofilms

The results are presented as log CFU reduction per biofilm compared with the negative controls (Fig. 1). The negative controls contained  $8.1 \pm 0.6$  log (CFU) microorganisms per biofilm, which gradually increased in time. The biofilms used for the CLSM experiment, 24 h, contained on average  $3.2 \times 10^7$  cells.

The mean log reduction in CFU in the hyperosmotic groups for both *E. faecalis* and *P. aeruginosa* after 72 h and after 168 h was 6 and 8, respectively, and was statistically significant compared with the negative controls. At 168 h, the control biofilms contained on average  $1.5 \times 10^9$  CFU of *P. aeruginosa*, of which none could be detected after 168 h of hyperosmotic treatment. *E. faecalis* was reduced to a mean of 40 CFU. The reductions were highly significant for both species (*P* < 0.001) and showed a significant linear trend in time.

Alkaline broth reduced *P. aeruginosa* significantly (P = 0.02); however, the reduction did not increase with an increased application time. Reduction of *E. faecalis* by alkali was not significant (P = 0.09). After 72 h, a log 2 reduction was determined for both species, but after 168 h, bacterial growth recovered until a mean log 1 reduction was obtained.

The NaOCl groups showed 100% kill after 10-min exposure. With the naked eye, the glass coverslips exposed to hyperosmotic broth showed no signs of a biofilm plaque after treatment. Increase in biofilm plaque was an obvious observation in the alkaline broth group (Fig. 2).

#### CLSM assessment of biofilm matrix

The CLSM images show the bacterial cells and the EPS. CW does not give an indication of the viability of the cells. The biofilm incubated in hyperosmotic broth (Fig. 3) showed reduced amount of biofilm mass compared with the control biofilm (Fig. 4). CLSM clearly indicated enhanced biofilm mass production in biofilms incubated in alkaline broth (Fig. 5) compared with the control biofilm. This is in line with the naked-eye observations.



**Figure 1** The reduction in colony-forming units (CFU) counts of bacterial species *Enterococcus faecalis* and *Pseudomonas aeruginosa* in log CFU/biofilm after exposure (hours) to hyperosmotic or alkaline broth. The negative controls contained on average  $2.0 \times 10^8$  CFU *E. faecalis* and  $3.5 \times 10^8$  CFU *P. aeruginosa*, which numbers increased in time.



**Figure 2** Illustrative photographs of glass cover slips fixed with green silicone impression material (President<sup>®</sup>) in white clamps on the custom-made stainless steel lid, with biofilms after 96 + 24 h control (left), after 24 h incubation in hyperosmotic broth (middle) or 24 h incubation in alkaline broth (right). The thickness of the plaque (right) increased throughout the experiment. The plaque seemed petrified and was firmly attached to the cover slip.

Furthermore, the appearance of the cells in the hyperosmotic group seems altered in size and shape.

#### Discussion

In this biofilm model, hyperosmotic broth reduced the viability of cells in a biofilm significantly compared with the negative controls (Fig. 1). When raising the osmotic value, there was a selective bactericidal effect with *P. aeruginosa* being more susceptible to raising the osmotic value than *E. faecalis* at earlier time points

(Fig. 1). This was expected because bacterial species respond differently to environmental stresses. It was reported that *P. aeruginosa* does not tolerate 6.5% NaCl (Gilardi 1969) whilst *E. faecalis* was reported to be more resistant to NaCl (Lee 1972, Faklam 1973).

The mechanism of hyperosmotic metabolic stress killing of bacteria has been demonstrated earlier (Lee 1972, Faklam 1973). Microorganisms counteract the emigration of free intracellular water owing to the extracellular hyperosmotic value, by restoring the intracellular solute concentration. Restoration





**Figure 3** Confocal laser scanning microscopy image of 24 h biofilm after incubation in hyperosmotic broth for 4 h. Cells show more fluorescence which suggests that they are reduced in size and the cells are not embedded in extracellular polysaccharides (EPS). EPS material is reduced to small spots on the substrate.

**Figure 4** Confocal laser scanning microscopy image of negative control biofilm stained with Calcofluor White. Bacteria are visible as rod and cocci-shaped structures embedded in extracellular polysaccharides.

occurs through the uptake or synthesis of small ions and molecules, which do not interfere with the cell's metabolism (Csonka 1989, Gutierrez *et al.* 1995,

**Figure 5** Confocal laser scanning microscopy image of 24 h biofilm after incubation in alkaline broth for 4 h shows a dense cluster of extracellular polysaccharides (EPS). Within the EPS, the cells are difficult to distinguish.

Record et al. 1998). Whether extracellular high osmotic value is lethal depends on the ability of the microorganisms to adapt to the changed osmotic environment, the available resources of compatible solutes and energy and duration of the hyperosmotic change. Storing and synthesizing compatible solutes are energy-demanding processes. Once the stocks of compatible solutes or energy resources are depleted, microbes enter the stationary phase, after which eventually bacterial death follows (Csonka & Hanson 1991). The phenomenon of osmosis is solely dependent on the concentration of solutes. The attraction of solvents by the solutes continues until certain equilibrium has been reached. In theory, a hyperosmotic root canal dressing would raise the osmotic value in the root canal. Because this high ionic concentration will be equilibrated with the surrounding tissues, some effect might perhaps be expected in microorganisms located more distant from the main root canal. The mode of action at a distance was described in a study by Wijnker et al. (2006), where multispecies bacterial cultures were inoculated inside pieces of intestine. The sample ends were tied, after which the samples were disinfected in  $6.0 \text{ mol } \text{L}^{-1}$ brine.

pH 12.1 did not significantly reduce bacterial numbers in this model. Calcium hydroxide has some bactericidal effect in planktonic cultures (Stuart *et al.* 1991, Ferreira *et al.* 2007). Microorganisms, however, reside in the root canal system in a biofilm configuration (Nair 1987, Chávez de Paz 2007a, Ricucci & Siqueira 2010). Studies investigating the antibacterial action of Ca(OH)<sub>2</sub> on *E. faecalis* biofilms show that *E. faecalis* is resistant to Ca(OH)<sub>2</sub> (Abdullah *et al.* 2005, Chávez de Paz *et al.* 2007b). Its decreased susceptibility not only might be caused by genetic adaption of the biofilm cell but might also be a result of increased production of EPS in which the cells can shelter from the alkaline environment.

A dual-species biofilm was used because of its increased resistance to antimicrobials (Kara et al. 2007, Özok et al. 2007, Jiang et al. 2011). E. faecalis, facultative gram-positive cocci, was chosen as the test organism because of its reported association with persistent periapical pathosis (Sundqvist et al. 1998, Siqueira & Rôças 2004, Stuart et al. 2006). Factors facilitating its presence in filled root canals with persistent lesions include the ability to invade dentinal tubules (Haapasalo & Ørstavik 1987, Weiger et al. 2002), to tolerate the alkaline pH of  $Ca(OH)_2$  (Evans et al. 2002, Nakajo et al. 2006) and to withstand long periods of starvation with subsequent recovery in the presence of serum (Figdor et al. 2003). E. faecalis grows easily in a laboratory setting. Furthermore, E. faecalis is known to efficiently accumulate the compatible solutes from the medium to counteract increased osmotic pressure (Pichereau et al. 1999). P. aeruginosa is an aerobic gram-negative motile rod. It is a notorious EPS producer and for this reason commonly chosen as model bacterium in biofilm studies (Bazire et al. 2006, Toté et al. 2010). No matter how smooth a surface, P. aeruginosa adheres within 30 s (Vanhaecke et al. 1990). Although classified as aerobic, P. aeruginosa survives anaerobic conditions (Yoon et al. 2002). Its presence in the root canal is associated with failed endodontic treatment (Ranta et al. 1988, Sirén et al. 1997).

In pilot studies performed prior to this study, varying ratios of *P. aeruginosa* and *E. faecalis* were cultured together to determine whether and in which composition the species would grow together in a biofilm. The ratio of 1 cell of *P. aeruginosa* to 100 cells of *E. faecalis* yielded a reproducible dual-species biofilm and was therefore used to inoculate this model.

Biocides and disinfectants work by killing microorganisms, a strategy that is rendered ineffective by the reduced susceptibility of biofilm microorganisms to antimicrobial challenges (Stewart *et al.* 2000). This study focused on biofilm killing, measured by the number of viable microorganisms attached to the glass coverslip after treatment. Plate count of CFU, however, combines the effects of biofilm removal and killing of bacterial cells (Chen & Stewart 2000).

The addition of  $Ca^{2+}$  ions to the growth medium is known to enhance the attachment of the biofilm cell to the substrate (Geesey et al. 2000) and to induce the production of EPS (Patrauchan et al. 2005) as well as to increase the Young's modulus of the EPS (Körstgens et al. 2001). It has also been demonstrated that raising the pH of the growth medium in the presence of divalent cations has an effect on the EPS of bacteria and yeasts. By raising the pH, divalent cations bind to binding sites of the polysaccharide strands formerly occupied by H<sup>+</sup> (Geddie & Sutherland 1993). These findings might explain the increase in biofilm plaque and its petrified appearance in the present model after incubation with Ca(OH)2containing broth (Fig. 2). The sterile coverslips were also incubated in alkaline broth and showed very little precipitate on the coverslips after a week's incubation (not reported here). This suggests an active role of the bacteria.

Sodium hypochlorite was used in this study in the positive control group and killed all bacteria in the biofilms. In the complexity of the root canal system, however, NaOCl does not remove all biofilm and microorganisms (Nair *et al.* 2005). Probably, this is because of the difficulty of dispersing NaOCl during irrigation into the complete root canal system, especially into the apical part of the root canal system (Boutsioukis *et al.* 2010).

Application of an inter-appointment root canal dressing, after chemo-mechanical preparation with NaOCl, with a mode of action based on hyperosmotic, metabolic stress might possibly result in a decrease in the microorganismal load either by killing in a biofilm or by removing the EPS. This concept needs further investigation in models that resemble the root canal.

#### Conclusions

Raising the osmotic value of growth medium has a bactericidal effect as well as an antibiofilm effect and might therefore seem promising for future use as an inter-appointment root canal dressing. High pH induced by calcium hydroxide was not effective in this model. Whether a hypertonic dressing affects biofilms in the complexity of a root canal system needs to be investigated in future studies.

#### References

- Abdullah M, Ng YL, Gulabivala K, Moles DR, Spratt DA (2005) Susceptibilities of two *Enterococcus faecalis* phenotypes to root canal medications. *Journal of Endodontics* **31**, 30–6.
- Bautista-Gallego J, Arroyo-López FN, Durán-Quintana MC, Garrido-Fernández A (2008) Individual effects of sodium, potassium, calcium, and magnesium chloride salts on *Lactobacillus pentosus* and *Saccharomyces cerevisiae* growth. *Journal of Food Protection* **71**, 1412–21.
- Bazire A, Diab F, Jebbar M, Haras D (2006) Influence of high salinity on biofilm formation and benzoate assimilation by *Pseudomonas aeruginosa. Journal of Industrial Microbiology and Biotechnology* **34**, 5–8.
- Boutsioukis C, Verhaagen B, Kastrinakis E, Versluis M, van der Sluis LWM (2010) Evaluation of irrigant flow in the root canal using different needle types by an unsteady Computational Fluid Dynamics model. *Journal of Endodontics* 36, 875–9.
- Byström A, Claesson R, Sundqvist G (1985) The antibacterial effect of camphorated paramonochlorophenol, camphorated phenol and calcium hydroxide in the treatment of infected root canals. *Endodontic and Dental Traumatology* **1**, 170–5.
- Chávez de Paz LE (2007a) Redefining the persistent infection in root canals: possible role of biofilm communities. *Journal* of Endodontics **33**, 652–62.
- Chávez de Paz L, Bergenholtz G, Dahlén G, Svensäter G (2007b) Response to alkaline stress by root canal bacteria in biofilms. *International Endodontic Journal* **40**, 344–55.
- Chen X, Stewart PS (2000) Biofilm removal caused by chemical treatments. Water Research 34, 4229–33.
- Csonka LN (1989) Physiological and genetic responses of bacteria to osmotic stress. *Microbiological Reviews* 53, 121– 47.
- Csonka LN, Hanson AD (1991) Prokaryotic osmoregulation: genetics and physiology. Annual Review of Microbiology 45, 569–606.
- Deng DM, Hoogenkamp MA, Exterkate AM et al. (2009) Influence of Streptococcus mutans on Enterococcus faecalis biofilm formation. Journal of Endodontics 35, 1249–52.
- Evans M, Davies JK, Sundqvist G, Figdor D (2002) Mechanisms involved in the resistance of *Enterococcus faecalis* to calcium hydroxide. *International Endodontic Journal* **35**, 221–8.
- Faklam RR (1973) Comparison of several laboratory media for presumptive identification of Enterococci and Group D Streptococci. Applied Microbiology 26, 138–45.
- Fan B, Cheung GS, Fan M, Gutmann JL, Bian Z (2004) Cshaped canal system in mandibular second molars: part I – anatomical features. *Journal of Endodontics* **30**, 899–903.
- Ferreira FB, Torres SA, Rosa OP et al. (2007) Antimicrobial effect of propolis and other substances against selected endodontic pathogens. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology 104, 709– 16.

- Figdor D, Davies JK, Sundqvist G (2003) Starvation survival, growth and recovery of *Enterococcus faecalis* in human serum. *Oral Microbiology and Immunology* **18**, 234–9.
- Geddie JL, Sutherland IW (1993) Uptake of metals by bacterial polysaccharides. *Journal of Applied Bacteriology* **74**, 467–72.
- Geesey GG, Wigglesworth-Cooksey B, Cooksey KE (2000) Influence of calcium and other cations on surface adhesion of bacteria and diatoms: a review. *Biofouling* **15**, 95–205.
- Gilardi LG (1969) Evaluation of media for differentiating nonfermenting gram-negative bacteria of medical significance. *Applied Microbiology* 18, 355–9.
- Gubler M, Brunner TJ, Zehnder M, Waltimo T, Sener B, Stark WJ (2008) Do bioactive glasses convey a disinfecting mechanism beyond a mere increase in pH? *International Endodontic Journal* **41**, 670–8.
- Gutierrez C, Abee T, Booth IR (1995) Physiology of the osmotic stress response in microorganisms. *International Journal of Food Microbiology* 28, 233–44.
- Haapasalo M, Ørstavik D (1987) In vitro Infection and disinfection of dentinal tubules. Journal of Dental Research 66, 1375–9.
- Jiang LM, Hoogenkamp MA, van der Sluis LWM, Wesselink PR, Crielaard W, Deng DM (2011) Resazurin metabolism assay for root canal disinfectant evaluation on dual-species biofilms. *Journal of Endodontics* **37**, 31–4.
- Kara D, Luppens SB, van Marle J, Özok R, Ten Cate JM (2007) Microstructural differences between single-species and dualspecies biofilms of *Streptococcus mutans* and *Veillonella parvula*, before and after exposure to chlorhexidine. *FEMS Microbiology Letters* 271, 90–7.
- Körstgens V, Flemming HC, Wingender J, Borchard W (2001) Influence of calcium ions on the mechanical properties of a model biofilm of mucoid *Pseudomonas aeruginosa*. Water Science and Technology **43**, 49–57.
- Lee WS (1972) Improved procedure for identification of group D Enterococci with two new media. *Applied Microbiology* **24**, 1–3.
- Lin LM, Skribner JE, Gaengler P (1992) Factors associated with endodontic treatment failures. *Journal of Endodontics* 18, 625–7.
- Nair PNR (1987) Light and electron microscopic studies of root canal flora and periapical lesions. *Journal of Endodontics* 13, 29–39.
- Nair PN, Henry S, Cano V (2005) Microbial status of apical root canal system of human mandibular first molars with primary apical periodontitis after 'one-visit' endodontic treatment. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics **99**, 231–52.
- Nakajo K, Komori R, Ishikawa S *et al.* (2006) Resistance to acidic and alkaline environments in the endodontic pathogen *Enterococcus faecalis*. Oral Microbiology and Immunology 21, 283–8.
- Neut D, Hendriks JGE, Van Horn JR, Van der Mei HC, Busscher HJ (2010) *Pseudomonas aeruginosa* biofilm formation and

1116

slime excretion on antibiotic-loaded bone cement. *Acta Orthopaedica* **76**, 109–14.

- Özok AR, Wu MK, Luppens SB, Wesselink PR (2007) Comparison of growth and susceptibility to sodium hypochlorite of mono- and dual-species biofilms of *Fusobacterium* nucleatum and Peptostreptococcus (micromonas) micros. Journal of Endodontics **33**, 819–22.
- Patrauchan MA, Sarkisova S, Sauer K, Franklin MJ (2005) Calcium influences cellular and extracellular product formation during biofilm-associated growth of a marine *Pseudoalteromonas* sp. *Microbiology* **151**, 2885–97.
- Peters LB, Van Winkelhoff AJ, Buijs JF, Wesselink PR (2002) Effects of instrumentation, irrigation and dressing with calcium hydroxide on infection in pulpless teeth with periapical bone lesions. *International Endodontic Journal* **35**, 13–21.
- Pichereau V, Bourot S, Flahaut S, Blanco C, Auffray Y, Bernard T (1999) The osmoprotectant glycine betaine inhibits salt-induced cross-tolerance towards lethal treatment in *Enterococcus faecalis*. *Microbiology* **145**, 427–35.
- Portenier I, Waltimo T, Ørstavik D, Haapasalo M (2005) The susceptibility of starved, stationary phase, and growing cells of *Enterococcus faecalis* to endodontic medicaments. *Journal of Endodontics* **31**, 380–6.
- Ranta K, Haapasalo M, Ranta H (1988) Monoinfection of root canal with *Pseudomonas aeruginosa*. *Endodontics and Dental Traumatology* **4**, 269–72.
- Record MT Jr, Courtenay ES, Cayley DS, Guttman HJ (1998) Responses of *E. coli* to osmotic stress: large changes in amounts of cytoplasmic solutes and water. *Trends in Biochemical Sciences* 23, 143–8.
- Ricucci D, Siqueira JF Jr (2010) Biofilms and apical periodontitis: study of prevalence and association with clinical and histopathologic findings. *Journal of Endodontics* **36**, 1277–88.
- Seltzer S, Bender IB, Smith J, Freedman I, Nazimov H (1967) Endodontic failures-an analysis bases on clinical, roentgenographic, and histologic findings I & II. Oral Surgery, Oral Medicine and Oral Pathology 23, 500–30.
- Siqueira JF Jr, Lopes HP (1999) Mechanisms of antimicrobial activity of calcium hydroxide: a critical review. *International Endodontic Journal* 32, 361–9.
- Siqueira JF Jr, Rôças IN (2004) Polymerase chain reactionbased analysis of microorganisms associated with failed endodontic treatment. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology 97, 85–94.
- Sirén EK, Haapasalo MP, Ranta K, Salmi P, Kerosuo EN (1997) Microbiological findings and clinical treatment procedures in endodontic cases selected for microbiological investigations. *International Endodontic Journal* **30**, 91–5.
- Sjögren U, Figdor D, Spångberg L, Sundqvist G (1991) The antimicrobial effect of calcium hydroxide as a short-term

intracanal dressing. International Endodontic Journal **24**, 119–25.

- Stewart PS, Murga R, Srinivasan R, De Beer D (1994) Biofilm structural heterogeneity visualized by three microscopic methods. *Water Research* 8, 2006–9.
- Stewart PS, McFeters GA, Huang CT (2000) Biofilm control by antimicrobial agents. In: Bryers JD, ed. Biofilms II: Process Analysis and Applications. New York: Wiley, pp. 373–405.
- Stuart KG, Miller CH, Brown CE Jr, Newton CW (1991) The comparative antimicrobial effect of calcium hydroxide. Oral Surgery, Oral Medicine and Oral Pathology 72, 101–4.
- Stuart CH, Schwartz SA, Beeson TJ, Owatz CB (2006) Enterococcus faecalis: its role in root canal treatment failure an current concepts in retreatment. Journal of Endodontics 32, 93–8.
- Sundqvist G, Figdor D, Persson S, Sjögren U (1998) Microbiologic analysis of teeth with failed endodontic treatment and the outcome of conservative re-treatment. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology 85, 86–93.
- Toté K, Horemans T, Vanden Berghe D, Maes L, Cos P (2010) Inhibitory effect of biocides on viable mass and matrix of *Staphylococcus aureus* and *Pseudomonas aeruginosa* biofilms. *Applied and Environmental Microbiology* **76**, 3135–42.
- Vanhaecke E, Remon JP, Moors M, Raes F, De Rudder D, Van Peteghem A (1990) Kinetics of *Pseudomonas aeruginosa* adhesion to 304 and 316-L stainless steel: role of cell surface hydrophobicity. *Applied Environmental Microbiology* 56, 788–95.
- Vertucci FJ (1984) Root canal anatomy of the human permanent teeth. Oral Surgery, Oral Medicine and Oral Pathology 58, 589–99.
- Weiger R, de Lucena J, Decker HE, Löst C (2002) Vitality status of microorganisms in infected human root dentine. *International Endodontic Journal* 35, 166–71.
- Wijnker JJ, Koop G, Lipman LJA (2006) Antimicrobial properties of salt (NaCl) used for the preservation of natural casings. *Food Microbiology* 23, 657–62.
- Wilson M, Patel H, Fletcher J (1996) Susceptibility of biofilms of *Streptococcus sanguis* to chlorhexidine gluconate and cetylpyridinium chloride. *Oral Microbiology and Immunology* 11, 118–92.
- Wu MK, R'oris A, Barkis D, Wesselink PR (2000) Prevalence and extent of long oval canals in the apical third. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology 89, 739–43.
- Yoon SS, Hennigan RF, Hilliard GM et al. (2002) Pseudomonas aeruginosa anaerobic respiration in biofilms: relationships to cystic fibrosis pathogenesis. Developmental Cell 3, 593–603.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.