

## Attitudes and opinions of oral health and oral care among community-dwelling elderly subjects in Sweden: an interview study

K Andersson

Karlstad University, Karlstad, Sweden

**Objective:** The aim of this study was to investigate attitudes and opinions of old, independently living people about their oral health and how it has affected them through life. **Material and methods:** A random selection of 12 individuals was made in a group of 79 individuals who earlier participated in a study about self-reported oral health. The average age was 78.4 years. An interview guide was used as support to focus on oral health. The respondents themselves decided what was important within the topics. The respondents were encouraged to speak freely about their childhood, family, social relations, memories from early dental care, general health situation, experience from health care, life and their future. The interviews were recorded and extended from 60 to 90 min. All of them were transcribed and analysed with phenomenological method inspired by Giorgi. The analysis was made by two researchers independently. After 11 interviews, no new information was found; similar opinions, answers and stories recurred. **Results:** Seven of 11 respondents were born and grown up in the countryside and had moved to Stockholm in the forties. Awareness about dental care was generally low. Many of them have terrible memories from early dental visits, and the dentist was authoritarian and rough. Many had no memories of brushing their teeth as a child. Most of the respondents went to dentists as adults and when they could afford to pay by themselves. After retirement, all have continuous dental care and have visited a dental hygienist during the last 20 years. Nine of 11 have regular contact with a dental hygienist. Most of them are satisfied with their oral health and want to continue being clean and healthy in the mouth. **Conclusions:** Most of the respondents experienced an improved oral health, information and instructions from the dental hygienist have affected their self-care. Many tell that they have become more aware and carry out the oral hygiene more carefully after retirement. All of them experience that oral health affects the quality of life.

## Ingredients in dentifrices and their effect on plaque, gingivitis and mutans streptococci

L Jannesson<sup>1</sup>, D Birkhed<sup>2</sup>, S Renvert<sup>1</sup>

<sup>1</sup>Kristianstad University, Kristianstad, Sweden, <sup>2</sup>The Sahlgrenska Academy at Göteborg University, Göteborg, Sweden

The main objectives of this thesis were to study: (i) the effect of an enzyme-containing dentifrice (Zendium Dentine<sup>®</sup>), with addition of xylitol on mutans streptococci (MS) in saliva and dental plaque (Paper I) (ii) the effect of a combination of triclosan

and xylitol in a dentifrice (Colgate Total<sup>®</sup>) on MS in saliva and dental plaque (Paper II), and (iii) the effect of oxybenzone on prostaglandin E<sub>2</sub> (PGE<sub>2</sub>)-production in cell culture and the effect of an oxybenzone-containing dentifrice on plaque, gingivitis and MS (Paper III). In Paper I, the subjects were divided into two test groups: one using a 10% xylitol and the other using a 5% xylitol dentifrice for 3 months. The addition of 10% xylitol to Zendium Dentine<sup>®</sup> had an inhibitory effect on MS in both saliva and dental plaque, and the effect of xylitol seemed to be dose dependent. In Paper II, three groups were using one of the following dentifrices: (i) Colgate Total<sup>®</sup> with addition of 10% xylitol; (ii) Colgate Total<sup>®</sup>; and (iii) Colgate Total<sup>®</sup> without triclosan and without xylitol. The results showed that the addition of 10% xylitol to Colgate Total<sup>®</sup> reduced the number of MS in saliva and plaque. This effect was more pronounced at 6 months than at 2 months. In Paper III, the effect of oxybenzone was studied *in vitro* and *in vivo*. Human Embryo Palatal Mesenchyme (HEPM) cells were used to test the inhibition of IL-1 $\beta$ -stimulated PGE<sub>2</sub> production by different concentrations of oxybenzone. The results revealed that there was no decrease of cell viability up to 50  $\mu$ M. A dose-dependent inhibition of stimulated PGE<sub>2</sub> production was found: 50% inhibition (IC<sub>50</sub>) was found at 0.6  $\mu$ M. Paper III also included a double-blind clinical trial testing two fluoride dentifrices: one with the addition of 0.5% oxybenzone and one without. Plaque index was reduced in both groups. There was no difference between the groups. A 25% reduction in gingival index was observed in the oxybenzone group after 6 weeks, compared to 2% in the placebo group, indicating an anti-inflammatory effect of oxybenzone.

## Research utilisation among Swedish dental hygienists

K Öhrn, C Olsson

Dalarna University, Falun, Sweden

Dental hygienists have to practise evidence-based decision making in the future, which means that actively seeking and utilising research findings will become more important. **Objective:** The objective of this study was to explore barriers to and facilitators of research utilisation in clinical practice among Swedish dental hygienists. **Methods:** The study was a descriptive, comparative cross-sectional survey including a random sample of 491 dental hygienists in Sweden. The response rate was 62%. A validated questionnaire covering different aspects of participation in research, support for and availability of research, and research utilisation was employed. **Results:** The most common research-related activities were: reading research projects in professional journals, 83%; participating in clinical audit, 67%; and sharing research findings with their own professional colleagues, 65%.

Copyright of International Journal of Dental Hygiene is the property of Blackwell Publishing Limited and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.

Copyright of International Journal of Dental Hygiene is the property of Wiley-Blackwell and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.