



HK Yuen
FN Hant
C Hatfield
LM Summerlin
EA Smith
RM Silver

Authors' affiliations:

HK Yuen, Department of Occupational Therapy, School of Health Professions, University of Alabama at Birmingham, Birmingham, AL, USA
FN Hant, C Hatfield, EA Smith, RM Silver, Division of Rheumatology & Immunology, Department of Medicine, Medical University of South Carolina, Charleston, SC, USA
LM Summerlin, Department of Craniofacial Biology, College of Dental Medicine, Medical University of South Carolina, Charleston, SC, USA

Correspondence to:

Hon K. Yuen
Department of Occupational Therapy
School of Health Professions
University of Alabama at Birmingham
1720 2nd Avenue South
Birmingham
AL 35294
USA
Tel.: +1 205 934 6301
Fax: +1 205 975 7787
E-mail: yuen@uab.edu

Factors associated with oral hygiene practices among adults with systemic sclerosis

Abstract: *Objective:* To identify factors associated with oral hygiene practices in adults with systemic sclerosis (SSc). *Methods:* In this cross-sectional study, 178 dentate adults with SSc received an oral examination which included measurement of oral aperture, assessment of manual dexterity to perform oral hygiene, as well as completion of the Center of Epidemiological Studies Depression (CES-D) Scale and an oral health-related questionnaire. *Results:* Multivariable logistic regression modelling showed male, minority and high CES-D scores (i.e. clinically significant symptoms of depression) were associated with less likelihood of participants brushing teeth at least twice daily, but the presence of self-reported dry mouth symptoms increased the likelihood of toothbrushing. Having a dental visit in the past 12 months and use of an adapted flossing or interdental cleaning device were significantly associated with daily dental flossing; however, having difficulty flossing teeth reduced the likelihood of daily flossing. *Conclusions:* Overall, demographic variables were strongly associated with toothbrushing frequency, whereas flossing self-efficacy and barriers were strongly associated with dental flossing frequency in adults with SSc. The results suggest that dental health professionals should take mental health into consideration when educating patients with SSc to improve their oral hygiene and consider making referrals for patients exhibiting suspected clinically significant depressive symptoms to mental health professionals for further evaluation and treatment. In addition, an appropriate adapted flossing or interdental cleaning device should be recommended to increase dental flossing practices in this patient population.

Key words: adapted devices; epidemiologic methods; mental health; oral hygiene; scleroderma

Dates:

Accepted 9 September 2013

To cite this article:

Int J Dent Hygiene 12, 2014; 180–186.
DOI: 10.1111/ijdh.12056
Yuen HK, Hant FN, Hatfield C, Summerlin LM, Smith EA, Silver RM. Factors associated with oral hygiene practices among adults with systemic sclerosis.

© 2013 John Wiley & Sons A/S. Published by John Wiley & Sons Ltd

Introduction

Systemic sclerosis (SSc, scleroderma) is one of a group of autoimmune connective tissue diseases that is characterized by inflammation, vascular dysfunction and excessive fibrosis of connective tissue supporting the skin and visceral organs (1, 2). Systemic sclerosis affects the skin and musculoskeletal tissue of the hands as well as the oral and perioral tissues (3–6). Manual dexterity impairments and orofacial dysfunction (e.g. microstomia and xerostomia) are two major clinical manifestations that may complicate oral hygiene and lead to oral health problems among people with SSc (7–9). Hand deformity and limited oral aperture may

affect oral hygiene performance as well as willingness to perform oral hygiene (7, 10). Xerostomia (excessive dry mouth resulting from salivary hypofusion) has been shown to promote the development of dental plaque and increase the risk of developing oral diseases (11–13).

A population survey of dentate adults in Detroit showed dental flossing frequency was associated with flossing self-efficacy and benefits–barriers of flossing (14). Adapted oral hygiene devices to accommodate microstomia and decreased manual dexterity may help reduce oral self-care barriers in adults with SSc (15–18). Studies comparing the use of an adapted flosser to finger flossing indicated that participants with no manual dexterity impairment preferred the adapted flosser over finger flossing (19, 20). Several authors concluded that the adapted flosser was significantly more effective in facilitating patient compliance and establishing long-term regular flossing habits (17, 21). In regard to xerostomia, patients with Sjogren's syndrome were found to have similar or even better oral hygiene habits than the control subjects without xerostomia (22, 23).

In addition to various SSc disease-related specific factors, such as dry mouth, decreased oral aperture and manual dexterity impairment, secondary psychosocial conditions such as depressive symptoms may also have an impact on oral hygiene practices. High levels of depressive symptoms are common in adults with SSc (24, 25). Patients exhibiting depressive disorders often demonstrate disinterest in performing oral hygiene (26). A population survey in Northern Finland found persons with a high number of depressive symptoms had lower toothbrushing frequency than persons with no or only a few depressive symptoms (27). In addition, poor oral hygiene as indicated by high plaque score was significantly associated with higher levels of depressive symptoms (28).

Few studies have investigated the association of SSc disease-related factors and secondary psychosocial conditions with oral hygiene practices in adults with SSc (7, 29–31). Therefore, the purpose of the present study was to identify factors associated with oral hygiene practices in adults with SSc. Identification of these factors (e.g. SSc disease-related and secondary conditions) is important in designing appropriate preventive oral health education and promotion for this patient population. Findings can help the dental health professional tailor oral care educational programs and provide specific evidence that can be used to educate patients with SSc to improve oral hygiene practices.

Methods

Participants

Participants eligible for this cross-sectional study were adults (aged >18 years) diagnosed with SSc and who fulfilled the American College of Rheumatology preliminary classification criteria for SSc (32). Exclusion criteria were localized scleroderma (e.g. morphea, linear scleroderma and *en coup de sabre*).

Recruitment

Study participants were recruited through the Medical University of South Carolina (MUSC) scleroderma clinic and a local connective tissue disease database (CTDD). The CTDD contains medical information on the majority of patients with SSc who received consultation and/or treatment at the MUSC scleroderma clinic beginning in 2001. At the time when this study started (which was in October of 2007), the CTDD contained contact information on 509 patients with SSc. Potential subjects listed in the CTDD and patients who visited the MUSC scleroderma clinic were invited (either in person or by phone) to participate in the dental survey study. The research coordinator, on behalf of the physicians at the MUSC scleroderma clinic, contacted the patients by phone and reminded them of their upcoming physician appointment and invited them to participate in the study. In addition, during the week (every Monday), the research coordinator was assigned to the scleroderma clinic for recruitment. The research coordinator explained the study to potential participants including their time commitment, obtaining verbal consent and scheduling an oral examination appointment. The study required a single dental visit of about 1.5-h duration.

Procedures

Informed consent was obtained, and the oral examination was conducted at the MUSC General Clinical Research Center. The examination included measurement of the oral aperture, assessment of manual dexterity to perform oral hygiene, and dental and periodontal health, as well as completion of the Center of Epidemiological Studies Depression (CES-D) Scale (33), a self-report instrument to assess depressive symptomatology and a package containing an oral health-related questionnaire. Two trained and calibrated dental hygienists were designated as the oral health examiners and conducted the mouth cavity assessments (34). The protocol was approved by the MUSC Institutional Review Board. The clinical trial protocol number was NCT01817361.

Outcome variables

Frequency of toothbrushing and frequency of dental flossing were the two outcome measures. Frequency of toothbrushing was defined as toothbrushing at least twice daily versus less often (coded as 1 or 0 respectively). Frequency of dental flossing was defined as flossing at least once a day versus less often (coded as 1 or 0, respectively).

Explanatory variables

Manual dexterity to perform oral hygiene was assessed by the Toothbrushing Ability Test (TAT) (35, 36). Detailed procedures and scoring have been described in the literature (35, 36). Briefly, participants were requested to brush the front surfaces of all lower front teeth (i.e. from canine to canine) in

30 s using a regular manual toothbrush. After being stained, three anterior mandibular teeth (i.e. #22, #25, and #27) were scored according to patient hygiene performance (PHP) criteria (37). The score of the TAT ranges from 0 to 19 with a score >6 indicating manual dexterity impairments.

Maximum oral aperture was measured, using a small metal ruler, as the distance in mm between the upper and lower incisal edges of the right central incisors when the participants were requested to open their mouth as wide as possible (38). If the right central incisors were absent, the left central, right lateral or left lateral incisors were substituted in that order (38). Three successive trials of maximum oral aperture measurement, with a 5-s pause in between each measurement, were conducted and recorded (39). An average of the three successive trials for each oral aperture measurement was computed.

The Center for Epidemiologic Studies Depression (CES-D) Scale is a self-report instrument to assess the presence and severity of depressive symptomatology in the general population (33). The CES-D consists of 20 items (symptoms) grouped into four multiitem subscales that assess general depression, positive affect, somatic/vegetative symptoms and interpersonal difficulties. The frequency of occurrence of each symptom during the past week is rated on a four-point Likert-type scale ranging from 'rarely' (0) to 'mostly' (3). After reverse coding of four items, scores from 20 items are summed to form a composite score with a possible range of 0–60. A higher score reflects a higher level of depression. The CES-D has been demonstrated to be a reliable and valid measure of depressive symptoms in patients with SSc (40). The CES-D score of ≥ 19 has shown to be the most accurate cut-off for identifying patients with rheumatoid arthritis who exhibit clinically significant symptoms of depression (41).

The oral health-related questionnaire contained items addressing the demographic and socio-economic characteristics of the participants, as well as their dental care visit patterns, oral hygiene behaviours, smoking habits and dry mouth symptoms. Demographic and socio-economic characteristics included age, gender, race [minority (i.e. African American and others) versus Caucasian American], marital status (married/live together versus unmarried), education (less than college versus college and beyond), employment status (paid employment versus not employed), annual household income (ranging from below \$10 000 to \$75 000 and above, with \$5000 incremental brackets) and dental insurance (private/public-funded versus none). Dental visits in the past year were recoded as 'yes/1' or 'no/0'. Mouth rinsing practices were indicated by using mouth rinse at least once a day versus less often. Other questions related to oral hygiene included use of adapted oral hygiene devices (1 = use or 0 = not use), having difficulty cutting a piece of dental floss from a floss box (1 = have difficulty, or 0 = no difficulty) and having difficulty flossing teeth (1 = have difficulty, or 0 = no difficulty). Participants' smoking was indicated by current smokers versus past smokers and those who never smoked.

The presence of dry mouth symptoms (coded as '1') was indicated by a positive response to one or more of the following four questions (42): (i) 'Do you sip liquids to aid in swallowing dry foods?'; (ii) 'Does your mouth feel dry when eating a meal?'; (iii) 'Do you have difficulties swallowing any foods?'; and (iv) 'Does the amount of saliva in your mouth seem to be too little?'

Data analysis

Prior to developing multivariable logistic regression models for predicting the probability of patients with SSc who complete daily oral hygiene practices (i.e. toothbrushing and dental flossing), bivariate analyses (using univariate regression models) were performed to determine the relationship between the explanatory variables and daily oral hygiene practices. Explanatory variables with a P -value < 0.2 in bivariate models were considered to be candidates for multivariable modelling (43). Certain demographic and socio-economic status variables such as gender, marital status, education and income have been shown to be positively associated with oral hygiene practices (14, 44), and we controlled for the confounding effect of these variables in our multivariable analysis regardless of its P -value in the univariate analysis. Age and SSc disease onset were highly correlated ($r = 0.65$). To avoid problems caused by collinearity, we selected age as the more appropriate variable to be included in the multivariable analysis. All P -values reported in the multivariable models were two-sided, and the type I error rate was set at 0.05. All statistical analyses were performed using IBM SPSS Statistics for Windows, version 20 (<http://www.spss.com>).

Results

We contacted all 509 patients in the CTDD and invited them to participate. Of these patients, 72 patients' phones were disconnected, 20 were deceased, 160 declined, 20 were not diagnosed with SSc and 47 gave verbal consent but did not show up to their appointment. In summary, 190 patients with SSc completed the oral examination with a recruitment rate of 37.3%.

Only dentate participants (i.e. those with at least one tooth) were included in the study. Eleven participants were edentulous and excluded for analysis. In addition, one African American male participant did not complete the oral health-related questionnaire. Therefore, the analytic sample consisted of 178 participants with usable data for this study.

Tables 1 and 2 show the characteristics of the 178 dentate participants, which includes their diagnosis, comorbidities, socio-demographics and oral health-related behaviour information. The majority of participants were Caucasian (66.3%), and 32.0% of them were African American; three participants (1.7%) identified themselves as Asian or Hispanic/Latino. The median disease duration and its interquartile range were 4.8 and 3–12 years, respectively. The mean and standard deviation (SD) number of participants' natural teeth (excluding third molars) was 23.4 (6.2) with a range from 2 to 28. The median number of missing teeth was two. The majority of the

Table 1. Characteristics of the participants (n = 178)

Characteristics	Mean \pm SD (range) or n (%)
Diagnosis and comorbidities	
Limited cutaneous subtype of SSc	106 (59.6)
Diffused cutaneous subtype of SSc	72 (40.4)
Disease duration (years)*	8.4 \pm 8.4 (0.5–40)
Oral aperture (mm) [†]	37.8 \pm 8.5 (10–60)
TAT score [‡]	9.2 \pm 2.7 (0–16)
CDS-D score	12.1 \pm 10.4 (0–52)
Self-report dry mouth symptoms [‡]	123 (71.5)
Socio-demographics	
Age (years)	54.5 \pm 13.0 (21.9–82.7)
Female	149 (83.7)
Caucasian American	118 (65.9)
African American	58 (32.4)
Married/live together	114 (64.0)
Education (less than college)	91 (51.1)
Employment (FT and PT)	63 (35.4)
Annual income (<\$20 000) [†]	39 (22.7)
Annual income (\geq \$55 000) [†]	73 (42.4)
No dental insurance	61 (34.3)

CDS-D, Center of Epidemiological Studies Depression; FT, full time, PT, part time; SD, standard deviation; SSc, systemic sclerosis; TAT, Toothbrushing Ability Test.

*Less than 9% of the data were missing.

[†]Less than 4% of the data were missing.

[‡]16% of the data were missing, as participants who required antibiotic therapy prior to dental examination did not engage in the TAT.

Table 2. Oral health-related behaviours of the participants (n = 178)

Variables	n (%)
Dental visit (in past 12 months)	138 (77.5)
Toothbrushing (at least twice a day)	125 (70.2)
Dental flossing (at least once a day)	72 (40.4)
Use mouth rinse (at least once a day)	66 (37.1)
Use an electric toothbrush*	64 (36.4)
Use an oral irrigator*	12 (7.0)
Use adapted flossing device*	82 (46.9)
Have difficulty flossing teeth*	95 (54.6)
Have difficulty cutting a piece of dental floss from a floss box*	22 (12.7)
Non-smoker (current)	165 (92.7)

*Less than 4% of the data were missing.

participants (85.3%) demonstrated impairment in manual dexterity to perform oral hygiene as indicated by TAT scores above six; about three quarters (71.5%) reported having dry mouth symptoms; and one quarter (24.7%) were classified as having clinically significant symptoms of depression as indicated by their CES-D score above 18. The mean (SD) of participants' size of oral aperture was 37.8 (8.5) mm with about 16% of the participants having an oral aperture of <30 mm.

Factors affecting toothbrushing frequency among adults with SSc

Results of the bivariate analyses indicated that toothbrushing at least twice daily was shown to be significantly related to the

following variables: race, gender, income, had a dental visit in the past 12 months, daily dental flossing, CES-D score and dry mouth (all P -values < 0.05). Table 3 shows the results of the bivariate analyses and variables that were included in the multivariable logistic regression model. After adjusting for other explanatory variables, only race, gender, CES-D score and dry mouth were included in the final multivariable model for toothbrushing at least twice daily (see Table 3). Compared to the minority group (95% of them were African Americans), Caucasians had a 2.5 times greater odds of brushing teeth at least twice daily (adjusted OR = 2.52, 95% CI = 1.18–5.37, P < 0.05). Females with SSc had a 4.4 times greater odds of brushing teeth at least twice daily compared to their male counterpart (adjusted OR = 4.41, 95% CI = 1.70–11.40, P < 0.005). Dentate adults with SSc who reported dry mouth symptoms had a 2.8 times greater odds of brushing teeth at least twice daily compared to those who did not complain of dry mouth (adjusted OR = 2.77, 95% CI 1.22–6.29, P < 0.05). Dentate adults with SSc who reported having clinically significant symptoms of depression (i.e. CES-D \geq 19) had 71% less odds to perform toothbrushing at least twice daily than those who reported less severity of depressive symptoms (adjusted OR = 0.29, 95% CI = 0.12–0.67, P < 0.005).

Factors affecting dental flossing frequency among adults with SSc

Results of the bivariate analyses indicated that dental flossing at least once daily was shown to be significantly related to the following variables: age, race, had a dental visit in the past 12 months, daily toothbrushing and use of an adapted flossing device (all P -values < 0.05). Table 4 shows the results of the bivariate analyses and variables that were included in the multivariable logistic regression model. After adjusting for other explanatory variables, had a dental visit in the past 12 months, use an adapted flossing or interdental cleaning device and having difficulty flossing teeth were included in the final multivariable model for daily flossing (see Table 4). Participants who had a dental visit in the past 12 months had a 3.9 times greater odds of flossing teeth daily compared to those who did not have a dental visit in the past 12 months (adjusted OR = 3.88, 95% CI = 1.47–10.26, P < 0.01). Participants who used an adapted flossing or interdental cleaning device had a 2.5 times greater odds of flossing teeth daily compared to those who did not use an adapted or interdental cleaning device (adjusted OR = 2.50, 95% CI = 1.27–4.92, P < 0.01). Participants who reported having difficulty flossing teeth had 55% less odds to floss daily compared to those who did not report having difficulty flossing (adjusted OR = 0.45, 95% CI 0.23–0.89, P < 0.05).

Discussion

As indicated in the results, factors that influence toothbrushing frequency differed from flossing frequency in dentate adults with SSc. For toothbrushing, race, gender, self-perceived

Table 3. Univariate and multivariable analyses examining factors associated with toothbrushing of at least twice daily in dentate adults with systemic sclerosis

Explanatory variables	OR (95% CI)	P-value	Adj OR (95% CI)	P-value
Age (years)	1.02 (0.99, 1.05)	0.072		
Race (Caucasian)	3.26 (1.66, 6.38)	0.001	2.52 (1.18, 5.37)	0.017
Gender (female)	3.73 (1.64, 8.47)	0.002	4.41 (1.70, 11.40)	0.002
Marital status (married)	1.40 (0.72, 2.72)	0.316		
Education (college)	1.71 (0.89, 3.28)	0.109		
Income	1.07 (1.00, 1.15)	0.036		
Had a dental visit in the past 12 months	3.18 (1.53, 6.62)	0.002		
Flossing daily	2.14 (1.19, 4.88)	0.014		
Oral aperture	0.97 (0.93, 0.1.01)	0.151		
Dry mouth	2.23 (1.11, 4.47)	0.024	2.77 (1.22, 6.29)	0.015
Depression (CES-D \geq 19)	0.34 (0.17, 0.70)	0.003	0.29 (0.12, 0.67)	0.004

Adj OR, adjusted odds ratio; CES-D, Center of Epidemiological Studies Depression; CI, confidence interval; OR, odds ratio.

Table 4. Univariate and multivariable analyses examining factors associated with dental flossing of at least once a day in dentate adults with systemic sclerosis

Explanatory variables	OR (95% CI)	P-value	Adj OR (95% CI)	P-value
Age (years)	1.04 (1.01, 1.06)	0.006		
Race (Caucasian)	2.48 (1.26, 4.89)	0.008		
Gender (female)	1.35 (0.59, 3.11)	0.475		
Marital status (married)	1.66 (0.88, 3.14)	0.121		
Education (college)	1.08 (0.59, 1.96)	0.805		
Income	1.05 (0.99, 1.12)	0.102		
Had a dental visit in the past 12 months	5.19 (2.05, 13.17)	0.001	3.88 (1.47, 10.26)	0.006
Number of teeth	1.04 (0.99, 1.10)	0.123		
Toothbrushing frequency	2.41 (1.19, 4.88)	0.014		
Use adapted flossing device	2.21 (1.19, 4.09)	0.012	2.50 (1.27, 4.92)	0.008
Have difficulty flossing teeth	0.55 (0.30, 1.01)	0.055	0.45 (0.23, 0.89)	0.022
Have difficulty cutting a piece of dental floss from a floss box	0.51 (0.19, 1.38)	0.183		

Adj OR, adjusted odds ratio; CI, confidence interval; OR, odds ratio.

mouth dryness and depressive symptoms are the four explanatory variables that were significantly associated with toothbrushing at least twice daily, whereas had a dental visit in the past 12 months, use of adapted flossing or interdental cleaning devices and have difficulty flossing teeth are the three explanatory variables that were significantly associated with daily dental flossing.

Consistent with the literature on predicting oral hygiene habits in the general population (14), demographic variables were strongly associated with toothbrushing frequency, whereas flossing self-efficacy and barriers were strongly associated with dental flossing frequency in adults with SSc. For example, participants who were minority (African American) and male (14, 44) were less likely to brush teeth at least twice daily. Similar to patients with Sjogren's syndrome (22), participants who reported mouth dryness had a higher toothbrushing frequency than those who reported no mouth dryness. The clinical implications of the association between higher scores on the CES-D (i.e. having clinically significant symptoms of depression) and less likely to brush teeth at least twice daily are that significant neglect of oral hygiene in patients with SSc may alert dental healthcare professionals to be aware of potential underlying depressive symptoms (including diagnosis and

antidepressants used by patients with SSc), and to consider making timely referrals to mental health professionals for evaluation and treatment.

The dexterity to perform sufficient oral hygiene has been previously suggested as affecting oral health in people with SSc (7, 9). In this study, an objective measure of manual dexterity impairment, as indicated by the TAT score, and size of oral aperture were not significantly associated with frequency of oral hygiene practices. As the disease progressed, participants' hand function and size of oral aperture may deteriorate. Participants may maintain their daily frequency of toothbrushing routine, although the quality of plaque removal may be scarified. The majority of patients with SSc develop this disease in middle age and beyond, and toothbrushing habits have already been ingrained in their daily routine.

Having difficulty flossing teeth was significantly associated with being less likely to floss at least daily. In the present study, only 40% of the study participants flossed at least once daily, which is at the same rate as that of a clinical population (45). As patients with SSc are at a greater risk of dental disease (46, 47) and dental flossing, especially in the evening, was shown to be significantly related to less gingival inflammation in this patient population (9), it is important for dental health

professionals to educate patients with SSc on the importance of daily flossing. The strategies that may aid those who floss less than daily include the use of adapted flossing or interdental cleaning device to decrease the difficulty in flossing and increase daily flossing adherence.

The electric toothbrush can compensate for decreased manual dexterity and improve plaque control and reduce gingival inflammation (16), but the size of the brush head of most oscillating-rotating electric toothbrushes is either the same or bigger than that of a manual toothbrush. Dental health professionals should emphasize the importance of at least twice daily toothbrushing by recommending the use of an electric toothbrush with a child-sized brush head.

We acknowledge that the sample selection in this study was limited to one institution and the findings may or may not be generalizable to the whole SSc population; however, the participants' demographic characteristics (gender, type of SSc, incidence of limited and diffuse forms of SSc among African and Caucasian Americans, and mean age of SSc disease onset) of this large oral health study on SSc are similar to those reported in several population-based epidemiological studies conducted in the US (48–51). For example, about 80% of the participants were women (50, 51), the ratio of limited to diffuse form of SSc was 59.6% to 40.4%, with more African American participants diagnosed with the diffuse form (56.1%) and more Caucasian American participants diagnosed with the limited form (67.8%) (49, 51). The mean age of SSc disease onset among African American participants was about 10 years younger than that of Caucasian American participants (38.3 versus 50.2 years of age) (48). Finally, the incidence of having clinically significant symptoms of depression (i.e. CES-D \geq 19) in the present study (25%) was very similar to that reported in previous studies (26–29%) (52, 53).

Conclusion

Little was known about factors influencing oral hygiene practices in adults with SSc. Identification of these factors will provide evidence for clinicians to tailor patient education. Findings from this study revealed that clinically significant depressive symptoms reduced the likelihood of toothbrushing. The use of adapted interdental cleaning devices was associated with flossing frequency, whereas having difficulty flossing teeth reduced the likelihood of flossing. It is concluded that significant neglect of oral hygiene on patients with SSc may alert clinicians to be aware of potential underlying depressive symptoms and to consider making referrals to mental health professionals for evaluation and treatment. Recommendations on the use of adapted interdental cleaning device may increase flossing adherence.

Acknowledgements

The authors wish to thank the dental hygienist, Pemra L. Hudson, RDH for conducting the oral examination; and the project coordinator, Samantha Mahoney, for subject recruitment

and data entry. This study was supported by a grant awarded to the first author from the National Institute of Dental and Craniofacial Research (Grant # 1R21DE017360-01A2). Additional support was provided by the Multidisciplinary Clinical Research Center grant from the National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health (Grant # P60 AR049459), the South Carolina Centers of Biomedical Research Excellence (COBRE) for Oral Health with funding provided by the National Institutes of Health (NIH) and the National Center for Research Resources (NCRR) with a P20 RR-017696, and the South Carolina Clinical & Translational Research Institute, Medical University of South Carolina's CTSA, NIH/NCRR Grant Number UL1RR029882.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- 1 Derk CT, Jimenez SA. Systemic sclerosis: current views of its pathogenesis. *Autoimmun Rev* 2003; **2**: 181–191.
- 2 Dghoughi S, El Wady W, Taleb B. Systemic sclerosis. Case report and review of literature. *NY State Dent J* 2010; **76**: 30–35.
- 3 Alantar A, Cabane J, Hachulla E *et al*. Recommendations for the care of oral involvement in patients with systemic sclerosis. *Arthritis Care Res (Hoboken)* 2011; **63**: 1126–1133.
- 4 Albilia JB, Lam DK, Blanas N, Clokie CM, Sandor GK. Small mouths. Big problems? A review of scleroderma and its oral health implications. *J Can Dent Assoc* 2007; **73**: 831–836.
- 5 Fischer DJ, Patton LL. Scleroderma: oral manifestations and treatment challenges. *Spec Care Dentist* 2000; **20**: 240–244.
- 6 Skare TL, Toebe BL, Boros C. Hand dysfunction in scleroderma patients. *Sao Paulo Med J* 2011; **129**: 357–360.
- 7 Poole JL, Brewer C, Rossie K, Good CC, Conte C, Steen V. Factors related to oral hygiene in persons with scleroderma. *Int J Dent Hyg* 2005; **3**: 13–17.
- 8 Scardina GA, Messina P. Systemic sclerosis: description and diagnostic role of the oral phenomena. *Gen Dent* 2004; **52**: 42–47.
- 9 Yuen H, Weng Y, Reed S, Summerlin L, Silver R. Factors associated with gingival inflammation among adults with systemic sclerosis. *Int J Dent Hyg* 2013; doi: 10.1111/idh.12024 [Epub ahead of print].
- 10 Kelsey JL, Lamster IB. Influence of musculoskeletal conditions on oral health among older adults. *Am J Public Health* 2008; **98**: 1177–1183.
- 11 Guggenheimer J, Moore PA. Xerostomia: etiology, recognition and treatment. *J Am Dent Assoc* 2003; **134**: 61–69; quiz 118–9.
- 12 Hase JC, Birkhed D. Salivary glucose clearance, dry mouth and pH changes in dental plaque in man. *Arch Oral Biol* 1988; **33**: 875–880.
- 13 Sreebny LM, Valdini A. Xerostomia. A neglected symptom. *Arch Intern Med* 1987; **147**: 1333–1337.
- 14 Ronis DL, Antonakos CL, Lang WP. Usefulness of multiple equations for predicting preventive oral health behaviors. *Health Educ Q* 1996; **23**: 512–527.
- 15 Iacono VJ, Aldredge WA, Lucks H, Schwartzstein S. Modern supragingival plaque control. *Int Dent J* 1998; **48**(3 Suppl 1): 290–297.
- 16 Yuen HK, Weng Y, Bandyopadhyay D, Reed SG, Leite RS, Silver RM. Effect of a multi-faceted intervention on gingival health

- among adults with systemic sclerosis. *Clin Exp Rheumatol* 2011; **29** (2 Suppl 65): S26–S32.
- 17 Brothwell DJ, Jutai DK, Hawkins RJ. An update of mechanical oral hygiene practices: evidence-based recommendations for disease prevention. *J Can Dent Assoc* 1998; **64**: 295–306.
- 18 Choo A, Delac DM, Messer LB. Oral hygiene measures and promotion: review and considerations. *Aust Dent J* 2001; **46**: 166–173.
- 19 Carter-Hanson C, Gadbury-Amyot C, Killoy W. Comparison of the plaque removal efficacy of a new flossing aid (Quik Floss) to finger flossing. *J Clin Periodontol* 1996; **23**: 873–878.
- 20 Spolsky VW, Perry DA, Meng Z, Kissel P. Evaluating the efficacy of a new flossing aid. *J Clin Periodontol* 1993; **20**: 490–497.
- 21 Kleber CJ, Putt MS. Formation of flossing habit using a floss-holding device. *J Dent Hyg* 1990; **64**: 140–143.
- 22 Boutsis EA, Paikos S, Dafni UG, Moutsopoulos HM, Skopouli FN. Dental and periodontal status of Sjogren's syndrome. *J Clin Periodontol* 2000; **27**: 231–235.
- 23 Leung KC, McMillan AS, Leung WK, Wong MC, Lau CS, Mok TM. Oral health condition and saliva flow in southern Chinese with Sjogren's syndrome. *Int Dent J* 2004; **54**: 159–165.
- 24 Kwakkenbos L, van Lankveld WG, Vonk MC, Becker ES, van den Hoogen FH, van den Ende CH. Disease-related and psychosocial factors associated with depressive symptoms in patients with systemic sclerosis, including fear of progression and appearance self-esteem. *J Psychosom Res* 2012; **72**: 199–204.
- 25 Thombs BD, Hudson M, Taillefer SS, Baron M, Canadian Scleroderma Research Group. Prevalence and clinical correlates of symptoms of depression in patients with systemic sclerosis. *Arthritis Rheum* 2008; **59**: 504–509.
- 26 Friedlander AH, Mahler ME. Major depressive disorder. Psychopathology, medical management and dental implications. *J Am Dent Assoc* 2001; **132**: 629–638.
- 27 Anttila S, Knuuttila M, Ylostalo P, Joukamaa M. Symptoms of depression and anxiety in relation to dental health behavior and self-perceived dental treatment need. *Eur J Oral Sci* 2006; **114**: 109–114.
- 28 Kurer JR, Watts TL, Weinman J, Gower DB. Psychological mood of regular dental attenders in relation to oral hygiene behaviour and gingival health. *J Clin Periodontol* 1995; **22**: 52–55.
- 29 Chu CH, Yeung CM, Lai IA, Leung WK, Mok MY. Oral health of Chinese people with systemic sclerosis. *Clin Oral Invest* 2011; **15**: 931–939.
- 30 Leader DM. Scleroderma and dentistry: every dentist is a scleroderma specialist. *J Mass Dent Soc* 2007; **56**: 16–19.
- 31 Tolle SL. Scleroderma: considerations for dental hygienists. *Int J Dent Hyg* 2008; **6**: 77–83.
- 32 Masi AT. Preliminary criteria for the classification of systemic sclerosis (scleroderma). Subcommittee for scleroderma criteria of the American Rheumatism Association Diagnostic and Therapeutic Criteria Committee. *Arthritis Rheum* 1980; **23**: 581–590.
- 33 Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas* 1977; **1**: 385–401.
- 34 Hill EG, Slate EH, Wiegand RE, Grossi SG, Salinas CF. Study design for calibration of clinical examiners measuring periodontal parameters. *J Periodontol* 2006; **77**: 1129–1141.
- 35 Felder R, James K, Brown C, Lemon S, Reveal M. Dexterity testing as a predictor of oral care ability. *J Am Geriatr Soc* 1994; **42**: 1081–1086.
- 36 Felder R, Reveal M, Lemon S, Brown C. Testing toothbrushing ability of elderly patients. *Spec Care Dentist* 1994; **14**: 153–157.
- 37 Podshadley AG, Haley JV. A method for evaluating oral hygiene performance. *Public Health Rep* 1968; **83**: 259–264.
- 38 Naylor WP, Douglass CW, Mix E. The nonsurgical treatment of microstomia in scleroderma: a pilot study. *Oral Surg Oral Med Oral Pathol* 1984; **57**: 508–511.
- 39 Wood GD, Branco JA. A comparison of three methods of measuring maximal opening of the mouth. *J Oral Surg* 1979; **37**: 175–177.
- 40 Thombs BD, Hudson M, Schieir O, Taillefer SS, Baron M, Canadian Scleroderma Research Group. Reliability and validity of the center for epidemiologic studies depression scale in patients with systemic sclerosis. *Arthritis Rheum* 2008; **59**: 438–443.
- 41 Martens MP, Parker JC, Smarr KL, Hewett JE, Slaughter JR, Walker SE. Assessment of depression in rheumatoid arthritis: a modified version of the center for epidemiologic studies depression scale. *Arthritis Rheum* 2003; **49**: 549–555.
- 42 Fox PC, Busch KA, Baum BJ. Subjective reports of xerostomia and objective measures of salivary gland performance. *J Am Dent Assoc* 1987; **115**: 581–584.
- 43 Mickey J, Greenland S. A study of the impact of confounder-selection criteria on effect estimation. *Am J Epidemiol* 1989; **129**: 125–137.
- 44 Ronis DL, Lang WP, Farghaly MM, Passow E. Tooth brushing, flossing, and preventive dental visits by Detroit-area residents in relation to demographic and socioeconomic factors. *J Public Health Dent* 1993; **53**: 138–145.
- 45 Stoltzberg JL, Osborn JB, Pihlstrom BL et al. Prevalence of periodontal disease in a health maintenance organization and comparisons to the national survey of oral health. *J Periodontol* 1993; **64**: 853–858.
- 46 Leung WK, Chu CH, Mok MY, Yeung KW, Ng SK. Periodontal status of adults with systemic sclerosis: case-control study. *J Periodontol* 2011; **82**: 1140–1145.
- 47 Wood RE, Lee P. Analysis of the oral manifestations of systemic sclerosis (scleroderma). *Oral Surg Oral Med Oral Pathol* 1988; **65**: 172–178.
- 48 Laing TJ, Gillespie BW, Toth MB et al. Racial differences in scleroderma among women in Michigan. *Arthritis Rheum* 1997; **40**: 734–742.
- 49 Reveille JD, Fischbach M, McNearney T et al. Systemic sclerosis in 3 US ethnic groups: a comparison of clinical, sociodemographic, serologic, and immunogenetic determinants. *Semin Arthritis Rheum* 2001; **30**: 332–346.
- 50 Steen VD, Oddis CV, Conte CG, Janoski J, Casterline GZ, Medsger TA Jr. Incidence of systemic sclerosis in Allegheny County, Pennsylvania. A twenty-year study of hospital-diagnosed cases, 1963–1982. *Arthritis Rheum* 1997; **40**: 441–445.
- 51 Mayes MD. Scleroderma epidemiology. *Rheum Dis Clin North Am* 2003; **29**: 239–254.
- 52 Kwakkenbos L, Arthurs E, van den Hoogen FH et al. Cross-language measurement equivalence of the Center for Epidemiologic Studies Depression (CES-D) scale in systemic sclerosis: a comparison of Canadian and Dutch patients. *PLoS ONE* 2013; **8**: e53923.
- 53 Nietert PJ, Mitchell HC, Bolster MB, Curran MY, Tilley BC, Silver RM. Correlates of depression, including overall and gastrointestinal functional status, among patients with systemic sclerosis. *J Rheumatol* 2005; **32**: 51–57.

Copyright of International Journal of Dental Hygiene is the property of Wiley-Blackwell and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.