# Dental caries experience and use of dental services among preschool children in Ajman, UAE

R. HASHIM<sup>1</sup>, W. M. THOMSON<sup>2</sup>, K. M. S. AYERS<sup>2</sup>, J. D. LEWSEY<sup>3</sup> & M. AWAD<sup>4</sup>

<sup>1</sup>Ajman University, Ajman, UAE, <sup>2</sup>University of Otago, Dunedin, New Zealand, <sup>3</sup>University of London, London, UK and <sup>4</sup>University of Sharjah, Sharjah, UAE

**Summary.** *Objective*. The aim of this study was to estimate the prevalence and severity of dental caries in the primary dentition of young children in Ajman, UAE, and investigate its association with sociodemographic characteristics and use of dental services.

*Methods.* A cluster-sampling approach was used to randomly select children aged 5 or 6 years who were enrolled in public or private schools. Clinical examinations for caries were conducted by a single examiner using World Health Organization criteria. Parents completed questionnaires seeking information on socioeconomic background and dental service utilization. Zero-inflated negative binomial (ZINB) regression modelling was used to identify risk markers and risk indicators for caries experience. *Results.* The prevalence of dental caries in the sample was high 76.1%. The average dmfs score 10.2. Caries severity was greater among older children and among male children of less educated mothers. Emirati (local) children had higher caries severity than others. Children who had higher level of caries visited the dentist more frequently than other children whose visits were for check-up only.

*Conclusions.* Dental caries prevalence and severity in young children in Ajman are high, and socioeconomic characteristics and dental utilization are important determinants of their dental caries experience. There is an urgent need for oral health programmes targeted at the treatment and underlying causes of dental caries in these children.

# Introduction

The United Arab Emirates (UAE) lies on the eastern side of Arabia. It is bordered by the Arabian Gulf, the Gulf of Oman, and Sultanate of Oman, Qatar and Saudi Arabia. In 1995, the population of the UAE was officially estimated at 2.378 million residents [1]. The Emirati population represents some 25-30% of this number, with the remainder being expatriates who are living in the UAE for variable lengths of time. Children between 0 and 14 years (33.9%) and women in the childbearing age group of 15-45 years (20.6%) represent 54.5% of the total population, which emphasizes the importance of maternal and child health services. In general, dental services are spread throughout the UAE. The dental care system is well-developed and comprehensive, and has a predominantly curative emphasis. About half a million people sought dental care in 1995 [2]. On average, there is one dentist for every 13 000 people.

Correspondence: R. Hashim, Ajman University, PO Box 346, Ajman, UAE. E-mail: raghad69@yahoo.co.nz

Local dentists have all qualified abroad, since there was no dental school in the Emirates until very recently. A private dental school has recently been established by the University of Ajman. The total number of dentists registered with the Ministry of Health in the Emirate of Ajman is 10 (nine general practitioners and one prosthodontist). There is only one dental centre available in Ajman.

The Emirate of Ajman was selected for this study because there was no information available on the oral health of preschool children. Only three studies of child oral health had previously been conducted in the UAE: Al-Mughery *et al.* [3] reported from the Emirate of Abu Dhabi; Al-Hosani and Rugg-Gunn [4] reported from Abu Dhabi, Al-Ain and the Western Region; and Naqvi *et al.* [5] focused on the Al-Ain region only. Those three studies all reported caries estimates which were relatively high by international standards. The objectives of this study were to determine the prevalence and severity of dental caries among preschool children in Ajman Emirate, and to investigate the association of disease with sociodemographic characteristics and dental service utilization.

## Subjects and methods

Ethical approval for the study was obtained from both the Ministry of Health in the UAE and the ethics committee of Otago University, Dunedin, New Zealand. There are a total of 22 urban and rural schools (kindergartens) in the Emirate of Ajman. A complex sampling scheme was used, with the school (kindergarten) selected as the primary sampling unit. Half of those schools were selected randomly from updated lists obtained from the Ministry of Education, using a computer program to generate random numbers. All of the students in each sampled school were selected. A questionnaire was used to elicit information on ethnicity, level of parental educational attainment and parental income. Information was also collected on dietary habits, children's exposure to fluoride, and their utilization of dental services. In addition to items capturing sociodemographic data, another 18 sought information on the children's diet, oral hygiene, oral self-care and the use of dental services. For the child's dietary habits, information was sought on usual eating frequency, and on the number of times the eight most common UAE children's snack foods were consumed. The oral hygiene information collected included the frequency of brushing, the type of toothpaste used, and whether the child was helped with brushing. Fluoride use was also assessed (e.g. 'Has your child ever taken fluoride tablets or drops?'), as well as whether the child had received any fluoride from other sources. Parents were asked about child dental service use (e.g. 'Has your child visited the dentist during the past 12 months?'). The questionnaires were pre-tested before use in the field in order to examine the extent to which parents could easily understand their content.

At the field data collection stage, the questionnaire was sent to the parents, and written consent was obtained before each child was dentally examined. Dental examinations were carried out using a disposable mouth mirror. Children were examined at the school health clinic while sitting on an ordinary chair. Natural daylight was used for illumination, and no radiographs were taken. Children with food remnants on their teeth were asked to rinse with water before their examination. The World Health Organization (WHO) criteria [6] were used, and caries was diagnosed at the cavitation stage.

To determine the reliability of dental caries examination, the recorder was asked to arrange for one in 10 children to be re-examined. This was done without the examiner's knowledge, and the examination and re-examination were separated by at least one day. Re-examination of approximately 10% of the children resulted in 97 children being seen twice by the examiner. Test–retest reliability for decayed, filled or missed surface (dmfs) scores was very high, with an intraclass correlation coefficient (ICC) of 0·99 (indicating an error variance of 1%). In addition, the main researcher (R.H.) and the supervisor (M.A.) both examined 10 children; the ICC for dmfs was 1·00. The ICC was calculated using the reliability analysis command in the SPSS computer program (SPSS for Windows, Version 11·0, SPSS Inc., Chicago, IL, USA).

At the analysis stage, each child was categorized into one of three groups according to maternal education: 'low' (not able to read or write, and had attended primary school only); 'middle' (had attended high-school-level education); and 'high' (had obtained a university or college education). Children were also categorized according to monthly household income, using the three groups of: low (< 3000 Dirhams; Dhs 3.6 = US\$1.00); middle (3001–7000 Dirhams); and high (> 7000 Dirhams).

The information from the questionnaires and caries forms was coded and entered into a spreadsheet for analysis using SPSS and Stata (Intercooled Stata 8.0, Stata Corporation, College Station, TX, USA). The level of statistical significance was set at P < 0.05. Because taking the complex sampling scheme [7] into account by using the 'survey' procedures in Stata (and using the school as the strata variable) made no difference to the estimates, the decision was made to treat the sample as a simple random sample. Data analysis was undertaken, first, to determine dental caries prevalence (represented by the percentage of children who had one or more dmfs) and severity (represented by the mean dmfs), and secondly, to investigate possible risk markers and indicators for caries prevalence and severity. After computing descriptive statistics, bivariate analyses used Chi-square tests (for caries prevalence) and Mann–Whitney U- or Kruskal–Wallis H-tests where appropriate (for caries severity). P-values of less than 0.05 were considered to be statistically significant. Zero-inflated negative binomial (ZINB) regression was used to model dmfs and control for potential confounding factors. This approach was used because, unlike ordinary linear regression, the negative binomial model provides a superior 'fit' to

	Age	[n (%)]	Sex [n (%)]		
Variable	5 years	6 years	Male	Female	
Father's education	:				
primary school	104 (21.0)	119 (24.7)	109 (22.4)	114 (23.3)	
high school	212 (42.8)	208 (43.2)	204 (42.0)	216 (44.1)	
university	179 (36.2)	154 (32.0)	173 (35.6)	160 (32.7)	
Mother's educatio	n:				
primary school	157 (31.3)	106 (22.2)*	124 (25.6)	139 (28.1)	
high school	244 (48.7)	241 (50.4)	235 (48.5)	250 (50.6)	
university	100 (20.0)	131 (27.4)	126 (26.0)	105 (21.3)	
Monthly income (	Dhs):				
1000-3000	108 (21.5)	165 (33.4)*	136 (27.3)	137 (27.5)	
3001-7000	169 (33.7)	167 (33.8)	164 (32.9)	172 (34.5)	
> Dhs 7000	225 (44.8)	162 (32.8)	198 (39.8)	189 (38.0)	
Nationality:					
Emirati	367 (70.8)	334 (66.3)	360 (69.8)	341 (67.4)	
Arab	110 (21.2)	114 (22.6)	102 (19.8)	122 (24.1)	
other	41 (7.9)	56 (11.1)	54 (10.5)	43 (8.5)	
$\overline{*P < 0.01}$ .					

 Table 1. Age and sex of subjects by parents' education, monthly income and nationality.

the dmfs distribution, and it is more informative because it allows the simultaneous modelling of both the prevalence and severity of caries [8]. Variables which had shown statistically significant bivariate associations with dental caries experience were entered into the ZINB model (in Stata) for dmfs.

## Results

The total number of children sampled was 1297, of whom 1036 completed both the dental examination and the questionnaire by their parents, giving a participation rate of 79.9%. There were 518 males (50.0%) and 518 females (50.0%); 524 (50.6%)were 5-year-olds, and 512 (49.4%) were 6-year-olds. There were 248 (23.9%) 5-year-old and 270 (26.1%) 6-year-old males, and 276 (26.6%) 5-year-old and 242 (23.3%) 6-year-old females. Data on the distribution of participants by age, sex, parental education, monthly income and nationality are presented by age and sex in Table 1. A statistically significant association was found between maternal education and the child's age, with a higher percentage of the 6-year-old children in this study having mothers with a high education level. A higher percentage of the 5-year-old children (44.8%) came from high-income families.

The mean dmft was 4.4 (SD = 4.3, range = 0-20). Data on the severity of dental caries are presented in Table 2. Overall, the average dmfs score (10.2; SD = 13.0, range = 0-85) was dominated by the decayed component, with the ds contributing 91%

Table 2. Mean number  $(\pm SD)$  of dmfs, ds, ms and fs by age, sex, mother's education, monthly income and nationality.

$10.2 \pm 13.0$ $9.2 \pm 12.6*$	$9.3 \pm 12.1$	$0.5 \pm 2.4$	$0.4 \pm 1.4$
$9.2 \pm 12.6*$	0.2 + 11 (*		
$9.2 \pm 12.6*$	0.0 1 11 6*		
	$8.3 \pm 11.6^{*}$	$0.4 \pm 2.5$	$0.4 \pm 1.3$
$11.3 \pm 13.3$	$10.3 \pm 12.6$	$0.5 \pm 2.3$	$0.5 \pm 1.5$
$11.3 \pm 13.6**$	$10.2 \pm 12.7$	$0.6 \pm 2.6$	$0.5 \pm 1.5$
$9.3 \pm 12.3$	$8.4 \pm 11.4$	$0.4 \pm 2.2$	$0.4 \pm 1.2$
n:			
$13.2 \pm 14.3*$	$12.4 \pm 13.6*$	$0.5 \pm 2.6$	$0.3 \pm 1.2$
$9.6 \pm 12.3$	$8.5 \pm 11.2$	$0.6 \pm 2.6$	$0.4 \pm 1.3$
$8.5 \pm 12.3$	$7.5 \pm 11.7$	$0.4 \pm 1.9$	$0.6 \pm 1.8$
Dhs):			
$12.7 \pm 13.9*$	$11.8 \pm 13.0*$	$0.6 \pm 2.3$	$0.3 \pm 1.3b$
$10.6 \pm 14.3$	$9.8 \pm 13.6$	$0.4 \pm 1.9$	$0.4 \pm 1.3$
$8.3 \pm 10.8$	$7.2 \pm 9.7$	$0.5 \pm 2.7$	$0.6 \pm 1.5$
$10.8 \pm 13.3 **$	$9.9 \pm 12.5 **$	$0.5 \pm 2.5$	$0.4 \pm 1.4$
$8.7 \pm 11.8$	$7.6 \pm 10.9$	$0.5 \pm 2.2$	$0.5 \pm 1.6$
$9.6 \pm 13.3$	$8.9 \pm 12.4$	$0.5 \pm 2.0$	$0.2 \pm 0.6$
	$\begin{array}{c} 1.3 \pm 13.6^{**} \\ 9.3 \pm 12.3 \\ 1. \\ 3.2 \pm 14.3^{*} \\ 9.6 \pm 12.3 \\ 8.5 \pm 12.3 \\ \text{Dhs}): \\ 2.7 \pm 13.9^{*} \\ 0.6 \pm 14.3 \\ 8.3 \pm 10.8 \\ 0.8 \pm 13.3^{**} \\ 8.7 \pm 11.8 \\ 9.6 \pm 13.3 \end{array}$	$\begin{array}{c} 1.3 \pm 13.5 \\ 10.3 \pm 12.6 \\ 1.3 \pm 13.6 \\ 10.3 \pm 12.3 \\ 10.3 \pm 12.7 \\ 9.3 \pm 12.3 \\ 12.4 \pm 13.6 \\ 12.4 \pm 13.6 \\ 12.4 \pm 13.6 \\ 12.3 \\ 8.5 \pm 11.2 \\ 8.5 \pm 12.3 \\ 7.5 \pm 11.7 \\ 11.8 \\ 13.6 \\ 13.5 \\ 11.8 \pm 13.0 \\ 11.8 \pm 13.0 \\ 13.6 \\ 13.4 \\ 1$	$1\cdot3\pm 13\cdot 5^*$ $10\cdot 3\pm 12\cdot 6^*$ $0\cdot 3\pm 2\cdot 5^*$ $1\cdot3\pm 13\cdot 6^{**}$ $10\cdot 3\pm 12\cdot 7$ $0\cdot 6\pm 2\cdot 6$ $9\cdot 3\pm 12\cdot 3$ $8\cdot 4\pm 11\cdot 4$ $0\cdot 4\pm 2\cdot 2$ $1:$ $3\cdot 2\pm 14\cdot 3^*$ $12\cdot 4\pm 13\cdot 6^*$ $0\cdot 5\pm 2\cdot 6$ $9\cdot 6\pm 12\cdot 3$ $8\cdot 5\pm 11\cdot 2$ $0\cdot 6\pm 2\cdot 6$ $9\cdot 6\pm 12\cdot 3$ $8\cdot 5\pm 11\cdot 2$ $0\cdot 6\pm 2\cdot 6$ $8\cdot 5\pm 12\cdot 3$ $7\cdot 5\pm 11\cdot 7$ $0\cdot 4\pm 1\cdot 9$ Dhs): $2\cdot 7\pm 13\cdot 9^*$ $11\cdot 8\pm 13\cdot 0^*$ $0\cdot 6\pm 2\cdot 3$ $0\cdot 6\pm 14\cdot 3$ $9\cdot 8\pm 13\cdot 6$ $0\cdot 4\pm 1\cdot 9$ $8\cdot 3\pm 10\cdot 8$ $7\cdot 2\pm 9\cdot 7$ $0\cdot 5\pm 2\cdot 7$ $0\cdot 8\pm 13\cdot 3^{**}$ $9\cdot 9\pm 12\cdot 5^{**}$ $0\cdot 5\pm 2\cdot 5$ $8\cdot 7\pm 11\cdot 8$ $7\cdot 6\pm 10\cdot 9$ $0\cdot 5\pm 2\cdot 2$ $9\cdot 6\pm 13\cdot 3$ $8\cdot 9\pm 12\cdot 4$ $0\cdot 5\pm 2\cdot 0$

\*\*P < 0.05.

of the total score. Caries severity was greater among males and children of less-educated mothers. Children in high-income families had the least dental caries. Emirati (local) children had higher caries severity than the others. Children from higher-income families had a greater fs score, on average, while the mean number of decayed surfaces was greater among children from lower-income households and those whose mothers had had less education.

Data on the use of dental services by age, sex, maternal education, monthly income and nationality are presented in Table 3. More 6-year-old children than 5-year-olds, more children of highly educated mothers and more Arab children had visited in the previous year. More of the 6-year-olds had had to visit the dentist because of a problem. There were no significant differences by the other sociodemographic characteristics.

The ZINB model for caries prevalence is presented in Table 4. Children from low-income families had substantially lower probability of being caries-free. Males had higher dmfs scores on average, as did children of mothers with lower levels of education and Emirati children. Children from high-income families or those whose last dental visit was for a check-up had considerably lower dmfs scores, on average. The latter were also more likely to be an extra zero (although this just fell short of statistical significance).

Variable	Visited dentist in previous year $[n \ (\%)]$	Problem was reason for dental visit $[n (\%)]$
All children	332 (32.0)	260 (25.1)
Age (years):		
5	163 (31.2)	109 (66.9)*
6	190 (37.4)	151 (79.5)
Sex:		
male	179 (34.8)	132 (73.7)
female	174 (33.7)	128 (73.6)
Mother's education	n:	
primary school	77 (28.8)	51 (66.2)
high school	166 (34.3)	127 (76.5)
university	109 (41.4)	81 (74.3)
Monthly income (	Dhs):	
1000-3000	92 (33.9)	67 (72.8)
3001-7000	113 (33.7)	85 (75.2)
> 7000	135 (34.9)	99 (73.3)
Nationality:		
Emirati	223 (31.9)	166 (74.4)
Arab	96 (43.2)	70 (72.9)
other	28 (28.9)	19 (67.9)

Table 3. Dental utilization of subjects by age, sex, mother's education, monthly income and nationality.

 $*P < 0{\cdot}05.$ 

 $*P < 0{\cdot}01.$ 

#### Discussion

This cross-sectional study has investigated the influence of sociodemographic factors on the dental caries experience of preschool children in order to provide information which would be useful in planning for a dental preventive programme. The main reason for selecting 5-year-olds was to enable comparison of UAE children dental caries status with previously published findings [3–5]. Although questionnaires are an accepted way of obtaining information in dental epidemiology, they have not been used widely in the UAE, and never before in the Emirate of Ajman. The questions were designed in English, and then translated by a linguist into

Arabic and back into English again ('back translation') for checking to ensure that the meaning of the questions stayed the same. The questionnaire used in this study was based on other published questionnaires previously used for the same purpose [9,10]. It was assumed that parental responses were accurate and honest, but this might be questioned since their validity and reliability were not examined; however, every effort was made to gain truthful answers. The parents were asked to respond to the best of their knowledge, and were told beforehand that there were no right or wrong answers; they just had to give honest responses.

In this study, over three-quarters of the children had experienced caries. This estimate is very close to that of Al-Mughery et al. [3], who reported that 72% of Abu Dhabi children had experienced caries. Estimates by Al-Hossani and Rugg-Gunn [4] were higher - they reported that 94%, 90% and 82% of Abu Dhabi, Al-Ain and Western Region children, respectively, had experienced caries - while Naqvi et al. [5] reported 79% among a similar age group in Al-Ain. Comparison with recent findings from studies in countries neighbouring UAE suggests that caries levels may be similar [11,12]. This estimate is considerably higher than the WHO [13] target for the year 2000 for 5-year-olds (i.e. 50% caries-free). In this study, the mean dmft was 4.4, which is only slightly less than the 5.1 dmft in 5-years-olds reported approximately 12 years previously [3], but somewhat lower than the 8.4, 8.6 and 5.7 dmft scores observed in Abu Dhabi. Al-Ain and the Western Region, respectively, among 5-year-old children [4]. Naqvi et al. [5] reported that 4- to 5-year-old Al-Ain children had a mean dmft of 5.8. The overall estimate of 7.1 dmft reported by Al-Hosani and Rugg-Gunn [4] is identical to the 7.1 dmft reported for 5-year-olds in Riyadh, Saudi Arabia [14]. It is

Table 4. Outcome of zero-inflated negative binomial modelling\* for decayed, filled or missed surface (dmfs) scores (95% confidence intervals in brackets).

	Probability of being					Adjusted
	Logit	P-value	caries free	Negative binomial	P-value	dmfs
Intercept	-1.235 (-1.752, -0.718)	0.00	0.23	2.285 (2.086, 2.484)	0.00	9.83
Male	0.136 (-0.270, 0.542)	0.51	0.25	0.201 (0.055, 0.347)	0.01	12.01
Low maternal education	-0.195 (-0.721, 0.330)	0.47	0.19	0.227 (0.054, 0.400)	0.01	12.33
High maternal education	0.092 (-0.393, 0.577)	0.71	0.24	0.014 (-0.178, 0.207)	0.88	9.97
Low-income family	-0.888 ( $-1.534$ , $-0.242$ )	0.00	0.11	0.088 (-0.098, 0.274)	0.35	10.73
High-income family	0.179 (-0.293, 0.651)	0.46	0.26	-0.262 ( $-0.442$ , $-0.081$ )	0.00	7.60
Emirati nationality	-0.459 (-0.960, 0.043)	0.07	0.16	0.207 (0.024, 0.389)	0.03	12.09
Last visit was for check-up	0.609 (-0.007, 1.225)	0.05	0.35	-0.342 (-0.619, -0.065)	0.02	6.98

\*Used to model dmfs and control for potential confounding factors.

Journal compilation © 2006 BSPD, IAPD and Blackwell Publishing Ltd

worth noting that all these studies utilized the WHO system for diagnosing and recording dental caries. These Arab-world estimates are much higher than those reported from eight European countries, which range from 0.8 dmft in Sweden to 3.1 dmft in Scotland [15]. It appears that, by international standards, caries prevalence is high among young Arab children.

Not surprisingly, caries experience in the current study was greater among the older children, suggesting that caries activity continues in these 5- to 6-year-old children. This is consistent with other studies conducted in various countries [16,17], and reflects the cumulative nature of caries. The observed gender difference in caries severity was in line with the findings of Al-Khateeb et al. [18], Al-Hosani and Rugg-Gunn [4], and Maciel et al. [19], all of whom found that, on average, girls had cleaner teeth and lower caries experience in the primary dentition. This could be related to the traditional practice of overindulging sons in Arabic culture, and exploration of this phenomenon using non-quantitative approaches is worthwhile. Further support comes from a study in Jeddah, Saudi Arabia, which showed that males had slightly higher dental caries scores than females [20]. However, a number of studies [21,22] have offered mixed findings in the presence of other non-biological exposure factors.

There was no existing classification of 'social class' or 'socioeconomic status' in the Emirate. Thus, arbitrary classifications of children by parental education and income were made. Notwithstanding this problem, this study showed a clear relationship between maternal education and caries occurrence, with children of highly educated mothers (and highincome families) having lower levels of caries. This finding is consistent with those from numerous other investigations in industrialized countries [23,24]. In Jordan [25] and in Saudi Arabia, the same relationship between social class and caries has now been well documented [12,26].

The nationality of children was also investigated as a risk marker for caries experience. On average, non-local children had lower caries scores than the local (indigenous) children, after controlling for factors such as education and family income might be responsible. Ethnic differences in caries experience have also been reported from a wide range of settings [27,28]. The majority of the non-local children included in this study were from different Arab countries and possibly different cultural backgrounds. Clarification of this point in order to compare the influence of social factors and dietary practices among foreign workers and indigenous Emiratis would be interesting and useful. However, it is beyond the scope of this study.

In many developing countries, access to oral health services is limited, and teeth are often left untreated or are extracted because of pain or discomfort. Control of oral disease depends on many factors, one of which is the availability and accessibility of oral health care. However, the reduction of disease is only possible if services are orientated towards primary health care and prevention [29]. In the current study, children who had received dental care had greater caries experience, on average. It is probable that these children had attended because of pain or symptoms, rather than as a result of any greater dental awareness of their parents. The same observation has been reported by Wong et al. [30] among 5- to 6-year-old children in China. Parents may have been unaware of the need of treatment, or perhaps, felt that their children were too young to attend a dental clinic, and therefore, they were not taken to the clinic until pain was experienced. It has been shown that the oral habits of families are formative, since parents are role models for their children [31]. Therefore, if parents do not, or cannot, attend for their own dental care, then this is likely to be the pattern for these children in the future [32].

Several studies have shown that children with caries in the primary dentition are more likely to develop caries in their permanent teeth [33,34]. Thus, it seems reasonable to presume that the most important target groups for instituting preventive dental care are infants and toddlers (and their families). The high caries prevalence and severity in young children in the Emirate of Ajman (and the surrounding countries) is of great concern; this study's mean of 9.2 dmfs at the age of 5 years is among the highest recorded in the literature. In addition, the level of disease appears to have increased in recent years [5]. Public health efforts are needed to reduce this high level of dental caries and improve the dental health of Ajmani children. While this study's findings offer many challenges for UAE dental policy planners and health promoters, they may be useful as a platform on which to raise the profile of child oral health. It is very important to convey the message that caries in preschoolers is a health problem that warrants the attention and resources of the community.

#### What this paper adds

• The findings of this study emphasized the importance of health promotion strategies, which target new and expectant mothers, and to promote a more positive attitude towards oral health.

### Why this paper is important to paediatric dentists

• Preventive strategy for specific populations should take into account social and cultural factors.

#### Acknowledgements

The authors would like to thank the staff of Ajman University for their assistance in this study.

#### References

- Kronfol NM. Perspectives on the health care system of the United Arab Emirates. *Eastern Mediterranean Health Journal* 1999; 5: 149–167.
- 2 Ministry of Health, Department of Planning. Annual Statistical Report. Abu Dhabi: Ministry of Health, 2001.
- 3 Al-Mughery AS, Attwood D, Blinkhorn A. Dental health of 5-year-old children in Abu Dhabi, United Arab Emirates. *Community Dentistry and Oral Epidemiology* 1991; **19**: 308–309.
- 4 Al-Hosani E, Rugg-Gunn A. Combination of low parental educational attainment and high parental income related to high caries experience in pre-school children in Abu Dhabi. *Community Dentistry and Oral Epidemiology* 1998; **26**: 31–36.
- 5 Naqvi A, Othman SA, Thabit MG. Baseline oral conditions in preschool children in Al-Ain Medical District. *Dental News* 1999; 6: 17–21.
- 6 World Health Organization. Oral Health Surveys: Basic Methods, 4th edn. Geneva: World Health Organization, 1997.
- 7 Caplan DJ, Slade GD, Gansky SA. Complex sampling: implications for data analysis. *Journal of Public Health Dentistry* 1999; **59**: 52–59.
- 8 Lewsey JD, Thomson WM. The utility of the zero-inflated Poisson and zero-inflated negative binomial models: a case study of cross-sectional and longitudinal MDF data examining the effect of socio-economic status. *Community Dentistry and Oral Epidemiology* 2004; **32**: 183–189.
- 9 Borssen E, Stecksen-Blicks C. Risk factors for dental caries in 2-year-old children. Swedish Dental Journal 1998; 22: 9–14.
- 10 Milgrom P, Riedy CA, Weinstein P, Tanner ACR, Manibusan L, Bruss J. Dental caries and its relationship to bacterial infection, hypoplasia, diet, and oral hygiene in 6- to 36-month-old children. *Community Dentistry and Oral Epidemiology* 2000; 28: 295–306.
- 11 Al-Amoudi N, Salako NO, Massoud I. Caries experience of children aged 6–9 years in Jeddah, Saudi Arabia. *International Journal of Paediatric Dentistry* 1996; 6: 101–105.
- 12 Al-Malik MI, Holt RD, Bedi R. The relationship between erosion, caries and rampant caries and dietary habits in preschool children in Saudi Arabia. *International Journal of Paediatric Dentistry* 2001; **11**: 430–439.
- 13 World Health Organization. *Prevention Methods and Programs for Oral Disease*. Technical Report Series 713. Geneva: World Health Organization, 1984.
- 14 Paul T, Maktabi A. Caries experience of 5-year-old children in Alkharj, Saudi Arabia. *International Journal of Paediatric Dentistry* 1997; 7: 43–44.
- 15 Bolin AK, Bolin A, Koch G. Children's dental health in Europe: caries experience of 5- and 12-year-old children from

eight EU countries. International Journal of Paediatric Dentistry 1996; 6: 155–162.

- 16 Hattab FN, Al-Omari MAO, Angmar-Mansson B, Daoud N. The prevalence of nursing caries in one-to-four-year-old children in Jordan. *Journal of Dentistry for Children* 1999; 1: 53–58.
- 17 Douglass JM, Tinanoff N, Tang JMW, Altman DS. Dental caries patterns and oral health behaviors in Arizona infants and toddlers. *Community Dentistry and Oral Epidemiology* 2001; 29: 14–22.
- 18 Al-Khateeb TL, Darwish SK, Bastawi AE, O'Mullance DM. Dental caries in children residing in communities in Saudi Arabia with differing levels of natural fluoride in the drinking water. *Community Dentistry Health* 1990; 7: 165–171.
- 19 Maciel SM, Marcenes W, Sheiham A. The relationship between sweetness preference, level of salivary mutans streptococci and caries experience in Brazilian pre-school children. *International Journal of Paediatric Dentistry* 2001; 11: 123–130.
- 20 Gandeh MBS, Milaat WA. Dental caries among schoolchildren: report of a health education campaign in Jeddah, Saudi Arabia. *Eastern Mediterranean Health Journal* 2000; 6: 396–401.
- 21 Harris R, Nicoll AD, Adair PM, Pine CM. Risk factors for dental caries in young children: a systematic review of the literature. *Community Dental Health* 2004; 21: 71–85.
- 22 Ismail AL, Shoveller J, Langille D, MacInnis WA, McNally M. Should the drinking water of Truro, Nova Scotia, be fluoridated? Water fluoridation in the 1990s. *Community Dentistry* and Oral Epidemiology 1993; 21: 118–126.
- 23 O'Brien M. Children's Dental Health in the United Kingdom 1993. London: HMSO, 1994.
- 24 Hinds K, Gregory JR. National Diet and Nutrition Survey: Children Aged 1-5 to 4-5 Years, Vol. 2: Report of the Dental Survey. London: HMSO, 1995.
- 25 Sayegh A, Dini EL, Holt RD, Bedi R. Food and drink consumption, sociodemographic factors and dental caries in 4–5-year-old children in Amman, Jordan. *British Dental Journal* 2002; **193**: 37–42.
- 26 Al-Dashti AA, Williams SA, Curzon MEJ. Breast feeding, bottle feeding and dental caries in Kuwait, a country with low-fluoride levels in the water supply. *Community Dental Health* 1995; 12: 42–47.
- 27 Wendt L-K, Birkhed D. Dietary habits related to caries development and immigrant status in infants and toddlers living in Sweden. Acta Odontologica Scandinavica 1995; 53: 339–344.
- 28 Gray M, Morris AJ, Davies J. The oral health of South Asian five-year-old children in deprived areas of Dudley compared with White children of equal deprivation and fluoridation status. *Community Dental Health* 2000; **17**: 243–245.
- 29 Petersen PE. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century the approach of the WHO Global Oral Health Programme. *Community Dentistry and Oral Epidemiology* 2003; **31**: 3–24.
- 30 Wong MCM, Lo ECM, Schwarz E, Zhang HG. Oral health status and oral health behaviors in Chinese children. *Journal of Dental Research* 2001; **80**: 1459–1465.
- 31 Mattila ML, Rautava P, Sillanpää M, Aunio P. Caries in five-year-old children and associations with family-related factors. *Journal of Dental Research* 2000; **79**: 875–881.
- 32 Evans RW, Lo ECM, Darvell BW. Determinants of variation in dental caries experience in primary teeth of Hong Kong children aged 6–8 years. *Community Dentistry and Oral Epidemiology* 1993; **21**: 1–3.
- 33 Kaste LM, Marianos D, Chang R, Phipps KR. The assessment of nursing caries and its relationship to high caries in the permanent dentition. *Journal of Public Health Dentistry* 1992; **52**: 64–68.
- 34 Thomson WM, Poulton R, Milne BJ, Caspi A, Broughton JR, Ayers KMS. Socio-economic inequalities in oral health in childhood and adulthood in a birth cohort. *Community Dentistry and Oral Epidemiology* 2004; **32**: 345–353.

Copyright of International Journal of Paediatric Dentistry is the property of Blackwell Publishing Limited and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.