Anxiety and Depression in Adolescents and Their Relationship with Signs and Symptoms of Temporomandibular Disorders

Leonardo Rigoldi Bonjardim, DDS, MS^a/Maria Beatriz Duarte Gavião, DDS, MS, PhD^b/ Luciano José Pereira, DDS, MS^a/Paula Midori Castelo, DDS, MS^a

> Purpose: Several clinical studies suggest that psychologic factors may play an important role in the etiology and maintenance of temporomandibular disorder (TMD) signs and symptoms. The goal of this study was to verify the prevalence of anxiety and depression in adolescents, and their relationship with signs and symptoms of TMD. Materials and Methods: The sample comprised 217 nonpatient adolescents between 12 and 18 years of age. The subjective symptoms and clinical signs of TMD were evaluated, respectively, using a self-report questionnaire and the Craniomandibular Index (CMI, which has 2 subscales), the Dysfunction Index, and the Palpation Index. The Hospital Anxiety and Depression Scale (HADS), a 14-item selfadministered rating scale that was developed specifically to identify anxiety and depression in nonpsychiatric medical outpatients, assessed levels of anxiety and depression. Results: In the total sample, anxiety and depression were present in 16.58% and 26.71% of subjects, respectively, including all levels of HADS. The results showed that there were positive correlations between CMI and Palpation Index and anxiety (HADSa) (P < .01) but not with depression (HADSd). An association between the number of TMD subjective symptoms and HADSa/HADSd was found (P < .01). **Conclusion:** Anxiety and depression, although of mild intensity, are common in adolescents. In this study both HADSa and HADSd were associated with an increasing number of TMD subjective symptoms. However, only anxiety was correlated with clinical signs of TMD (CMI), primarily muscle tenderness (Palpation Index). Int J Prosthodont 2005;18:347-352.

Temporomandibular disorders (TMD) is an all-inclusive term referring to a heterogeneous group of psychophysiologic disorders with the common characteristics of orofacial pain, masticatory dysfunction, or both.^{1,2} The etiology of TMD is considered to be multifactorial,²⁻⁴ but the relative importance of individual factors is still controversial. Psychosocial variables may play an important role in adaptation to pain and eventual recovery. TMD patients exhibit a variety of psychologic and behavioral characteristics, including increased somatization, stress, anxiety, and depression.^{2,5}

Some investigations have examined the relationship between psychologic factors and TMD.^{6–11} Studies of anxiety and depression in chronic pain patients have established that despite methodologic problems, there is evidence of greater prevalence of anxious and depressive symptoms in chronic pain populations than in controls.^{5,11,12} Since there is compelling evidence that chronic pain patients do not have a particular personality structure,^{13,14} it seems likely that anxiety and depression develop as sequelae to pain. Mood may, however, contribute to the endurance and impact of pain.¹⁵

^aPostgraduate Student, Department of Oral Physiology, Piracicaba Dental School, State University of Campinas (FOP/UNICAMP), Piracicaba, São Paulo, Brazil.

^bProfessor, Department of Pediatric Dentistry, Piracicaba Dental School, State University of Campinas (FOP/UNICAMP), Piracicaba, São Paulo, Brazil.

Correspondence to: Prof Dr Maria Beatriz Duarte Gavião, Faculdade de Odontologia de Piracicaba/UNICAMP, Departamento de Odontologia Infantil, Área de Odontopediatria, Av. Limeira 901 CEP 13414-903 Piracicaba, SP/Brasil. Fax: +55-19-3412-5218. E-mail: mbgaviao@fop.unicamp.br

A small but clinically challenging population of children and adolescents become chronic pain patients who report not only pain, but also associated emotional distress and disability.¹⁶⁻¹⁸ A picture is developing of the adolescent with chronic pain who also reports general emotional distress and a heightened sense of vulnerability.¹⁹ Functional TMD is often accompanied by mental symptoms such as depression, anxiety, and/or somatization on various levels.

This investigation was conducted to verify the prevalence and relationship of anxious-depressive symptoms with signs and symptoms of TMD and gender in adolescents. Furthermore, this research sought to identify whether these psychologic variables were associated with the severity of this disorder.

Material and Methods

Sample

Two hundred seventeen nonpatient adolescents (120 girls/97 boys), aged between 12 and 18 years, were randomly selected from public schools in Piracicaba, Brazil. Those who had received any type of orthodontic treatment were excluded before selection. The Ethics Committee of Piracicaba Dental School approved the research. The parents/guardians and the adolescents were informed about the purpose of this research, and signed informed consents were obtained.

Prior to the examination of clinical signs and symptoms, a questionnaire was filled out; this included questions about each individual's general state of health, illness, diseases, oral hygiene, and occlusion characteristics. The questionnaire and the clinical examination were conducted in appropriate rooms in the Public Schools following calibration of two of the author examiners (LRB and LJP).

Subjective Symptoms Interview

A self-report questionnaire was used to assess subjective symptoms according to Riolo et al²⁰ regarding pain in the jaws when functioning (eg, chewing), unusually frequent headaches (more than once a week), stiffness/tiredness in the jaws, difficulty in jaw opening, grinding teeth, and sounds from the temporomandibular joints (TMJs). Each question could be answered with a "yes" or a "no."

Evaluation of Clinical Signs

The signs of TMD were assessed by the authors on 1 occasion according to the Craniomandibular Index (CMI), as described by Fricton and Schiffman.²¹ The CMI has a 0 to 1 scale that measures tenderness and

dysfunction in the stomatognathic system and includes all currently recognized signs of TMD.^{21,22} It has 2 subscales: the Dysfunction Index (DI) and the Palpation Index (PI). The DI is designed to measure limitation in mandibular movement, pain and deviation in movement, TMJ noise, and TMJ tenderness. The PI measures the prevalence of muscle tenderness in the stomatognathic system. Thus, this index separates joint problems from muscle problems. The clinical examinations were conducted by two calibrated examiners (Kappa = .936).

Hospital Anxiety and Depression Scale

The level of anxiety and depression was self-rated by the Hospital Anxiety and Depression Scale (HADS). HADS consists of 7 items for depression (HADSd) and 7 items for anxiety (HADSa). The scale originated in the study of Zigmond and Snaith,²³ and a Portuguese version was validated by Botega et al.²⁴ The main characteristic of HADS is that items covering somatic symptoms of anxiety and depression have been eliminated. By defining cutoff values, the HADS subscales can give an indication of mental disorder: 0 to 7 = normal; 8 to 10 = mild; 11 to 14 = moderate; and 15 to 21 = severe disorder.²³ HADS has been extensively tested and has well-established psychometric properties.²⁵

Statistical Analysis

Data were computerized and the SAS package (version 8.2; SAS Institute) was used for their analysis. The presence of anxiety and depression in both genders was calculated by percentage and the Chi-square test. Correlation between CMI, PI, DI, and HADSa and HADSd was tested using the Spearman correlation. Associations between the number of symptoms and HADSa and HADSd were calculated using the Mantel-Haenszel Chi-square test. The significance level was set at P < .05 for all analyses.

Results

Table 1 shows that ages 12 to 14 were most prevalent in the studied sample, and the number of males and females were similar in all age groups. Maximum, minimum, and mean values for CMI, PI, and DI in both genders are expressed in Table 2. The mean scores for the indexes were low, because the majority of the adolescents did not show significant signs and symptoms; this decreased the total mean. In fact, there was no statistical difference between genders (Mann-Whitney, P >.06). However, in Table 2 it can be seen that some subjects presented with high scores. The distribution of anxiety and depression in both genders is presented in Table 3 and Table 4, respectively. Anxiety was more

	Age (y)															
		12		13		14		15		16		17		18	Т	otal
Gender	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)
Female Male	38 38	17.51 17.51	47 28	21.65 12.9	24 18	11.05 8.29	4 4	1.84 1.84	1	0.46 0.46	4	1.84 1.38	2 5	3.68 2.30	120 97	55.3 44.7
Total	76	35.02	75	34.56	42	19.35	8	3.68	2	0.92	7	3.22	7	3.22	217	100

Table 1Age and Gender Distribution (n = 217)

 Table 2
 Maximum, Minimum, and Mean Values for DI, PI, and CMI

	Во	ys	Girls				
Scale	Mean	Range	Mean	Range			
DI	0.083	0-0.214	0.097	0-0.428			
PI	0.056	0-0.477	0.087	0-0.8181			
CMI	0.075	0-0.582	0.093	0-0.532			

Table 3 Gender Distribution According to Level of Anxiety (HASDa)

	Bo	ys	Gir	ls	To	tal
Anxiety level	n	(%)	n	%	n	(%)
Free of anxiety	73	75.3	108	90	181	83.4
Mild	22*	22.7	11*	9	33	15.20
Moderate	1	1.03	1	0.83	2	0.92
Severe		1.03	0	0	1	0.46

*P<.05.

Table 5Correlations Between CMI, DI, PI, and
HADSa/HADSd (r value)

	HADSa	HADSd
DI	0.08299	-0.00898
PI	0.25448*	0.03251
CMI	0.28093 [†]	0.00283

**P* < .0002; [†]*P* < .0001.

Table 7	Association Between Number of Subjective
Symptoms	and Depression (HADSd)

No. of	Free of depression			Mild		Mod	erate	Se	Severe		
symptor	ns	n	(%)	n	(%)	n	(%)	n	(%)		
0		94	43.32	13	5.99	1	0.46	0	0		
1		51	23.50	10	4.61	0	0	0	0		
2		23	10.60	5	2.30	0	0	1	0.46		
3		11	5.07	4	1.84	1	0.46	0	0		
4		2	0.92	1	0.46	0	0	0	0		

P = .0232.

classified as mild according to HADS. The correlation coefficients for CMI, PI, DI, and HADSa and HADSd are expressed in Table 5. A significant correlation was found between CMI/PI scores and anxiety (P<.01), but this correlation was not found for depressive symptoms. Thus, the correlation with anxiety was significant for muscle tenderness. The Mantel-Haenszel Chi-square test was used to examine the association among HADSa and HADSd and the number of subjective symptoms of TMD (Tables 6 and 7). The results showed that the number of subjective TMD symptoms was associated with anxiety and depression (P < .01). It

Table 4	Gender	Distribution	Accordi	ng to l	Level	of
Depressio	n (HASD)d)				

	Boys			Gir	ls	Tot	Total	
Depression level	n	(%)	_	n	%	n	(%)	
Free of depression	73	75.3		86	71.6	159	73.2	
Mild	20	20.6		17	14.1	37	17.04	
Moderate	4*	4.1		15*	13.3	20	9.21	
Severe	0	0		1	0.83	1	0.46	

*P<.05.

Table 6Association Between Number of SubjectiveSymptoms and Anxiety (HADSa)

No. of	Free o	of anxiety	N	lild	Mod	erate	Severe	
symptoms	n	(%)	n	(%)	n	(%)	n	(%)
0	89	41.01	13	5.99	6	2.76	0	0
1	40	18.43	14	6.45	7	3.23	0	0
2	21	9.68	3	1.38	4	1.84	1	0.46
3	8	3.69	6	2.76	2	0.92	0	0
4	1	0.46	1	0.46	1	0.46	0	0

P = .0013.

prevalent in boys (24.7%) than in girls (9.83%) (P<.05), and the degree "mild" was the most prevalent in both genders. According to HADSd, depression was found in 27.4% of girls and in 24.7% of boys, with no statistically significant difference (P>.05). Mild depression was predominant in boys, whereas in girls the "mild" and "moderate" degrees were most common. There was a significant difference in depression severity between genders, since the percentage of girls presenting with a moderate degree was significantly higher than boys (13.3% and 4.1%, respectively). The majority of subjects presenting anxiety and depression were

showed that subjects with a greater number of subjective TMD symptoms could present with more anxiety and depression symptoms.

Discussion

This study evaluated the presence of anxiety and depression in adolescents and the relationship of these characteristics with signs and symptoms of TMD. The HADS was chosen to assess anxiety and depression symptoms. The method used is considered an easy-to-use, self-reporting questionnaire to detect anxiety and depression in medical outpatients in either a nonpsy-chiatric hospital setting or in the general population.^{25,26} It has been extensively researched and validated, and findings have demonstrated good sensitivity, specificity, and correlation between HADS, other questionnaires, and structured interviews used to diagnose depression and anxiety disorders.^{27–29}

Subjects who were free of anxiety and depression comprised the majority of the sample (Tables 2 and 3). However, 16.58% and 26.71% presented with anxiety and depression respectively, predominantly of mild intensity. These results show that there is evidence of psychologic disturbances in the general adolescent population, as verified also by Wight et al.³⁰ Anxiety and depressive disorders have been found in children and adolescents, with the ratio ranging from 5.7% to 17.7%³¹ for anxiety and from 3% to 40%^{32,33} for depression, corroborating the present results. In this study, 15.2% and 1.38% of subjects presented with mild and moderate/severe anxiety symptoms, respectively. Since these results could be explained by the age of the sample, it may be that such anxiety symptoms might not yet be affecting their quality of life. With regard to depressive symptoms, moderate/severe depressive symptoms were found in 9.67% of the subjects; this is in agreement with Rushton et al,³⁴ who found that over 9% of 13,568 adolescents reported moderate to severe depressive symptoms. In the present study, the majority of the symptoms were mild; nevertheless, girls presented more moderate depression than boys. The larger number of adolescents in this sample who were age 12 to 14 suggests that the difference between genders could be a result of biologic variables (eg, hormonal characteristics), as maturation in girls typically occurs earlier than in boys. However, it is necessary to consider that adolescents experience many developmental challenges as they strive to separate from their parents, become independent, and establish their own identities. This period of biopsychosocial maturation creates the conditions for people experiencing a greater sense of hopelessness and despair than when they were younger.³⁵ List et al⁶ verified in adolescents with TMD a higher degree of stress and behavior problems than healthy individuals. The TMD group also reported more somatic complaints and sought health care more frequently compared to controls. Thus, it appears that adolescents suffering from TMD pain seem to be more sensitive and more somatically focused than their healthy peers.

The debate is ongoing regarding the possible predisposing, triggering, and/or worsening role played by some psychological disorders in TMD subjects.^{5,36} There are also numerous reports of greater levels of depression and anxiety in persons presenting temporomandibular complaints as compared to normal controls.^{5–7,10,37–39}

TMD patients are more emotionally reactive to environmental challenge than normal subjects,³⁹ and approximately one third of the persons seeking treatment for TMD were experiencing depression.^{40,41} A significant correlation was found between CMI/PI and anxiety (HADSa) but not DI and anxiety; however, the same results were not found for depression (HADSd). Because the subjects examined were not seeking treatment, most of them probably did not present great evidence of clinical signs of TMD, which was found by the low mean CMI score of 0.085, on a scale of 0 to 1, and their lives were not yet affected. Although the mean CMI score was low, there were some high scores in both genders for CMI, PI, and DI, as presented in Table 2. In addition, the scores for CMI and PI were greater than those for DI in both genders, meaning that muscle tenderness could be the factor in determining the difference.

Therefore, individuals with high CMI and PI scores also had the highest HADSa scores, showing that these score indexes, especially those that involve muscle tenderness, tend to be enhanced in anxious subjects. It was found that the frequency of muscle groups painful to palpation increased in proportion to the frequency of reported anxiety symptoms. Such observations are in keeping with the hypothesized link between psychologic factors and muscular activity in the facial area, as myofascial pain patients have shown the highest prevalence of both mood and anxiety psychopathology when compared with TMD-free, disc displacement, and joint disorder subjects.^{5,9,42}

Furthermore, anxiety may be an important factor in the perception of pain, and anxious subjects may increase attention to pain, thereby amplifying their perceived intensity. This evidence was confirmed by other studies,^{7,11,43-45} which indicated that anxiety is related to increased pain reports in clinical settings. In addition, the evaluation of the psychologic dimension of muscle tenderness and its associated disability could provide valuable information for the adequate management of these patients and for assessing treatment outcome. Nevertheless, the results of this research are in accordance with a number of studies that also suggest that the association between myofascial pain and anxiety disorders is stronger than the association between pain and depression.^{46–50}

Furthermore, these results determined a significant association between anxiety/depression and the number of symptoms, as shown in Tables 6 and 7, suggesting that as the number of subjective symptoms increases, the same happens with the proportion of individuals presenting anxiety and depression complaints; this corroborates the findings of Sipila et al.⁵¹ Such associations might be explained by the fact that TMD symptoms and orofacial pain could have a psychosocial impact on the adolescent's quality of life, as is the case in adult populations.⁵² The number of nonspecific pain conditions reported may be a predictor of psychosocial problems, depression, and somatization.⁵³ It has been observed that small elevations in anxiety, depression, and somatization have been consistently identified in TMD patients.⁵⁴ Although there is no consensus regarding the percentage of TMD patients in whom psychologic factors play a role, it is clear that such factors need to be taken into account, along with structural indicators, to properly diagnose and plan management strategies.

It was concluded that the signs and symptoms of TMD could be present in adolescents and psychologic variables may be related. However, these conclusions must be interpreted with caution, given the inherent shortcomings of correlation studies of this nature. Limitations include the selection protocol employed and the possible influence of gender on the recorded results. Additionally the known fluctuation in signs and symptoms of musculoskeletal disorders in a time-dependent context might have been better addressed by carrying out repeated clinical recordings. Nonetheless, the observed outcomes suggest the importance of recognizing the merits of psychologic screening of adolescents with a confirmed diagnosis of a TMD, where indicated.

Acknowledgments

Leonardo Rigoldi Bonjardim was the recipient of a CAPES (DF, Brazil) scholarship grant. The authors thank Prof Glaucia Maria Bovi Ambrosano for statistical analysis.

References

- Phillips JM, Gatchel RJ, Wesley AL, Ellis E III. Clinical implications of sex in acute temporomandibular disorders. J Am Dent Assoc 2001;132:49–57.
- McNeill C. Management of temporomandibular disorders: Concepts and controversies. J Prosthet Dent 1997;77:510–522.
- Pankhurst CL. Controversies in the aetiology of temporomandibular disorders. Part 1. Temporomandibular disorders: All in the mind? [review] Prim Dent Care 1997;4:25–30.

- Thilander B, Rubio G, Pena L, de Mayorga C. Prevalence of temporomandibular disorder and its association with malocclusion in children and adolescents: An epidemiologic study related to specified stages of dental development. Angle Orthod 2002;72: 146–154.
- Manfredini D, Bandettini di Poggio A, Cantini E, Dell'Osso L, Bosco M. Mood and anxiety psychopathology and temporomandibular disorders: A spectrum approach. J Oral Rehabil 2004;31:933–940.
- List T, Wahlund K, Larsson B. Psychosocial functioning and dental factors in adolescents with temporomandibular disorders: A case control study. J Orofac Pain 2001;15:218–227.
- Wahlund K, List T, Ohrbach R. The relationship between somatic and emotional stimuli: A comparison between adolescents with temporomandibular disorders (TMD) and a control group. Eur J Pain 2005;9:219–227.
- Wexler GB, Steed PA. Psychological factors and temporomandibular outcomes. Cranio 1998;16:72–77.
- Yap AU, Tan KB, Chua EK, Tan HH. Depression and somatization in patients with temporomandibular disorders. J Prosthet Dent 2002 ;88:479–484.
- Yap AU, Dworkin SF, Chua EK, List T, Tan KB, Tan HH. Prevalence of temporomandibular disorder subtypes, psychologic distress, and psychosocial dysfunction in Asian patients. J Orofac Pain 2003;17:21–28.
- de Leeuw R, Studts JL, Carlson CR. Fatigue and fatigue-related symptoms in an orofacial pain population. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2005;99:168–174.
- Brown GK. A causal analysis of chronic pain and depression. J Abnorm Psychol 1990; 99:127–137.
- Schnurr R, Brooke R, Rollman G. Psychosocial correlates of temporomandibular joint pain and dysfunction. Pain 1990;42:153–165.
- Wade J, Dougherty L, Hart R, Cook D. Patterns of normal personality structure among chronic pain patients. Pain 1992;48:37–43.
- Madland G, Feinmann C, Newman S. Factors associated with anxiety and depression in facial arthromyalgia. Pain 2000;84:225–232.
- Malleson PN, Connell H, Bennett SM, Eccleston C. Chronic musculoskeletal and other idiopathic pain syndromes. Arch Dis Child 2001;84:189–192.
- 17. Eccleston C, Malleson P. Management of chronic pain in children and adolescents. Br Med J 2003;326:1408–1409.
- Mohlin BO, Derweduwen K, Pilley R, Kingdon A, Shaw WC, Kenealy P. Malocclusion and temporomandibular disorder: A comparison of adolescents with moderate to severe dysfunction with those without signs and symptoms of temporomandibular disorder and their further development to 30 years of age. Angle Orthod 2004;74:319–327.
- Merlijn VPBM, Hunfeld JAM, van der Wouden JC, Hazebroek-Kampschreur AAJM, Koes BW, Passchier J. Psychosocial factors associated with chronic pain in adolescents. Pain 2002;101:33–43.
- Riolo ML, Brandt D, TenHave TR. Associations between occlusal characteristics and signs and symptoms of TMJ dysfunction in children and young adults. Am J Orthod Dentofacial Orthop 1987;92:467–477.
- Fricton JR, Schiffman EL. Reliability of a craniomandibular index. J Dent Res 1986;65:1359–1364.
- 22. Fricton JR, Schiffman EL. The craniomandibular index: Validity. J Prosthet Dent 1987;58:222–228.
- Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. Acta Psychiatr Scand 1983;67:361–370.
- Botega NJ, Bio MR, Zomignani MA, Garcia Jr C, Pereira WAB. Mood disorders among medical in-patients: A validation study of the hospital anxiety and depression scale (HAD). Rev Saúde Pública 1995;29:359–363.

- Mykletun A, Stordal E, Dahl AA. The hospital anxiety and depression scale (HADS): Factor structure, item analyses, and internal consistency in a large population. Br J Psychiatry 2001;179: 540–544.
- Hinz A, Schwarz R. Anxiety and depression in the general population: Normal values in the Hospital Anxiety and Depression Scale. Psychother Psychosom Med Psychol 2001 ;51: 193–200.
- Herman C. International experiences with the Hospital anxiety and depression scale—A review of validation data and clinical results. J Psychosom 1997;42:17–41.
- Bjelland I, Neckelmann D, Tangen Haug T, Dahl AA. A review of validation studies of the hospital anxiety and depression scale. J Psychosom Res 2002;2:69–77.
- Stordal E, Mykletun A, Dahl AA. The association between age and depression in the general population: A multivariate examination. Acta Psychiatr Scand 2003 Feb;107(2):132–141.
- Wight RG, Sepulveda JE, Aneshensel CS. Depressive symptoms: How do adolescents compare with adults? J Adolesc Health 2004;34:314–323.
- Yorbik O, Birmaher B. Pharmacological treatment of anxiety disorders in children and adolescents. Bull Clin Psychopharmacol 2003;13:133–141.
- Kashani JH, Carlson GA, Beck NC, et al. Depression, depressive symptoms and depressed mood among a community sample of adolescents. Am J Psychiatry 1987;144:931–934.
- Garrison C, Schluchter Md, Schoenbach VJ, Kaplan BK. Epidemiology of depressive symptoms in young adolescents. J Am Acad Child Adolesc Psychiatry 1989;28:343–351.
- Rushton JL, Forcier M, Schectman RM. Epidemiology of depressive symptoms in the National Longitudinal Study of Adolescent Health. J Am Acad Child Adolesc Psychiatry 2002;41:199–205.
- Ryan ND, Puig-Antich J, Ambrosini P, et al. The clinical picture of major depression in children and adolescents. Arch Gen Psychiatry 1987;44:854–861.
- Manfredini D, Cantini E, Romagnoli M, Bosco M. Prevalence of bruxism in patients with different research diagnostic criteria for temporomandibular disorders (RDC/TMD) diagnoses. Cranio 2003;21:279–285.
- McCreary CP, Clark GT, Merril RL, Flack V, Oakley ME. Psychological distress and diagnostic subgroups of temporomandibular disorder patients. Pain 1991;44:29–34.
- Carlson CR, Okeson JP, Falace DA, Nitz AJ, Curran SL, Anderson D. Comparison of psychologic and physiologic functioning between patients with masticatory muscle pain and matched controls. J Orofac Pain 1993;7:15–22.

- Curran SL, Carlson CR, Okeson JP. Emotional and physiologic responses to laboratory challenges: Patients with temporomandibular disorders versus matched control subjects. J Orofac Pain 1996;10:141–150.
- Kinney RK, Gatchel RJ, Ellis E, Holt C. Major psychological disorders in chronic TMD patients: Implications for successful management. J Am Dent Assoc 1992;123:49–54.
- 41. Glaros AG. Emotional factors in temporomandibular joint disorders. J Indiana Dent Assoc 2000–01;79:20–23.
- Sieber M, Grubenmann E, Ruggia GM, Palla S. Relation between stress and symptoms of craniomandibular disorders in adolescents. Schweiz Monatsschr Zahnmed 2003;113:648–654.
- Palermo TM, Drotar D. Prediction of children's postoperative pain: The role of presurgical expectations and anticipatory emotions. J Pediatr Psychol 1996;21:683–698.
- 44. Rhudy RL, Meagher, MW. Fear and anxiety: Divergent effects on human pain thresholds. Pain 2000;84:65–75.
- Ferrando M, Andreu Y, Galdon MJ, Dura E, Poveda R, Bagan JV. Psychological variables and temporomandibular disorders: Distress, coping, and personality. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2004;98:153–160.
- Lipchik GL, Holroyd KA, O'Donnell FJ, et al. Exteroceptive suppression periods and pericranial muscle tenderness in chronic tension-type headache: Effects of psychopathology, chronicity and disability. Cephalalgia 2000;20:638–646.
- 47. Velly AM, Gornitsky M, Philippe P. Contributing factors to chronic myofascial pain: A case-control study. Pain 2003;104:491–499.
- McWilliams LA, Cox BJ, Enns MW. Mood and anxiety disorders associated with chronic pain: An examination in a nationally representative sample. Pain 2003;106:127–133.
- 49. Esenyel M, Caglar N, Aldemir T. Treatment of myofascial pain. Am J Phys Med Rehabil 2000;79:48–52.
- Auerbach SM, Laskin DM, Frantsve LM, Orr T. Depression, pain, exposure to stressful life events, and long-term outcomes in temporomandibular disorder patients. J Oral Maxillofac Surg 2001;59:628–633, discussion 634.
- Sipila K, Veijola J, Jokelainen J, et al. Association between symptoms of temporomandibular disorders and depression: An epidemiological study of the Northern Finland 1966 Birth Cohort. Cranio 2001;19:183–187.
- Segu M, Lobbia S, Canale C, Collesano V. Quality of life in patients with temporomandibular disorders. Minerva Stomatol 2003;52:279–287.
- Yap AU, Chua EK, Dworkin SF, Tan HH, Tan KB. Multiple pains and psychosocial functioning/psychologic distress in TMD patients. Int J Prosthodont 2002;15:461–466.
- Dahlström L. Psychometrics in temporomandibular disorders. An overview [review]. Acta Odontol Scand 1993;51:339–352.

Prosthodontist

Department of Comprehensive Care Case Western Reserve University School of Dental Medicine Cleveland, OH

The Department of Comprehensive Care at Case Western Reserve University School of Dental Medicine invites applications for a full-time tenure track faculty position at the assistant/ associate professor level. Responsibilities include didactic and clinical teaching and research. Participation in the CWRU Dental Faculty practice is available. Candidates must have a DMD/DDS degree or equivalent, and advanced training in prosthodontics or equivalent. Research and clinical interest and/or experience in fixed and implant prosthodontics and adhesive dentistry is desired. Salary and rank commensurate with qualifications and experience.



SCHOOL OF DENTAL MEDICINE

Case Western Reserve University is an equal opportunity/affirmative action employer.

Applicants should send a curriculum vitae, and names of three references to:

Dr. Avishai Sadan, Chairman Department of Comprehensive Care Case Western Reserve University School of Dental Medicine 10900 Euclid Avenue Cleveland, OH 44106-4905 Copyright of International Journal of Prosthodontics is the property of Quintessence Publishing Company Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.