

The use of rodent models to investigate host–bacteria interactions related to periodontal diseases

Dana T. Graves¹, Daniel Fine², Yen-Tung A. Teng³, Thomas E. Van Dyke¹ and George Hajishengallis⁴

¹Department of Periodontology and Oral Biology, Boston University School of Dental Medicine, Boston, MA, USA; ²Department of Oral Biology, Division of Periodontology of the Eastman Department of Dentistry, University of Medicine and Dentistry of New Jersey, Newark, NJ, USA; ³Department of Microbiology and Immunology, Center for Oral Biology, School of Medicine and Dentistry, University of Rochester, Rochester, NY, USA; ⁴Center for Oral Health and Systemic Disease and Department of Periodontics, University of Louisville Health Sciences Center, Louisville, KY, USA

Graves DT, Fine D, Teng Y-TA, Van Dyke TE, Hajishengallis G. The use of rodent models to investigate host–bacteria interactions related to periodontal diseases. *J Clin Periodontol* 2008; 35: 89–105. doi: 10.1111/j.1600-051X.2007.01172.x.

Abstract

Even though animal models have limitations, they are often superior to in vitro or clinical studies in addressing mechanistic questions and serve as an essential link between hypotheses and human patients. Periodontal disease can be viewed as a process that involves four major stages: bacterial colonization, invasion, induction of a destructive host response in connective tissue and a repair process that reduces the extent of tissue breakdown. Animal studies should be evaluated in terms of their capacity to test specific hypotheses rather than their fidelity to all aspects of periodontal disease initiation and progression. Thus, each of the models described below can be adapted to test discrete components of these four major steps, but not all of them. This review describes five different animal models that are appropriate for examining components of host–bacteria interactions that can lead to breakdown of hard and soft connective tissue or conditions that limit its repair as follows: the mouse calvarial model, murine oral gavage models with or without adoptive transfer of human lymphocytes, rat ligature model and rat *Aggregatibacter actinomycetemcomitans* feeding model.

Key words: animal model; bacteria; bone; cytokine; infection; inflammation; mouse; murine; osteoclast; periodontitis; rat

Accepted for publication 30 October 2007

Animal models are necessary to prove cause and effect relationships and to test the potential of novel therapeutics. In periodontal disease, human studies are limited by the difficulty in establishing causal relationships between various factors and periodontal breakdown. Although animal models including those of periodontal disease are not perfect, the limitations in addressing mechanis-

tic questions are usually not as severe as those of in vitro or clinical studies. The periodontium is a complex set of tissues that is chronically exposed to large numbers of bacteria (Pihlstrom et al. 2005, Paster et al. 2006). There are many uncertainties about which models are appropriate due in part to open questions about the initiation and progression of periodontitis. It is well known that periodontal diseases are initiated by bacteria, and yet there is still controversy about which bacteria actually initiate the events that lead to irreversible breakdown of periodontal tissues (Pihlstrom et al. 2005, Sakamoto et al. 2005). Considerable attention has been paid to *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*, which have been linked to adult

periodontitis and localized aggressive periodontitis, respectively (Holt & Ebersole 2005, Fine et al. 2006). However, there is only indirect evidence that these bacteria are important periodontal pathogens. Similarly, it is not known whether they represent important constituents in a polymicrobial infection, and if the infection is polymicrobial what other bacteria participate. The choice of bacteria has important implications as does the decision to study a single or polymicrobial infection.

It is recognized that the host response plays an important role in periodontal tissue breakdown and that it is likely to involve both innate and acquired immunity (Assuma et al. 1998, Baker et al. 1999, Teng et al. 2000, Delima et al. 2002). One of the most important

Conflict of interest and source of funding statement

The authors declare that they have no conflict of interests.

The authors were supported by NIH grants DE07559, DE17732, DE016306, DE14473, DE15786, DE16191 and DE15566, DE015254 and DE018292.

uncertainties regarding periodontal disease is its chronic nature. By analogy, periodontal diseases may be like a visit from unwanted relatives. They may move in for visits that are short, with each visit representing an acute insult that accumulates so that it appears to be chronic over time. Alternatively, the relatives may move next door, representing a relatively constant mild insult that becomes problematic with time. The former is consistent with the burst model of periodontal breakdown and the latter is consistent with the slow continuous model (Socransky et al. 1984, Cohen & Ralls 1988). To date, it is not known whether either or both models are accurate (Gilthorpe et al. 2003). If the burst model is more accurate, it is also not known whether the burst lasts for days or weeks, and if the chronic model is more appropriate, it is not known how long the chronic destructive period lasts. In fact, these two scenarios are not mutually exclusive. This is relatively important because the time frame in which periodontal breakdown takes place is not known and hence, assumed advantages of a given model regarding chronicity may be illusory. Another significant issue is whether the response represents a primary or a secondary exposure to a pathogen. In some models such as the oral gavage and *A. actinomycetemcomitans* rat-feeding model, exposure to the pathogen occurs over a period of several weeks, suggesting that there is secondary exposure. Similarly, in the rat ligature model the ligature facilitates invasion of endogenous flora. It may be presumed that the response represents a secondary exposure because it is likely that the animals have been previously exposed to their endogenous flora. In the calvarial and airpouch models, there is a capacity to examine primary and secondary responses specifically. A secondary response can be initiated by immunizing the mice before inoculation while a primary response would be initiated without prior immunization.

Rodent models have several useful features for investigating the molecular mechanisms involved in the pathogenesis of infection-driven inflammatory diseases. Indeed, there is considerable background information on mouse and rat immune systems and a wide range of immunologic and cellular reagents are available. Mouse models are particularly informative in studying downstream events associated with the host

because of the number of transgenic mice and mice with targeted genetic deletions. Both mouse and rat models provide considerable insight when examining the impact of overlying systemic issues such as diabetes.

In designing studies that assess the various steps required for infection and periodontal disease, several parameters need to be considered. It is clear that the infectious process occurs in several distinct but interconnected steps. Examining each step independently and then interdependently can provide a unique window into the disease process. Many lessons have been learned from experiments using rats and hamsters designed to assess the role of *Streptococcus mutans* in a caries infection (Fitzgerald & Keyes 1960). In a serendipitous finding, Keyes discovered that albino hamsters were caries free as compared with golden hamsters who were caries prone (Keyes 1959). After isolating an *S. mutans*-like, acid-producing microorganism, the authors were able to show that transferring faeces (containing *S. mutans*) from golden hamster cages to cages containing albino hamsters would result in newly formed carious lesions when these albino hamsters were given a soft sucrose containing a chow diet (Keyes 1960). These studies were then supplemented by other studies using a large number of different in vivo and in vitro models to investigate the mechanisms by which environmental factors (fluoride, etc.) and host factors (saliva, plaque fluid, etc.) can modulate caries (Aoba 2004, Caufield et al. 2005). These elegant experiments illustrate that (a) animal models can determine the role of both the appropriate host and a pathogenic microbe in a transmissible infection, (b) models are useful in understanding the mechanisms by which a bacterium can cause disease and (c) models are useful in understanding how the host can modify the disease process. With some modification, the lessons learned from caries models can be applied to animal studies of periodontal disease.

The limitation of all animal models is that there is no single model that represents all aspects of human periodontal disease. On the other hand, mechanistic questions cannot be typically addressed in human studies due to important ethical considerations. Thus, animal studies are critical for establishing cause and effect relationships and just as importantly, for initial tests of principle for the

development of advanced therapeutics. The goal here is not to define which model comes closest to human disease because periodontal diseases have multiple components but rather to better define how various models can be used to provide insight into the mechanisms of periodontal diseases. Thus, the most important issue is whether a given model is suitable for studying a specific hypothesis.

Most animal models of periodontal disease tend to focus on specific clinical features of gingivitis and periodontitis or the growth of potential etiologic bacteria in situ. However, in other disease systems, a series of models have emerged that emphasize measurement of the host response to various challenges. Several of these have been adapted for use as models to study processes relevant to host–bacteria interactions that shed light on similar processes that take place in the periodontium. The impact of bacteria on the host can be studied at several levels including colonization, host–bacteria interactions, loss of connective tissue attachment, bone resorption and coupling, the process whereby bone formation occurs following resorption to limit bone loss. It may be that one model is not suitable to study all aspects. For example, if the goal is to examine the potential of gingipains to reduce cytokine levels in connective tissue through degradation, it may not be necessary to examine the complicating effect of bacterial colonization. Owing to the complicating effect of multiple variables (e.g., colonization, connective tissue invasion, etc.), the hypothesis may be tested most directly by injecting known levels of bacteria or bacterial mutants lacking gingipains directly into connective tissue rather than relying on a more complex model where the variables are difficult to control. In some cases, the goal may be to study the bacterial genes needed in early biofilm formation so that periodontal breakdown may be irrelevant. The selection of a model should therefore be optimized to the goals of the study and should not necessarily reflect all aspects of periodontal disease. That said, an appropriate model is likely to provide more information than in vitro studies where complex cellular interactions are typically ignored or human studies where mechanisms are difficult to prove. Thus, we propose that one should not evaluate animal models with respect to whether they are closest

to human periodontal disease, but whether they are appropriate for testing a specific hypothesis.

Calvarial Model

The calvarial or scalp model was developed by Brendan Boyce to study the effect of cytokines on osteoclastogenesis (Boyce et al. 1989). It has been adapted to study the effect of bacteria on bone resorption (Zubery et al. 1998) or host–bacteria interactions in connective tissue in vivo (Graves et al. 2001, Leone et al. 2006). In this model, a stimulus is injected directly into the connective tissue overlying the calvarial bone in a small volume (e.g., 25 µl) of carrier. Classic inflammatory events occur including the rapid expression of pro-inflammatory cytokines within a few hours and the recruitment of PMNs within 24 h (Graves et al. 2005b). Bone resorption is induced within 3–5 days depending on the size of the stimulus (Li et al. 2002b, He et al. 2004). If the stimulus is small there is an inflammatory event that resolves with relatively little tissue destruction. On the other hand, a large inoculum can create a perturbation large enough to induce a soft tissue wound and considerable bone resorption that subsequently undergoes repair. New bone formation can then be examined as part of the healing process and can be measured 8–12 days following inoculation of the stimulus (He et al. 2004, Liu et al. 2006a). Thus, a potentially important use of this model is the ability to examine coupling, the process of reparative bone formation that occurs after bone resorption. The precise time frame in which events are measured may be modified by the strain of mouse (Battaglino et al. 2007). The use of an outbred mouse strain, CD-1, avoids some of the extremes of inbred strains. Thus, the calvarial model is useful in examining the sequence of inflammation, destruction and repair. This may be particularly important in periodontal disease where the net loss of tissue may be related to loss of balance between destruction of bone and connective tissue and the degree of repair that occurs. A summary of hypotheses that have been tested in the calvarial model and potential uses of the model are described in Table 1.

It is well known that bacterial products are chemotactic for leucocytes. An important issue that can be addressed

Table 1. Hypotheses tested in the calvarial model

Impact of the host response on hard and soft tissue damage	
Interleukin 1 (IL-1) induces osteoclastogenesis and bone resorption	Boyce et al. (1989)
Periodontal pathogens induce osteoclastogenesis and bone resorption partly through induction of prostaglandins	Zubery et al. (1998)
<i>Porphyromonas gingivalis</i> induces an inflammatory infiltrate and osteoclastogenesis indirectly via TNF	Graves et al. (2001)
Periodontal pathogens <i>P. gingivalis</i> and <i>Aggregatibacter actinomycetemcomitans</i> induce greater cytokine expression than a commensal bacterium, <i>Streptococcus gordonii</i>	Kesavalu et al. (2002)
Activation of the acquired immune response significantly increases expression of innate immune cytokines and an inflammatory infiltrate stimulated by <i>P. gingivalis</i>	Leone et al. (2006)
Bacterial virulence factors	
Low dose <i>P. gingivalis</i> LPS induces osteoclast formation through IL-11 and prostaglandins	Li et al. (2002b)
High dose <i>P. gingivalis</i> LPS induces osteoclasts through IL-1 and TNF activity	Chiang et al. (1999)
<i>P. gingivalis</i> fimbriae are pro-inflammatory but do not play an essential role in the innate immune response to <i>P. gingivalis</i> in connective tissue	Graves et al. (2005a)
<i>P. gingivalis</i> LPS, fimbriae and intact bacteria exhibit a qualitatively different stimulation of the innate immune response	Zhou et al. (2005)
LPS indirectly stimulates fibroblast apoptosis through TNF activity	Alikhani et al. (2003)
Impact of systemic disease (diabetes)	
Diabetes types 1 and 2 both cause prolonged inflammation in response to <i>P. gingivalis</i>	Graves et al. (2005b), Naguib et al. (2004)
Diabetes enhanced apoptosis limits the repair of a bacteria induced wound	Al-Mashat et al. (2006)
Diabetes enhanced TNF increases fibroblast & osteoblast cell death initiated by <i>P. gingivalis</i> , which significantly interferes with repair of connective tissue and bone	Liu et al. (2006a)
Diabetes interferes with the coupling of bone resorption and formation following bacteria induced bone loss	Al-Mashat et al. (2006)
Potential and future uses of this model	
To investigate the role of specific virulence factors by genetic manipulation of bacterial genes in stimulating or evading a host response in connective tissue in vivo	
To investigate the role of specific genes in mounting an innate immune response to oral bacteria in connective tissue in vivo using genetically modified mice	
To investigate the role of specific genes in mounting an acquired immune response to oral bacteria in connective tissue in vivo using genetically modified mice	
To investigate host mechanisms or systemic factors that modulate repair following bacteria induced injury to connective tissue	
To investigate mechanisms or systemic factors that affect osseous repair following bacteria induced bone resorption	

with the calvarial model is the relative contribution of bacterial products or induced host cytokines in stimulating the recruitment of inflammatory cells. This was tested by inoculation of *P. gingivalis* into the connective tissue overlying calvarial bone in wild-type or knockout mice lacking TNF receptors and measuring the impact of osteoclast formation (Fig. 1). The results indicate that there were significantly fewer osteoclasts in the absence of TNF receptor signalling compared with the normal mice, representing a substantial and significant decrease. This indicates that the TNF activity plays a critical role in bone resorption induced by *P. gingivalis*. TNF activity also played an impor-

tant role in the formation of an inflammatory infiltrate and fibroblast cell death stimulated by *P. gingivalis* (Graves et al. 2001). Furthermore, the results are consistent with the data obtained with canine and non-human primate models, which indicate that the host response plays an essential and vital role in periodontal disease destruction and is likely to be more important than the direct effect of bacteria in the loss of connective tissue and bone (Williams et al. 1985, Offenbacher et al. 1992, Assuma et al. 1998). These experiments also illustrate the advantage of murine models; a large number of genetically modified mice are available.




Fig. 1. Use of the calvarial model to demonstrate that *Porphyromonas gingivalis*-induced osteoclastogenesis in vivo is largely modulated through TNF. *P. gingivalis* or vehicle alone was inoculated into the scalp of TNFRp55^{-/-}/p75^{-/-} or control mice. Osteoclasts were identified as TRAP-positive multinucleated cells lining bone and quantified from sections taken 5 days after *P. gingivalis* inoculation. * $p < 0.05$ for *P. gingivalis* versus vehicle alone, ** $p < 0.05$ for TNFR^{-/-} versus WT. This figure was previously published in Graves et al. (2001).

Another approach to identifying the functional role of specific molecules in the host response to bacteria involves inhibitors to investigate cause and effect relationships. For example, TNF and caspase inhibitors have been used to establish the role in TNF and apoptosis in orchestrating the events that follow a *P. gingivalis*-induced insult (Al-Mashat et al. 2006, Liu et al. 2006a). Likewise, this model is useful in examining reparative bone formation following bone loss. For example, the injection of LPS causes bone resorption, followed by repair with net bone loss due to uncoupling. When fluoxetine is given, the amount of bone formation is sufficiently enhanced so that there is no net bone loss, demonstrating the anabolic effect of fluoxetine (Battaglino et al. 2007). If bone resorption and formation had not been measured separately, the simplest interpretation of the results would have been that fluoxetine blocks bone resorption. However, in this example, fluoxetine affected coupling so that reparative bone formation was the principal factor affected rather than resorption.

A second advantage of the calvarial model is the ability to deliver a precise amount of inoculum and to examine qualitative changes in the response to different levels of the same stimulus.

This has been investigated in the calvarial model using mice with targeted deletions as well as inhibitory antibodies. It has been shown that high doses of enteric LPS induce osteoclastogenesis that is substantially mediated by interleukin 1 (IL-1) and TNF receptor signalling. Lower doses of LPS, in contrast, induce osteoclastogenesis through other pathways that involve IL-11 (Chiang et al. 1999, Li et al. 2002a).

In addition to studying bone repair, the calvarial model can be used to examine soft tissue healing. As mentioned above, the inoculation of a sufficiently large dose of bacteria causes an inflammation-induced soft tissue wound that follows the classic steps of healing including the migration of fibroblasts into the wounded site, proliferation, production of extracellular matrix and reorganization (Liu et al. 2004). If the bacteria are killed by formaldehyde fixation, the inflammation-induced wound is sterile and healing is not affected by the ability to kill bacteria. We have used this model to quantify differences in healing in diabetic and normoglycaemic mice and found that one of the principal differences is a higher rate of fibroblast apoptosis that is related to high levels of TNF production in diabetic animals (Liu et al. 2006a). In fact, when fibroblast apoptosis is reduced with a caspase inhibitor, there is a significant improvement in soft tissue healing (Al-Mashat et al. 2006). It is striking that elevated fibroblast apoptosis has been implicated in poor healing that occurs in diabetic wounds (Darby et al. 1997). It is noteworthy that loss of fibroblasts is one of the most distinctive cellular changes that occurs in progressing periodontal disease (Zappa et al. 1992). In patients with periodontitis, fibroblasts have one of the highest rates of apoptosis in the gingiva and are observed predominantly in areas where inflammatory cells have been recruited (Koulouri et al. 1999). Thus, studies in the calvarial model point to potential novel difference in the response to a bacteria-induced injury in diabetic animals. In addition, the results of increased osteoblast and fibroblast apoptosis with concomitantly reduced numbers of these cells in diabetics are similar to the results obtained in other diabetic complications. For example, neuronal apoptosis is increased in diabetic neuropathy, cardiomyocyte apoptosis in cardiomyopathy associated with diabetes, mesangial

cell apoptosis in diabetic nephropathy and microvascular cell apoptosis in diabetic retinopathy (Graves et al. 2006).

By immunizing mice against a bacterium, the calvarial model can be used to examine events in connective tissue when the acquired immune response is added to the innate response (Leone et al. 2006). When *P. gingivalis* is inoculated, there is increased expression of several cytokines including IFN- γ , IL-6, macrophage inflammatory protein 2 and monocyte chemoattractant protein 1. Immunization against *P. gingivalis* enhances the mRNA levels of all these cytokines. Interestingly, greater cytokine and chemokine expression is associated with an enhanced and prolonged inflammatory cell infiltrate ($p < 0.05$), increased connective tissue destruction, greater osteoclastogenesis, increased bone loss, enhanced expression of proapoptotic genes and higher levels of fibroblast apoptosis ($p < 0.05$). The results indicate that activation of acquired immunity increases the destructive events stimulated by *P. gingivalis* in part, through up-regulating the innate immune response. These studies illustrate the utility of the calvarial model in examining specific aspects of the host-bacteria interactions in a connective tissue environment that retains complex cellular interactions.

Experimental periodontitis in a humanized non-obese diabetic (NOD)/severe combined immunodeficient (SCID) gavage model

Because most human periodontal pathogens are not inhabitants in mice and laboratory mouse strains, rendering an in vivo study of human periodontal disease difficult, humanization of mouse models provides an excellent opportunity to investigate the underlying processes or mechanisms regarding specific aspects of immune-bacteria interactions related to aspects of periodontal pathogenesis. Here, we summarize some specific hypotheses tested in a humanized immunodeficient NOD/SCID model in order to study human immunity-mediated pathology in vivo. Despite being considered exogenous pathogen(s) in mice under study, the secondary (or recall) immune responses and periodontal tissue inflammation against *A. actinomycetemcomitans* infection can be readily observed and measured in a humanized model using the oral gavage approach (see Table 2).

Immunodeficient SCID mice provide a unique in vivo model to study human lympho-haematopoietic development. Humanization of this model is achieved by engraftment of human peripheral blood leucocytes (HuPBL) into SCID mice (Mosier et al. 1988). An advantage of this mouse model is the ability to assess directly the function of human lymphoid cells (i.e., CD4⁺ and CD8⁺ T cells and B cells) in vivo. However, there are limited engraftment rates for other human cell lineages such as IgA⁺ B cells, granulocytes and neutrophils. In order to establish and boost T-cell immune responses, the host can be treated with an anti-NK (sialo-GM1) antibody that depletes NK cells to increase xenograft acceptances and chimerism (Sandhu et al. 1994, Shpitz et al. 1994). However, the depletion of NK cells with the antibody results in a much shorter life span (<4–5 weeks; Hozumi et al. 1994), which limits long-term in vivo studies. HuPBL-SCID mice can be used to test the protective effect of vaccines (i.e., for *P. gingivalis* infection) and to identify human B-cell epitopes for immune protection against *S. mutans*-induced dental caries (Senpuku et al. 1996, Choi et al. 1998). The fidelity of the adoptive transfer is supported by findings that HuPBL from aggressive periodontitis patients placed into SCID mice produce increased anti-leukotoxin IgG production in vivo (Shenker et al. 1993).

Another strain of SCID mice, NOD/SCID, has been generated, which is immunodeficient with multiple defects

in innate and adaptive immunity and lacking functional B and T cells, NK cells and haemolytic complement activity (Shultz et al. 1995). A prerequisite to achieving a high engraftment rate in humanized NOD/SCID mice involves screening of donor HuPBL; successful adoptive transfer of HuPBL-NOD/SCID mice can reach as high as 40–70% of cells in lymphoid organs/tissues such as the spleen (Mosier et al. 1991, Rohane et al. 1995). This is significantly higher than that detected in HuPBL-SCID mice, where endogenous mouse NK cells would attack the xenografts. It should be noted that both humanized SCID and NOD/SCID mice may not be useful to study the initiation of periodontal infection, as the majority of human lymphocytes transferred to the mouse in vivo exhibit a memory phenotype (being CD45RO⁺). Therefore, they are not suitable to study the early interactions between antigen-presenting cells, other innate immune cells and naïve T cells that are important in the initial stages of disease pathogenesis.

Oral infection of mice in this model is achieved by gavage without prior antibiotic treatment (Baker et al. 1994, Teng et al. 1999). Gavage with *A. actinomycetemcomitans* results in the formation of inflammatory infiltrates in the periodontal pocket and gingival connective tissue area consisting of T and B lymphocytes and CD14⁺ monocytes/macrophages. Like most of the small mouse models reported, the jaw samples can be examined by histology to assess bone

loss, formation of inflammatory infiltrates and by immunohistochemistry to identify the cell types involved (Teng et al. 1999, 2000). Early alveolar bone loss can be observed around 5 weeks with a peak loss at the end of 7–8 weeks. T cells can be collected and isolated from local mucosal tissues or cervical lymph nodes as described previously (Teng 2002b) and re-stimulated in vitro to further examine their responses and phenotypes. The humanized SCID or NOD/SCID models can also be used to examine immunization strategies or to test the contribution of specific aspects of lymphocyte function in vivo. For example, human B or T cells harvested from HuPBL-SCID or NOD/SCID mice can be screened by various techniques. This approach has been used to identify the genes expressed in *A. actinomycetemcomitans* that are associated with aggressive periodontitis such as the CagE-homologue and OPM-1 (Teng & Hu 2003, Teng & Zhang 2005). Alternatively, the induction of genes in lymphocytes can be examined by various mRNA profiling or proteomic techniques (see Table 2; and Y. T. Teng et al. unpublished data).

SCID mice infected with periodontal pathogens (i.e., *P. gingivalis*) produce significantly less alveolar bone loss than the immuno-competent wild-type mice (Baker et al. 1994, 1999). Similar findings occur in humanized NOD/SCID mice (Teng et al. 1999, 2000), suggesting the common features and validity of both model systems. Thus, the humanized NOD/SCID chimeras offer a unique approach to study specific aspects of periodontal pathogenesis such as cytokine profiling and a detailed analysis of lymphoid cell recruitment and participation in alveolar bone loss in vivo (Teng et al. 2000, Teng & Hu 2003, Teng & Zhang 2005, Zhang & Teng 2006). This model has been used to demonstrate that RANKL-mediated osteoclastogenesis and activation plays a significant role in alveolar bone loss in vivo (Teng et al. 2000, 2002a, 2005, Zhang & Teng 2006). Similarly, using a specific inhibitor, it was shown that diabetic NOD mice that manifest characteristics of human type-1 diabetes exhibit bone loss that is RANKL dependent (Mahamed et al. 2005). This suggests that the humanized mice can serve as a useful model to assess the efficacy of experimental therapeutics for specific aspects of periodontal pathogenesis related to bone loss in vivo (see Table 2

Table 2. Experimental periodontitis in humanized non-obese diabetic (NOD)/severe combined immunodeficient (SCID) gavage model

T cells significantly contribute to periodontal bone loss and tissue inflammation	Teng et al. (1999, 2000), Teng (2002b)
The cytokine, RANKL plays an important role in periodontal bone loss and its production is associated with both Th1 and Th2 responses during periodontal pathogenesis	Teng et al. (2000), Teng (2002b), Mahamed et al. (2005)
OPG administration significantly abolishes alveolar bone destruction in vivo; interferon- γ (IFN- γ) treatment increases periodontal bone loss while treatment with interleukin 10 (IL-10) reduces alveolar bone loss	Teng, Mahamed, & Singh (2005), Zhang & Teng (2006)
The antigen CagE found on <i>Aggregatibacter actinomycetemcomitans</i> , which is pro-apoptotic, plays a significant role in periodontal tissue and bone loss	Teng & Hu (2003), Teng (2003), Teng & Zhang (2005)
CD11c ⁺ dendritic cells can participate in modulating inflammation-induced bone loss by acting as osteoclast precursors	Alnaeeli et al. (2006, 2007)
Potential and future uses of this model	
(i) To investigate mechanisms by which diabetes exacerbates periodontal bone loss	
(ii) To identify susceptibility genes that lead to enhancing periodontal bone loss	
(iii) To examine the role of leucocyte subsets in promoting periodontal inflammation and alveolar bone loss	

and Fig. 2). For instance, engraftment of human foetal liver or thymic tissues into NOD/SCID facilitates testing human MHC-I and II-restricted immune responses against specific infections as well as super-antigens (Melkus et al. 2006). Lymphocyte recruitment in vivo may also be tested in humanized NOD/SCID mice in which the mice or the engrafted lymphoid tissue may be genetically modified to express specific genes (e.g., chemokines or chemokine receptors) to study dynamic processes in the acquired immune response (Shultz et al. 2007).

Oral gavage model of experimental mouse periodontitis

The introduction of human strains of bacteria by oral gavage and subsequent impact on the periodontium has been

studied in various rodent models (Chang et al. 1988, Klausen et al. 1991). A mouse periodontitis model was developed in which oral gavage with *P. gingivalis* induces reproducible alveolar bone loss (Baker et al. 1994, 1999, 2000b, Lalla et al. 1998). Bone loss induction in this model has been replicated using other periodontal pathogens, including *A. actinomycetemcomitans* (Garlet et al. 2006), *Tannerella forsythia* (Sharma et al. 2005) and *Porphyromonas gulae*, an animal periodontal pathogen considered to be equivalent to human *P. gingivalis* (Hardham et al. 2005). Typically, a susceptible mouse strain, such as BALB/cByJ, is orally administered a viscous suspension (2% carboxymethyl-cellulose) containing bacteria. Reproducibility and predictability are enhanced by prior treatment with antibiotics to

reduce the oral flora and with repeated inoculations of the implanted organism (e.g., three times at 2-day intervals with 10^9 CFU). Substantial bone loss typically takes more than 4 weeks to develop and can be measured histologically, by morphometric analysis as shown in Fig. 3, or by microcomputed tomography (Hart et al. 2004, Wilensky et al. 2005, Garlet et al. 2006). For bone loss assessment, the investigators most often examine the bone around the maxillary molars because induction of bone loss in the mandible is slower, whereas incisors are not included in the assessment due to their continuous eruption. In many reports, mice are euthanized 6 weeks after the final infection dose (Baker et al. 1999, 2000a, Yu et al. 2007), although it seems reasonable that antecedents could be investigated at earlier time points. In one analytical approach, the buccal and lingual gingival tissue surrounding the six maxillary molars are dissected for determination of cytokine induction at the mRNA (Hart et al. 2004) and/or the protein level (Garlet et al. 2006) and bone loss quantified by macro- or microscopic examination (Baker et al. 2000a, b, Burns et al. 2006, Garlet et al. 2006).

In the oral gavage model, *P. gingivalis* is undetectable before infection but can be recovered from the mouse oral cavity during the infection period (Baker et al. 2000b). The ability of *P. gingivalis* to establish an oral infection is facilitated by prior antibiotic-mediated suppression of the murine oral flora and may exploit inter-bacterial interactions with early colonizers as has been proposed for the human oral cavity (Jeon et al. 2005, Daep et al. 2006). Although vendors may claim that mice are pathogen-free, this does not necessarily mean that they are free of potential periodontal pathogens. In fact, at least one mechanism whereby *P. gingivalis* may contribute to periodontitis is perhaps by influencing inter-bacterial interactions so that it modifies the endogenous subgingival biofilm to become destructive. Such alterations may function synergistically to promote colonization, persistence and virulence of multi-species microbial populations, which may thereby acquire enhanced virulence. This is a distinct possibility, given that the oral gavage approach does not constitute a monoinfection model (unless mice are germ free). Indeed, the antibiotic treatment transiently




Fig. 2. Control of alveolar bone loss by *Aggregatibacter actinomycetemcomitans*-reactive CD4⁺ T cells. Groups of mice as indicated were either left untreated or inoculated with *A. actinomycetemcomitans*. CD4⁺ T, CD8⁺ T, or B cells were depleted from mice using antibody and complement as described (Teng et al. 2000). Data shown are the mean (\pm SD) of alveolar bone loss by time up to 8 weeks, expressed as percentage of bone loss detected in positive control, *A. actinomycetemcomitans*-infected BALB/c mice (as 100%). Group I, sham-infected, nondepleted NOD/SCID mice ($n = 10$); group II, *A. actinomycetemcomitans*-infected, nondepleted HuPBL-NOD/SCID mice ($n = 32$); group III, *A. actinomycetemcomitans*-infected, CD4⁺ T cell-depleted NOD/SCID mice ($n = 20$); group IV, sham-infected, CD4⁺ T cell-depleted NOD/SCID mice ($n = 9$); group V, *A. actinomycetemcomitans*-infected, CD8⁺ T cell-depleted NOD/SCID mice ($n = 12$); group VI, *A. actinomycetemcomitans*-infected, B cell-depleted NOD/SCID mice ($n = 16$); group VII, *A. actinomycetemcomitans*-infected NOD/SCID mice bearing adoptively transferred *A. actinomycetemcomitans*-reactive CD4⁺ T cells (AT-CD4T) plus autologous MO/MQ as APCs ($n = 10$); group VIII, *A. actinomycetemcomitans*-infected NOD/SCID mice bearing adoptively transferred autologous MO/MQ as APCs (irr-APC; $n = 6$). Statistical differences for bone loss between group III and groups II, IV, V, and VI were significant ($p < 0.005$). HuPBL, human peripheral blood leucocytes; NOD, non-obese diabetic; SCID, severe combined immunodeficient.


Mouse *P. gingivalis* Oral Gavage Model

Fig. 3. Periodontal bone loss in the oral gavage model. Assessment of bone levels is made by measuring the distance from the cemento-enamel junction (CEJ) to the alveolar bone crest (ABC) at the seven indicated buccal sites on the three molars on the left side and the symmetrical seven sites on the right side of the maxilla. To determine bone loss, the 14-site total CEJ–ABC distance for each infected mouse is subtracted from the mean CEJ–ABC distance from the group of sham-infected mice. (a) Sham-infected BALB/cByJ mouse (b) *Porphyromonas gingivalis*-infected BALB/cByJ mouse demonstrating increased CEJ–ABC distances at the molar sites, compared with sham infection. (c) Sham-infected C57BL/6J mouse (d) *P. gingivalis*-infected C57BL/6J mouse demonstrating negligible bone loss compared with sham infection, due to genetically determined resistance to bone loss. Originally published by Baker et al. (2000b).

suppresses but does not eliminate the normal oral flora, which gradually re-emerges. Additionally, *P. gingivalis* may modify the host response in ways that favour pathogen survival and disease development.

The oral gavage model has been used to establish proof of principle in vaccine development and to examine the impact of different components of the host response. For instance, immunization against RgpA and Kgp gingipain peptides or intact RgpA gingipain reduces *P. gingivalis*-induced bone resorption (Gibson & Genco 2001, O'Brien-Simpson et al. 2005). In an early study, Baker et al. showed a greater degree of *P. gingivalis*-induced bone loss in immunocompetent mice than in immunodeficient (SCID) mice lacking both T and B cells (Baker et al. 1994). In a follow-up study by the same group, mice deficient in MHC class II-responsive CD4⁺ T cells displayed increased resistance to *P. gingivalis*-induced bone loss, compared with normal mice or mice deficient in MHC class I-responsive CD8⁺ T cells (Baker et al. 1999). These findings indicate that at least certain aspects of the host response contribute to periodontal tissue destruction. Moreover, the severity of induced bone loss in this periodontitis model is dependent on both the genetic background of the mice and the particular *P. gingivalis* strain used, suggesting that the interplay of host and

microbial factors determines disease activity (Baker et al. 2000a, b).

Because of the complexity of this model, there are instances where the results need to be interpreted with caution because genetic modifications in mice could affect a number of parameters including colonization, invasion, downstream destructive events, repair or all four. In this regard, the observation that SCID mice display resistance to *P. gingivalis*-induced bone loss (Baker et al. 1994) could be interpreted to suggest that innate immunity may sufficiently protect against periodontitis, although the study was not specifically designed to test this hypothesis. This is consistent with another report that periodontitis-susceptible IL-17-receptor-knockout mice have reduced recruitment of neutrophils to the periodontium, which may be interpreted to mean that neutrophils represent a critical protective component of the host response (Yu et al. 2007). An alternative interpretation is that genetic modification indirectly leads to a more pathogenic flora as discussed in the same study (Yu et al. 2007).

Whereas more studies are warranted to conclusively associate the innate response with protection, it should be borne in mind that pathogens in general have evolved strategies to exploit and undermine innate immunity (Finlay & McFadden 2006). In this context, the oral gavage model appears to be appro-

priate for investigating aspects of immune evasion in periodontitis. Specifically, it was recently shown that TLR2 activation leads to enhanced *P. gingivalis* persistence when inoculated into chambers implanted into the abdomen, and TLR2-deficient mice display increased resistance to *P. gingivalis*-induced periodontal bone loss compared with normal controls (Burns et al. 2006). Although the precise underlying mechanisms are uncertain, at least partly, they may involve the ability of *P. gingivalis* to co-opt TLR2 inside-out signalling for activation of CD11b/CD18 (also known as CR3 for complement receptor-3) outside-in signalling, which inhibits IL-12 p70 production (Hajishengallis et al. 2005, 2006, Harokopakis & Hajishengallis 2005). Because IL-12 p70 plays a key role in intracellular bacterial clearance, this mechanism may presumably undermine the ability of the host to control an infection (Trinchieri 2003). In support of this notion, in vivo blockade of CR3 leads to enhanced IL-12 p70-mediated clearance of *P. gingivalis* and inhibition of its capacity to cause periodontal bone loss in the oral gavage model (Hajishengallis et al. 2007). Moreover, oral infection with *P. gingivalis* expressing a mutant version of fimbriae that interacts poorly with CR3 leads to severely attenuated bone loss, compared with the wild-type control (Wang et al. 2007). Interestingly, these in vivo findings from the oral gavage model are correlated with the in vitro virulence characteristics of the tested strains or the in vitro effects of CR3 blockade. Therefore, the oral gavage model seems to be appropriate for addressing hypotheses related to putative immune evasion strategies, identified from in vitro host-pathogen mechanistic studies.

The oral gavage model was moreover successfully used to link *P. gingivalis*-induced periodontitis to systemic conditions mechanistically. For example, oral infection of hyperlipidemic mice with *P. gingivalis* mice accelerates atherosclerotic plaque formation (Lalla et al. 2003, Gibson et al. 2004). Moreover, *P. gingivalis*-induced alveolar bone loss is enhanced in diabetic mice unless the receptor for advanced glycation end products is blocked (Lalla et al. 1998, 2000). It becomes obvious that the oral gavage model is useful for addressing a wide variety of hypotheses related to periodontal pathogenesis, ranging from the role of the host response

Table 3. Hypotheses tested in the oral gavage model

	References
Genetic basis for host susceptibility or resistance to periodontal disease	
Different genetic mouse strains display differential susceptibility to <i>Porphyromonas gingivalis</i> -induced periodontal bone loss	Baker (2005)
<i>Il1b</i> , <i>Tnf</i> , and <i>Stat6</i> are associated with susceptibility, whereas <i>Il15</i> and <i>Selp</i> with resistance to <i>P. gingivalis</i> -induced periodontal bone loss	Hart et al. (2004)
Role of specific cytokines or antimicrobial molecules in periodontal disease	
Increased susceptibility of interleukin 10 (IL-10) knockout mice to <i>P. gingivalis</i> -induced periodontal bone loss	Sasaki et al. (2000)
IL-17-receptor-knockout mice display increased susceptibility to <i>P. gingivalis</i> -induced periodontal bone loss	Yu et al. (2007)
Inducible nitric oxide synthase (iNOS)-knockout mice exhibit increased periodontal bone loss in response to <i>P. gingivalis</i> infection	Alayan et al. (2006)
Identification of potential virulence factors	
Wild-type <i>P. gingivalis</i> , but not a FimA-deficient mutant, induces periodontal bone loss and accelerates atherosclerosis	Gibson et al. (2004)
Diminished induction of periodontal bone loss by <i>P. gingivalis</i> mutant expressing FimA fimbriae lacking the FimCDE accessory components	Wang et al. (2007)
Diminished induction of periodontal bone loss by <i>P. gingivalis</i> mutants lacking expression of Kgp and/or RgpB	Pathirana et al. (2007)
Diminished induction of periodontal bone loss by <i>T. forsythia</i> mutant lacking BspA adhesin	Sharma et al. (2005)
Immune evasion through exploitation of innate receptors	
TLR2 deficiency attenuates <i>P. gingivalis</i> -induced periodontal bone loss	Burns et al. (2006)
Blockade of complement receptor-3 inhibits <i>P. gingivalis</i> -induced periodontal bone loss	Hajishengallis et al. (2007)
Periodontal disease connection with systemic diseases	
Oral infection with <i>P. gingivalis</i> accelerates atherosclerosis	Gibson et al. (2004), Lalla et al. (2003)
<i>P. gingivalis</i> -induced periodontal bone loss is enhanced in diabetes	Lalla et al. (1998)
Blockade of RAGE (receptor for advanced glycation end products) suppresses periodontal bone loss in diabetic mice	Lalla et al. (2000)
Proof-of-principle immunization studies to identify candidate vaccine antigens	
Systemic immunization using RgpA and Kgp gingipain peptides protects against <i>P. gingivalis</i> -induced periodontal bone loss	O'Brien-Simpson et al. (2005)
Systemic immunization using intact RgpA protects against <i>P. gingivalis</i> -induced periodontal bone loss	Gibson & Genco (2001)
Oral immunization with FimA-expressing <i>Streptococcus gordonii</i> vector protects against <i>P. gingivalis</i> -induced periodontal bone loss	Sharma et al. (2001)
Potential and future uses of this model	
To determine the role of pattern-recognition receptors in protection or susceptibility to periodontitis using genetically modified mice	
To determine the impact of aging in periodontal disease susceptibility using young and aged mice	
To investigate the protective potential of specific antagonists of disease-promoting mechanisms (e.g., destructive inflammation, bacterial evasion of immunity)	
To investigate the role of specific bacterial genes in interbacterial co-adherence and biofilm formation	

to virulence traits of suspected periodontopathogens and the interconnection of those factors with systemic parameters. A summary of such tested hypotheses is presented in Table 3, along with additional future uses of this model.

***A. actinomycetemcomitans* Rat-Feeding Model**

Colonization is a critical process in infection and is dependent on surface-related, tissue-specific adhesins that interact with host receptors. The

A. actinomycetemcomitans rat model of periodontal disease is particularly well suited to examine aspects of bacterial behaviour that promote colonization and initiation of periodontal disease (see Fig. 4). Scannapieco et al. (1987) demonstrated that clinical isolates of *A. actinomycetemcomitans* when grown on agar had a rough surface colonial morphology that attached with great avidity when grown in broth. Further, when these rough wild-type strains were passaged in the laboratory, the adherent strains often transformed into a smooth, gum drop-like colony that no longer attached to surfaces. These characteristics are seen in many *A. actinomycetemcomitans* strains in the ATCC culture collection. Moreover, it appears that conversion is irreversible and thus smooth colonies do not have the capacity to revert to their original rough adherent phenotype (Fine et al. 1999). In order to investigate the significance of rough versus smooth strains, a suitable infectious model is needed, as described below.

Seminal experiments by Mekalanos and colleagues highlighted the differences observed when microbes were removed from their natural environment and grown in the laboratory (Mahan et al. 1993). The use of promoter-reporter constructs established differences in gene expression between in vivo as compared with in vitro bacterial gene expression. This has led to identification of novel virulence gene sequences in the bacteria tested (Mahan et al. 1993). To test this concept in *A. actinomycetemcomitans*, a smooth laboratory strain of *A. actinomycetemcomitans* was placed in the mouths of rats to determine whether growth in an animal host could lead to conversion of smooth to rough by expression or up-regulation of genes in the smooth strain. To make this comparison, wild-type *A. actinomycetemcomitans* was passaged in broth to achieve a smooth, afibrillar, minimally adherent isogenic *A. actinomycetemcomitans* mutant. Both wild-type and smooth *A. actinomycetemcomitans* strains were made rifampicin resistant so that they could be readily identified in the mouths of rats. The results indicated that the smooth strain could survive in some animals for a 2–3-week period but not more than 4 weeks. On the other hand, the rough adherent strain was found over the 12-week experiment period and in some cases 7 months after


A. actinomycetemcomitans rat feeding model

Fig. 4. Time line for studying *Aggregatibacter actinomycetemcomitans* colonization and bone loss in the rat.

inoculation. No reversion was seen in any strain. Thus, none of the isogenic smooth strains recovered from the mouths of rats were found to be rough and none of the rough strains converted to smooth even though rough strains gave rise to the smooth strains in the laboratory in less than four passages (Fine et al. 2001).

At about the same time, it was discovered that avid attachment of *A. actinomycetemcomitans* to tooth surfaces related to a group of genes called the *tad* genes (for *right adherence*) (Kachlany et al. 2001). The assumption was that these genes were capable of enabling *A. actinomycetemcomitans* to attach to the teeth of rats because the *tad* genes conferred on these *A. actinomycetemcomitans* strains the ability to bind to abiotic and biological surfaces. A study testing this hypothesis showed that knocking out these *tad* genes and thus creating the equivalent of a smooth strain prevented the inoculated strains from colonizing and inducing bone loss. In contrast, the *A. actinomycetemcomitans* wild-type strain colonized, developed an antibody to *A. actinomycetemcomitans*, and stimulated bone loss. Each animal that was colonized by *A. actinomycetemcomitans* showed an increase in antibody titre and bone loss, while the *A. actinomycetemcomitans* knock-out strains showed neither

antibody titre nor bone loss (Schreiner et al. 2003).

The lessons learned from these experiments indicated that it is important to consider the existing flora, tagging the exogenous inoculated species, the housing of animals, the methods of recovery and the methods of feeding of animals. Experiments designed to add exogenous *A. actinomycetemcomitans* to an existing oral flora requires that the existing flora be reduced to facilitate colonization of the implanted strain. We also found it beneficial to use a tag so that the implanted strain could be distinguished from the existing flora. This is particularly useful to distinguish between rough and smooth forms of *A. actinomycetemcomitans*. Thus, when tested against the existing flora of the host animal, only the implanted rifampicin-resistant strain would survive in rifampicin-containing culture media because no other isolates obtained from the normal oral flora could survive. In this manner, the organism recovered is derived from the organism implanted. In addition to these considerations, housing of animals is a critical issue in this colonization and transmission model of disease. In this case, separate cages are required to avoid transmission of microorganisms from one animal to another. In feeding experiments where some animals are more aggressive than

others, co-habiting can lead to inadvertent transmission of microbes via, licking, touching, etc. Further considerable ports should be made to deliver a standard dose over a standard period of time (Fine et al. 2001). Mixing a given amount of *A. actinomycetemcomitans* in with the animal feed and caging animals separately allowed for a more quantitative estimate of the dosage given and the colonization efficiency.

More recently, we have become aware of the fact that both species and tissue specificity can be critical issues with respect to the choice of an animal model. With respect to *A. actinomycetemcomitans*, wild-type strains have been shown to have a narrow host range that includes humans, old world monkeys, cows and rats when tested against the buccal epithelium (Fine et al. 2005). Human and old-world monkey specificity was found to be due to two *A. actinomycetemcomitans* adhesions. Thus, *A. actinomycetemcomitans* adhesions Aae and ApiA, in addition to species specificity, demonstrated a narrow range of tissue tropism that limited attachment to buccal epithelial cells and the tongue and not to alveolar, nasal, skin or vaginal epithelium. Moreover, wild-type *A. actinomycetemcomitans* binds to rat buccal epithelial cells, and a review of the literature indicates that *A. actinomycetemcomitans* is a normal

inhabitant of a species of rice rat, making the rat an appropriate model for studies of *A. actinomycetemcomitans* colonization and infection (Fine et al. 2005). However, *A. actinomycetemcomitans* has difficulty in colonizing the oral cavity of mice. Thus, the mouse is not an appropriate model to study *A. actinomycetemcomitans* colonization, and studies with *A. actinomycetemcomitans*-induced bone loss in mice should keep in mind that the infection is transient and as a result, self-limiting. While *Actinobacillus actinomycetemcomitans*, recently renamed *A. actinomycetemcomitans*, has been isolated from the mouths of rats (MacDonald et al. 1959; Shaw et al. 1961), the exact species was only tentatively identified. Moreover, in our feeding model experiments, an extensive survey of the mouths of the Sprague–Dawley rats before feeding failed to reveal any pre-existing *A. actinomycetemcomitans*. As such, animals used in this rat-feeding model were infected by an exogenous source of *A. actinomycetemcomitans*. Nevertheless, the ability of rat buccal cells and salivary-coated rat teeth to bind *A. actinomycetemcomitans* in vitro, coupled with the finding of *Actinobacillus* in the mouths of laboratory rodents (Simpson & Simmons 1980), suggests that, unlike mice, *A. actinomycetemcomitans* can infect and inhabit the mouths of rats. As a result, one can extrapolate and suggest that *A. actinomycetemcomitans* is well suited to cause an indigenous infection (Macdonald et al. 1959) that can be altered by antibiotic administration (Shaw et al. 1961).

To overcome this issue of tissue and species specificity, Finlay and coworkers in a diarrhoea model have used *Citrobacter rodentium*, an enteropathogenic rat microbe with a phenotype similar to *Escherichia coli*. As a non-invasive gram-negative species of bacteria that colonizes the intestine of rodents and causes diarrhoea, these studies have advanced the understanding of human intestinal disease. These bacteria colonize by direct contact with the intestinal tissue, making it feasible to test adhesin/receptor interactions. It is also not surprising that different strains of rats may have different levels of susceptibility to infection and bone loss. Preliminary data from our laboratory suggest that three different strains of rats show different levels of bone loss even if the same *A. actinomycetemcomitans* strain and the same method of

inoculation of *A. actinomycetemcomitans* are used. This may well reflect differences in rat strains manifested as an expression of host genetic variability that could result in strain variation in the production of cytokine and in the robustness of the innate or acquired immune responses as has been shown in mice (Adelman et al. 1978, Darville et al. 1997, Gemmell et al. 2000). A summary of the hypotheses that have been tested and potential future uses of the *A. actinomycetemcomitans* rat-feeding model are described in Table 4.

Rat Ligature Model

The placement of ligatures around teeth to initiate periodontal tissue loss has been used in various types of animals. The ligature leads to greater accumulation of plaque and ulceration of the sulcular epithelium, facilitating invasion of connective tissue. Although this model was once commonly used with non-human primates, it is used less frequently due to the expense and difficulty in carrying out experiments with these animals. Because many of the same series of events occur in the rat as in the non-human primate, there has been renewed interest in the rat ligature model, particularly because it is simpler to work with and less expensive (Rovin et al. 1966). In the rat, loss of attachment and bone occurs predictably in a

7-day period (Nowotny & Sanavi 1983, Lohinai et al. 1998, Bezerra et al. 2000, Bezerra et al. 2002) although investigators have conducted experiments over much longer periods of time (Rovin et al. 1966, Weiner et al. 1979, Nociti et al. 2001, Kuhr et al. 2004). Like human periodontitis, alveolar bone loss in the ligature model is dependent on bacteria. In gnotobiotic rats, placement of ligatures does not induce significant gingival inflammation or periodontal bone loss (Rovin et al. 1966). Topical treatment with chlorhexidine reduces bone resorption and antibiotics reduce loss of attachment and loss of bone, supporting the role of bacteria in initiating destruction in this model (Weiner et al. 1979, Kenworthy & Baverel 1981). In contrast, increasing gram-negative bacterial burden enhances osteoclastogenesis and bone resorption (Samejima et al. 1990). A summary of the hypotheses that have been tested in the rat ligature model are described in Table 5 in addition to examples of hypotheses that could be tested in the future.

Like human periodontitis, the destructive phase of ligature-induced experimental periodontitis is associated with a host response as shown by the formation of an inflammatory infiltrate in the gingiva before bone resorption (Gyorfi et al. 1994). When the host response is diminished by inducing endotoxin tolerance, bone loss is

Table 4. Hypotheses tested *Aggregatibacter actinomycetemcomitans* rat feeding model

Ability of microbe delivered via oral feeding to induce an infection	Fine et al. (2001), Schreiner et al. (2003)
Ability of microbe to attach, colonize, and integrate into an established (albeit reduced) oral flora	Fine et al. (2001), Schreiner et al. (2003)
Recovery of microbe from animal and analysis of traits unique to its growth in that animal as compared to growth seen in laboratory cultures of same organism	Fine et al. (2001), Schreiner et al. (2003)
Microbial attachment factors essential for colonization of animal tooth and tissue Can be analysed	Schreiner et al. (2003), Sharma et al. (2001)
Microbial attachment and colonization can be related to antibody response and bone loss	Burckhardt & Guggenheim (1980), Schreiner et al. (2003), Sharma et al. (2001)
Analysis of effect of antagonistic or symbiotic organisms on growth and survival of test microbe	Hillman et al. (2000)
Potential and future uses of this model	
To investigate how mutants in known <i>A. actinomycetemcomitans</i> virulence traits (i.e., leukotoxin, cytolethal distending toxin) effect colonization, antibody responses and bone loss	
To identify relationship between specific and non-specific adhesions on attachment and colonization	
To determine how different strains of <i>A. actinomycetemcomitans</i> and other ‘‘pathogens’’ colonize in the rat-feeding model	
To determine how different strains of rats are affected by <i>A. actinomycetemcomitans</i> colonization; which strains are more susceptible to colonization; does colonization correlate to antibody response and bone loss	
To determine the immunopathological response in animals colonized by <i>A. actinomycetemcomitans</i> in different rat species	

decreased (Samejima et al. 1990). The functional role of the inflammatory response is further demonstrated by decreasing periodontal destruction in rats treated with prostaglandin inhibitors. The COX-1/COX-2 inhibitor, indomethacin, reduces gingival inflammation, osteoclast formation and bone loss in this model (Bezerra et al. 2000). The COX-2 inhibitor, meloxicam, has a similar effect (Bezerra et al. 2000). Low-dose doxycycline, which inhibits MMP activity, also reduces ligature-enhanced alveolar bone loss (Bezerra et al. 2002). The role of cytokines in the disease process is demonstrated by enhanced bone loss when ligature-induced periodontitis is augmented by application of IL-1 or TNF (Koide et al. 1995, Gaspersic et al. 2003). And lastly, the ligature-induced model is sensitive to systemic effects such as smoking, so that rats exposed to cigarette smoke have greater bone loss and higher MMP-2 activity than ligated rats not exposed to cigarette smoke (César Neto et al. 2004). Similar results have been obtained when ligated rats were injected with nicotine (Nociti et al. 2001, Benatti et al. 2003).

An advantage of the rat ligature model is the availability of systemic disease models. For example, the type-2 Zucker diabetic fatty (ZDF) rat can be used to examine the impact of diabetes. The ZDF rat develops obesity and insulin resistance, which results in diabetes by

8 weeks of age (Clark et al. 1983, Peterson 1995). When ligature-induced periodontal bone loss is induced in the ZDF rats and normoglycaemic littermates, diabetes increases the intensity and duration of the inflammatory infiltrate, enhances osteoclastogenesis and activity and increases bone loss (Liu et al. 2006b). Furthermore, by removing the ligatures, a healing response ensues in which coupling may be examined. When tested, the amount of new bone formation following resorption was 2.4–2.9-fold higher in normoglycaemic *versus* diabetic rats. This suggests that the amount of bone formed after resorption, which represents ‘‘coupling’’, a process where bone mass is maintained, is deficient in the diabetic animals and is illustrated in Fig. 5. Thus, diabetes may have more net bone loss because resorption is greater and coupling is less. Thus, this model is useful in investigating the mechanisms by which diabetes could aggravate periodontal tissue loss, including more persistent inflammation, greater bone resorption and decreased reparative bone formation leading to uncoupling and greater net bone loss.

Airpouch/Chamber Models

The airpouch model was originally developed in mice and rats by the Willoughby group to study the function of the synovial membrane (Edwards et al.

1981). To produce an airpouch, 5 cm³ of air is injected subcutaneously into the back of a mouse or a rat near the midline. A second injection is usually administered after 3 days to maintain the patency of the airpouch. After 6 days, the airpouch will have become filled with fluid and it is lined by epithelium. The model was used extensively in the rheumatoid arthritis (RA) literature to investigate the dynamics of synovial inflammation and the initiation of inflammation. Many of these studies were seminal in the identification of mast cells and histamine as important players in the early inflammatory response (Sin et al. 1986). The model also proved to be useful in developing clinical correlates for the role of cytokines, chemokines and lipid mediators in the pathogenesis of RA (Ferrandiz & Foster 1991).

More recently, the murine airpouch model has been adapted for the evaluation of potential pharmacologic agents. The injection of proinflammatory agents including irritants, cytokine or bacteria into the airpouch elicits a quantifiable inflammatory response measured as the number of inflammatory cells, secretion of cytokines or other common parameters (Sin et al. 1986). Hachicha et al. (1999) adapted Willoughby's model to assess neutrophil responses to TNF- α and IL-1 β and how the acute inflammatory response is modified by lipoxins and aspirin-triggered lipoxins. The model has the advantage of allowing both analysis of the exudate fluid and the epithelial lining of the airpouch. Lipoxins and their aspirin-triggered analogues induced a profound dampening of acute inflammation, demonstrating a dose–response reduction in the number of leucocytes in the exudate as well as a reduction of leucocytes in the epithelial lining. The experiment is performed in the viable whole animal, which reduces in vitro artefacts.

The airpouch model was adapted for periodontitis by Pouliot et al. (2000) to assess the acute inflammatory response to *P. gingivalis*. This report provides an excellent example of the benefit of whole-animal experiments to minimize in vitro artefacts. In previous work, Van Dyke et al. (1982) had reported that *P. gingivalis* produced a factor that inhibited neutrophil chemotaxis, prompting the suggestion that *P. gingivalis* was evading the host response by inhibiting its first line of defence. However, injection of *P. gingivalis* into the

Table 5. Hypotheses tested in the rat ligature model

Bacteria play an essential role in initiating gingival inflammation and periodontal bone loss	Rovin et al. (1966)
Chlorhexidine and antibiotics reduce bone resorption induced by oral bacteria	Kenworthy & Baverel (1981), Weiner et al. (1979)
A gram-negative bacterial burden enhances osteoclastogenesis and bone resorption	Samejima et al. (1990)
Endotoxin tolerance reduces periodontal bone loss	Samejima et al. (1990)
Cox-1 and -2 contribute to gingival inflammation, osteoclast formation and bone loss	Bezerra et al. (2000), Bezerra et al. (2002)
Application of interleukin 1 (IL-1) or TNF enhance periodontal bone loss	Gaspersic et al. (2003), Koide et al. (1995)
Cigarette smoke or nicotine enhances alveolar bone loss	César Neto et al. (2004), Benatti et al. (2003), Nociti et al. (2001)
Diabetes prolongs gingival inflammation, increases osteoclast activity and enhances fibroblast and osteoblast cell death	Al-Mashat et al. (2006)
Diabetes interferes with osseous coupling limiting reparative bone formation following periodontal bone resorption.	Al-Mashat et al. (2006)
Potential and future uses of this model	
To investigate systemic conditions that modulate loss of connective tissue attachment or alveolar bone	
To investigate the role of specific factors in the loss of attachment or alveolar bone by the use of well defined molecules including inhibitors	
To investigate mechanisms that affect coupling of alveolar bone resorption and formation	
mRNA or protein profiling of factors modulated during the progression of alveolar bone loss	




Fig. 5. Diabetes inhibits reparative alveolar bone formation following resorption. To induce periodontal bone loss ligatures were placed around the second molar teeth of type 2 diabetic (ZDF) and normoglycemic matched littermates. After 7 days ligatures were removed which allowed coupling to proceed involving the formation of new bone. The amount of bone formed following resorption was significantly less in the diabetic group suggesting that net bone loss may be greater in the diabetic group due to the failure to form new alveolar bone following an episode or resorption. *Significant difference between diabetic and normoglycemic control rats ($p < 0.05$). This figure was previously published in Liu et al. (2006b). ZDF, Zucker diabetic fatty.

airpouch induced a profound neutrophil infiltrate. In retrospect, it is likely that gingipains were degrading either the chemotactic agent or the neutrophil receptor for the agent, but importantly, it is now clear that this does not appear to occur in situ and *P. gingivalis* is apparently a strong pro-inflammatory stimulus. Another important observation relevant to the pathogenesis of periodontitis was also uncovered in this model. The major source of PGE₂ in periodontal tissues has been widely held to originate from monocyte/macrophages; these experiments demonstrated that neutrophils are a rich source of PGE₂ in the inflammatory lesion.

The shortcoming of the airpouch model is the duration of the experiment; it is not practical to conduct a chronic inflammation experiment with the airpouch, with maintenance of the airpouch being the limiting factor. This problem was solved in part by the use of a tissue cage and chamber models. The use of the airpouch and chamber models to test specific hypotheses is described in Table 6, along with potential future uses of this model.

The mouse subcutaneous chamber model was adapted for use in perio-

Table 6. Hypotheses tested in airpouch/chamber models

Eicosanoids control TNF- α expression in acute inflammation	Ferrandiz & Foster (1991)
Lipoxins and ATL regulate chemokines and cytokines and damped acute inflammation	Hachicha et al. (1999)
Lipoxin analogues dampen the neutrophil response to <i>Porphyromonas gingivalis</i> . Neutrophils are a rich source of PGE ₂	Pouliot et al. (2000)
<i>P. gingivalis</i> induces an acquired immune response	Genco et al. (1991)
Serologic diversity of <i>Neisseria</i> can provide protective immunity	Wong et al. (1979)
<i>P. gingivalis</i> evades phagocyte clearance by failing to activate or degrading complement	Schenkein (1989)
Immunization with <i>P. gingivalis</i> protects against disseminating infection	Dahlen & Slots (1989), Genco et al. (1992)
<i>P. gingivalis</i> activates blood kinin pathways to escape the vasculature creating a disseminating infection	Hu et al. (2006)
Potential and future uses of this model	
Determine the role of pharmacologic agents in acute and chronic inflammation	
Assess the role of putative virulence factors of single bacteria and mixed infection in acute and chronic inflammation	
Using transgenic animals, determine the role of specific host pathways, enzymes, receptors and ligands in the acute and chronic inflammatory response	

dontal studies by Genco et al. (1991). Originally used to study immunogenicity and strain diversity of *Neisseria gonorrhoeae* in mice (Wong et al. 1979), the method was also used to assess the pathogenesis of foreign-body infections in guinea-pigs (Zimmerli et al. 1982). Teflon tissue cages have also been used in guinea-pigs (Schenkein 1989) and New Zealand White Rabbits (Dahlen & Slots 1989). The chamber is a coiled stainless-steel wire that is surgically implanted subcutaneously in the back of the mouse (or other animal) and the wound is allowed to heal for 10 days. Like the airpouch model, the interior of the coil becomes epithelialized, allowing for the injection of bacteria or other substances into the lumen of the chamber. Fluid can be aspirated from the chamber at intervals or the entire chamber can be excised and histology can be performed.

The chamber model and the tissue cage variants used as periodontal models have, for the most part, assessed the virulence of bacteria. It has the advantage over abscess models of allowing assessment of host parameters as well as bacterial colonization, growth and tissue damage. Genco et al. (1991) were able to demonstrate the development of an acquired immune response to *P. gingivalis* injected into the chamber and to demonstrate strain differences within the bacteria. Dahlen and Slots (Zimmerli et al. 1982) obtained similar results with the tissue cage in rabbits. Schenkein (1989) evaluated the role of complement in the clearance of *P. gingivalis*.

The long-term chamber model was also effectively used in immunization studies (Genco et al. 1992). Mice were

immunized with non-invasive and invasive strains of heat-killed *P. gingivalis*, followed by a chamber challenge with invasive strains. The investigators were able to demonstrate that immunization with any strain provided protection against tissue damage and invasion by invasive (virulent) strains. Interestingly, immunization did not prevent colonization of the chamber by any *P. gingivalis* strain, but it did prevent tissue damage and spread of the colonizing bacteria. In another related experiment, Hu et al., demonstrated that activation of the kinin system is involved in the breach of the vascular barrier that allows for dissemination of *P. gingivalis*. The investigators demonstrated that *P. gingivalis* in the chamber caused leakage through the epithelial barrier and activation of the kinin system resulted in more severe dissemination, while blocking the kinin system prevented dissemination of *P. gingivalis*.

The strength of the airpouch and chamber models in small animals is clearly the whole-animal aspect for the measurement of host response parameters. The strength of the model is enhanced, particularly in mice, by the availability of genetically modified knock-out or transgenic animals that can be used in the experiments. For short-term, acute inflammation experiments, the strength of the airpouch model is its simplicity and ease of use with no requirement for surgical interventions. The major advantage of the chamber models is the ability to perform long-term experiments that can be adapted to immunization and chronic inflammation and pathogenesis experiments.

Concluding Remarks

Rodent models have provided a great deal of information about the process of periodontal disease. The rat ligature model using gnotobiotic rats demonstrated that bacteria play an essential role in initiating gingival inflammation and periodontal bone loss. The oral gavage and *A. actinomycetemcomitans* rat models have established that certain bacterial virulence factors are essential in promoting an infection that causes periodontal bone loss. That the acquired immune response contributes significantly to periodontal tissue destruction induced by transient *A. actinomycetemcomitans* or *P. gingivalis* infection has been demonstrated in the gavage model using mice that are genetically modified or modified by adoptive transfer of human lymphocytes. The mechanisms by which diabetes affects the response to *P. gingivalis* by prolonging inflammation, enhancing the death of fibroblasts and osteoblasts and by interfering with bone coupling have been shown in the calvarial model and in the ligature model of periodontal bone loss. Much of this critical insight into the mechanisms through which oral pathogens lead to connective tissue and bone loss could have been obtained in human studies and would have been considerably more difficult or expensive in other animal models. Moreover, in vivo models come closer to providing an accurate assessment of cell behaviour compared with in vitro models. For example, the response of host cells to bacteria is much more likely to be accurately depicted by in vivo models. This is true in spite of the fact that the in vivo setting in animals is not identical to the human periodontium. It is therefore the conclusion of the authors that well-designed animal models are essential to understanding the complex processes of periodontal diseases.

Acknowledgements

We would like to thank Alicia Ruff for help in preparing this manuscript.

References

- Adelman, N., Cohen, S. & Yoshida, T. (1978) Strain variations in murine MIF production. *Journal of Immunology* **121**, 209–212.
- Al-Mashat, H. A., Kandru, S., Liu, R., Behl, Y., Desta, T. & Graves, D. T. (2006) Diabetes enhances mRNA levels of proapoptotic genes and caspase activity, which contribute to impaired healing. *Diabetes* **55**, 487–495.
- Alayan, J., Ivanovski, S., Gemmell, E., Ford, P., Hamlet, S. & Farah, C. S. (2006) Deficiency of iNOS contributes to *Porphyromonas gingivalis*-induced tissue damage. *Oral Microbiology and Immunology* **21**, 360–365.
- Alikhani, M., Alikhani, Z., He, H., Liu, R., Popek, B. I. & Graves, D. T. (2003) Lipopolysaccharides indirectly stimulate apoptosis and global induction of apoptotic genes in fibroblasts. *Journal of Biological Chemistry* **278**, 52901–52908.
- Alnaeeli, M., Park, J., Mahamed, D., Penninger, J. M. & Teng, Y. T. (2007) Dendritic cells at the osteo-immune interface: implications for inflammation-induced bone loss. *Journal of Bone and Mineral Research* **22**, 775–780.
- Alnaeeli, M., Penninger, J. M. & Teng, Y. T. (2006) Immune interactions with CD4+ T cells promote the development of functional osteoclasts from murine CD11c+ dendritic cells. *Journal of Immunology* **177**, 3314–3326.
- Aoba, T. (2004) Solubility properties of human tooth mineral and pathogenesis of dental caries. *Oral Diseases* **10**, 249–257.
- Assuma, R., Oates, T., Cochran, D., Amar, S. & Graves, D. (1998) IL-1 and TNF antagonists inhibit the inflammatory response and bone loss in experimental periodontitis. *Journal of Immunology* **160**, 403–409.
- Baker, P., Dixon, M., Evans, R., Dufour, L., Johnson, E. & Roopenian, D. (1999) CD4(+) T cells and the proinflammatory cytokines gamma interferon and interleukin-6 contribute to alveolar bone loss in mice. *Infection and Immunity* **67**, 2804–2809.
- Baker, P., Evans, R. & Roopenian, D. (1994) Oral infection with *Porphyromonas gingivalis* and induced alveolar bone loss in immunocompetent and severe combined immunodeficient mice. *Archives of Oral Biology* **39**, 1035–1040.
- Baker, P. J. (2005) Genetic control of the immune response in pathogenesis. *Journal of Periodontology* **76**, 2042–2046.
- Baker, P. J., Dixon, M., Evans, R. T. & Roopenian, D. C. (2000a) Heterogeneity of *Porphyromonas gingivalis* strains in the induction of alveolar bone loss in mice. *Oral Microbiology and Immunology* **15**, 27–32.
- Baker, P. J., Dixon, M. & Roopenian, D. C. (2000b) Genetic control of susceptibility to *Porphyromonas gingivalis*-induced alveolar bone loss in mice. *Infection and Immunity* **68**, 5864–5868.
- Battaglino, R., Vokes, M., Schulze-Spate, U., Sharma, A., Graves, D., Kohler, T., Muller, R., Yoganathan, S. & Stashenko, P. (2007) Fluoxetine treatment increases trabecular bone formation in mice. *Journal of Cellular Biochemistry* **100**, 1387–1394.
- Benatti, B. B., Nogueira-Filho, G. R., Diniz, M. C., Sallum, E. A., Sallum, A. W. & Nociti, F. H. Jr. (2003) Stress may enhance nicotine effects on periodontal tissues. An in vivo study in rats. *Journal of Periodontal Research* **38**, 351–353.
- Bezerra, M. M., Brito, G. A., Ribeiro, R. A. & Rocha, F. A. (2002) Low-dose doxycycline prevents inflammatory bone resorption in rats. *Brazilian Journal of Medical and Biological Research* **35**, 613–616.
- Bezerra, M. M., de Lima, V., Alencar, V. B., Vieira, I. B., Brito, G. A., Ribeiro, R. A. & Rocha, F. A. (2000) Selective cyclooxygenase-2 inhibition prevents alveolar bone loss in experimental periodontitis in rats. *Journal of Periodontology* **71**, 1009–1014.
- Boyce, B., Aufdemorte, T., Garrett, I., Yates, A. & Mundy, G. (1989) Effects of interleukin-1 on bone turnover in normal mice. *Endocrinology* **125**, 1142–1150.
- Burckhardt, J. J. & Guggenheim, B. (1980) Increased smooth-surface caries incidence in gnotobiotic rats immunized with *Actinomyces viscosus*. *Caries Research* **14**, 56–59.
- Burns, E., Bachrach, G., Shapira, L. & Nussbaum, G. (2006) Cutting edge: TLR2 is required for the innate response to *Porphyromonas gingivalis*: activation leads to bacterial persistence and TLR2 deficiency attenuates induced alveolar bone resorption. *Journal of Immunology* **177**, 8296–8300.
- Caufield, P., Li, Y. & Dasanayake, A. (2005) Dental caries: an infectious and transmissible disease. *Compendium of Continuing Education in Dentistry* **26**, 10–16.
- César Neto, J. B., de Souza, A. P., Barbieri, D., Moreno, H. Jr., Sallum, E. A. & Nociti, F. H. Jr. (2004) Matrix metalloproteinase-2 may be involved with increased bone loss associated with experimental periodontitis and smoking: a study in rats. *Journal of Periodontology* **75**, 995–1000.
- Chang, K. M., Ramamurthy, N. S., McNamara, T. F., Genco, R. J. & Golub, L. M. (1988) Infection with a gram-negative organism stimulates gingival collagenase production in non-diabetic and diabetic germfree rats. *Journal of Periodontal Research* **23**, 239–244.
- Chiang, C., Kyritsis, G., Graves, D. & Amar, S. (1999) Interleukin-1 and tumor necrosis factor activities partially account for calvarial bone resorption induced by local injection of lipopolysaccharide. *Infection and Immunity* **67**, 4231–4236.
- Choi, J. I., Schifferle, R. E., Yoshimura, F. & Kim, B. W. (1998) Capsular polysaccharide-fimbrial protein conjugate vaccine protects against *Porphyromonas gingivalis* infection in SCID mice reconstituted with human peripheral blood lymphocytes. *Infection and Immunity* **66**, 391–393.
- Clark, J., Palmer, C. & Shaw, W. (1983) The diabetic Zucker fatty rat. *Proceedings of the Society for Experimental Biology and Medicine* **173**, 68–75.
- Cohen, M. E. & Ralls, S. A. (1988) Distributions of periodontal attachment levels. Mathematical models and implications. *Journal of Periodontology* **59**, 254–258.
- Daep, C. A., James, D. M., Lamont, R. J. & Demuth, D. R. (2006) Structural characterization of peptide-mediated inhibition of *Porphyromonas gingivalis* biofilm formation. *Infection and Immunity* **74**, 5756–5762.

- Dahlen, G. & Slots, J. (1989) Experimental infections by *Bacteroides gingivalis* in non-immunized and immunized rabbits. *Oral Microbiology and Immunology* **4**, 6–11.
- Darby, I., Bisucci, T., Hewitson, T. & MacLellan, D. (1997) Apoptosis is increased in a model of diabetes-impaired wound healing in genetically diabetic mice. *The International Journal of Biochemistry and Cell Biology* **29**, 191–200.
- Darville, T., Andrews, C. W. Jr., Laffoon, K. K., Shymasani, W., Kishen, L. R. & Rank, R. G. (1997) Mouse strain-dependent variation in the course and outcome of chlamydial genital tract infection is associated with differences in host response. *Infection and Immunity* **65**, 3065–3073.
- Delima, A., Spyros, K., Amar, S. & Graves, D. T. (2002) Inflammation and tissue loss caused by periodontal pathogens is reduced by IL-1 antagonists. *The Journal of Infectious Disease* **186**, 511–516.
- Edwards, J. C., Sedgwick, A. D. & Willoughby, D. A. (1981) The formation of a structure with the features of synovial lining by subcutaneous injection of air: an in vivo tissue culture system. *Journal of Pathology* **134**, 147–156.
- Ferrandiz, M. L. & Foster, S. J. (1991) Tumour necrosis factor production in a rat airpouch model of inflammation: role of eicosanoids. *Agents Actions* **32**, 289–294.
- Fine, D. H., Furgang, D., Schreiner, H. C., Goncharoff, P., Charlesworth, J., Ghazwan, G., Fitzgerald-Bocarsly, P. & Figurski, D. H. (1999) Phenotypic variation in *Actinobacillus actinomycetemcomitans* during laboratory growth: implications for virulence. *Microbiology* **145** (Pt 6), 1335–1347.
- Fine, D. H., Goncharoff, P., Schreiner, H., Chang, K. M., Furgang, D. & Figurski, D. (2001) Colonization and persistence of rough and smooth colony variants of *Actinobacillus actinomycetemcomitans* in the mouths of rats. *Archives of Oral Biology* **46**, 1065–1078.
- Fine, D. H., Kaplan, J. B., Kachlany, S. C. & Schreiner, H. C. (2006) How we got attached to *Actinobacillus actinomycetemcomitans*: a model for infectious diseases. *Periodontology* **2000** **42**, 114–157.
- Fine, D. H., Velliyagounder, K., Furgang, D. & Kaplan, J. B. (2005) The *Actinobacillus actinomycetemcomitans* autotransporter adhesin Aae exhibits specificity for buccal epithelial cells from humans and old world primates. *Infection and Immunity* **73**, 1947–1953.
- Finlay, B. B. & McFadden, G. (2006) Anti-immunology: evasion of the host immune system by bacterial and viral pathogens. *Cell* **124**, 767–782.
- Fitzgerald, R. J. & Keyes, P. H. (1960) Demonstration of the etiologic role of streptococci in experimental caries in the hamster. *The Journal of American Dental Association* **61**, 9–19.
- Garlet, G. P., Cardoso, C. R., Silva, T. A., Ferreira, B. R., Avila-Campos, M. J., Cunha, F. Q. & Silva, J. S. (2006) Cytokine pattern determines the progression of experimental periodontal disease induced by *Actinobacillus actinomycetemcomitans* through the modulation of MMPs, RANKL, and their physiological inhibitors. *Oral Microbiology and Immunology* **21**, 12–20.
- Gaspersic, R., Stiblar-Martincic, D., Osredkar, J. & Skaleric, U. (2003) Influence of subcutaneous administration of recombinant TNF-alpha on ligature-induced periodontitis in rats. *Journal of Periodontal Research* **38**, 198–203.
- Gemmell, E., Winning, T. A., Grieco, D. A., Bird, P. S. & Seymour, G. J. (2000) The influence of genetic variation on the splenic T cell cytokine and specific serum antibody responses to *Porphyromonas gingivalis* in mice. *Journal of Periodontology* **71**, 1130–1138.
- Genco, C. A., Cutler, C. W., Kapczynski, D., Maloney, K. & Arnold, R. R. (1991) A novel mouse model to study the virulence of and host response to *Porphyromonas (Bacteroides) gingivalis*. *Infection and Immunity* **59**, 1255–1263.
- Genco, C. A., Kapczynski, D. R., Cutler, C. W., Arko, R. J. & Arnold, R. R. (1992) Influence of immunization on *Porphyromonas gingivalis* colonization and invasion in the mouse chamber model. *Infection and Immunity* **60**, 1447–1454.
- Gibson, F. C. III & Genco, C. A. (2001) Prevention of *Porphyromonas gingivalis*-induced oral bone loss following immunization with gingipain R1. *Infection and Immunity* **69**, 7959–7963.
- Gibson, F. C. III, Hong, C., Chou, H. H., Yumoto, H., Chen, J., Lien, E., Wong, J. & Genco, C. A. (2004) Innate immune recognition of invasive bacteria accelerates atherosclerosis in apolipoprotein E-deficient mice. *Circulation* **109**, 2801–2806.
- Gilthorpe, M. S., Zamzuri, A. T., Griffiths, G. S., Maddick, I. H., Eaton, K. A. & Johnson, N. W. (2003) Unification of the ‘burst’ and ‘linear’ theories of periodontal disease progression: a multilevel manifestation of the same phenomenon. *Journal of Dental Research* **82**, 200–205.
- Graves, D. T., Liu, R., Alikhani, M., Al-Mashat, H. & Trackman, P. C. (2006) Diabetes-enhanced inflammation and apoptosis – impact on periodontal pathology. *Journal of Dental Research* **85**, 15–21.
- Graves, D. T., Naguib, G., Lu, H., Desta, T. & Amar, S. (2005a) *Porphyromonas gingivalis* fimbriae are pro-inflammatory but do not play a prominent role in the innate immune response to *P. gingivalis*. *Journal of Endotoxin Research* **11**, 13–18.
- Graves, D. T., Naguib, G., Lu, H., Leone, C., Hsue, H. & Krall, E. (2005b) Inflammation is more persistent in Type 1 diabetic mice. *Journal of Dental Research* **84**, 324–328.
- Graves, D., Oskoui, M., Volejnikova, S., Naguib, G., Cai, S., Desta, T., Kakouras, A. & Jiang, Y. (2001) Tumor necrosis factor modulates fibroblast apoptosis, PMN recruitment, and osteoclast formation in response to *P. gingivalis* infection. *Journal of Dental Research* **80**, 1875–1879.
- Gyorfi, A., Fazekas, A., Suba, Z., Ender, F. & Rosivall, L. (1994) Neurogenic component in ligature-induced periodontitis in the rat. *Journal of Clinical Periodontology* **21**, 601–605.
- Hachicha, M., Pouliot, M., Petasis, N. A. & Serhan, C. N. (1999) Lipoxin (LX)A4 and aspirin-triggered 15-epi-LXA4 inhibit tumor necrosis factor alpha-initiated neutrophil responses and trafficking: regulators of a cytokine-chemokine axis. *Journal of Experimental Medicine* **189**, 1923–1930.
- Hajishengallis, G., Ratti, P. & Harokopakis, E. (2005) Peptide mapping of bacterial fimbrial epitopes interacting with pattern recognition receptors. *Journal of Biological Chemistry* **280**, 38902–38913.
- Hajishengallis, G., Shakhatreh, M. A., Wang, M. & Liang, S. (2007) Complement receptor 3 blockade promotes IL-12-mediated clearance of *Porphyromonas gingivalis* and negates its virulence in vivo. *Journal of Immunology* **179**, 2359–2367.
- Hajishengallis, G., Wang, M., Harokopakis, E., Triantafyllou, M. & Triantafyllou, K. (2006) *Porphyromonas gingivalis* fimbriae proactively modulate beta2 integrin adhesive activity and promote binding to and internalization by macrophages. *Infection and Immunity* **74**, 5658–5666.
- Han, X., Kawai, T., Eastcott, J. W. & Taubman, M. A. (2006) Bacterial-responsive B lymphocytes induce periodontal bone resorption. *Journal of Immunology* **176**, 625–631.
- Hardham, J., Reed, M., Wong, J., King, K., Laurinat, B., Sfintescu, C. & Evans, R. T. (2005) Evaluation of a monovalent companion animal periodontal disease vaccine in an experimental mouse periodontitis model. *Vaccine* **23**, 3148–3156.
- Harokopakis, E. & Hajishengallis, G. (2005) Integrin activation by bacterial fimbriae through a pathway involving CD14, Toll-like receptor 2, and phosphatidylinositol-3-kinase. *European Journal of Immunology* **35**, 1201–1210.
- Hart, G. T., Shaffer, D. J., Akilesh, S., Brown, A. C., Moran, L., Roopenian, D. C. & Baker, P. J. (2004) Quantitative gene expression profiling implicates genes for susceptibility and resistance to alveolar bone loss. *Infection and Immunity* **72**, 4471–4479.
- He, H., Liu, R., Desta, T., Leone, C., Gerstenfeld, L. & Graves, D. (2004) Diabetes causes decreased osteoclastogenesis, reduced bone formation, and enhanced apoptosis of osteoblastic cells in bacteria stimulated bone loss. *Endocrinology* **145**, 447–452.
- Hillman, J. D., Brooks, T. A., Michalek, S. M., Harmon, C. C., Snoep, J. L. & van Der Weijden, C. C. (2000) Construction and characterization of an effector strain of *Streptococcus mutans* for replacement therapy of dental caries. *Infection and Immunity* **68**, 543–549.
- Holt, S. C. & Ebersole, J. L. (2005) *Porphyromonas gingivalis*, *Treponema denticola*, and *Tannerella forsythia*: the ‘red complex’, a prototype polybacterial pathogenic consortium in periodontitis. *Periodontology* **2000** **38**, 72–122.

- Hozumi, N., Gorczynski, R., Peters, W. & Sandhu, J. S. (1994) A SCID mouse model for human immune response and disease. *Research in Immunology* **145**, 370–379.
- Hu, S. W., Huang, C. H., Huang, H. C., Lai, Y. Y. & Lin, Y. Y. (2006) Transvascular dissemination of *Porphyromonas gingivalis* from a sequestered site is dependent upon activation of the kallikrein/kinin pathway. *Journal of Periodontal Research* **41**, 200–207.
- Jeon, S., Kim, S., Kim, E., Lee, J. E., Kim, S. J., Juhn, Y. S., Kim, Y. S., Bae, C. D. & Park, J. (2005) Chloride conductance is required for the protein kinase A and Rac1-dependent phosphorylation of moesin at Thr-558 by KCl in PC12 cells. *Journal of Biological Chemistry* **280**, 12181–12189.
- Kachlany, S. C., Planet, P. J., DeSalle, R., Fine, D. H. & Figurski, D. H. (2001) Genes for tight adherence of *Actinobacillus actinomycetemcomitans*: from plaque to plague to pond scum. *Trends in Microbiology* **9**, 429–437.
- Kenworthy, R. & Baverel, M. (1981) Studies of a periodontal tissue lesion in the rat, untreated or treated with chlorhexidine digluconate. *Journal of Clinical Periodontology* **8**, 349–358.
- Kesavalu, L., Chandrasekar, B. & Ebersole, J. L. (2002) In vivo induction of proinflammatory cytokines in mouse tissue by *Porphyromonas gingivalis* and *Actinobacillus actinomycetemcomitans*. *Oral Microbiology and Immunology* **17**, 177–180.
- Keyes, P. H. (1959) Dental caries in the Syrian hamster. VIII. The induction of rampant caries activity in albino and golden animals. *Journal of Dental Research* **38**, 525–533.
- Keyes, P. H. (1960) The infectious and transmissible nature of experimental dental caries. Findings and implications. *Archives of Oral Biology* **1**, 304–320.
- Klausen, B., Evans, R. T., Ramamurthy, N. S., Golub, L. M., Sfintescu, C., Lee, J. Y., Bedi, G., Zambon, J. J. & Genco, R. J. (1991) Periodontal bone level and gingival proteinase activity in gnotobiotic rats immunized with *Bacteroides gingivalis*. *Oral Microbiology and Immunology* **6**, 193–201.
- Koide, M., Suda, S., Saitoh, S., Ofuji, Y., Suzuki, T., Yoshie, H., Takai, M., Ono, Y., Taniguchi, Y. & Hara, K. (1995) In vivo administration of IL-1 beta accelerates silk ligature-induced alveolar bone resorption in rats. *Journal of Oral Pathology and Medicine* **24**, 420–434.
- Koulouri, O., Lapping, D., Radvar, M. & Kinane, D. (1999) Cell division, synthetic capacity and apoptosis in periodontal lesions analyzed by in situ hybridization and immunohistochemistry. *Journal of Clinical Periodontology* **26**, 552–559.
- Kuhr, A., Popa-Wagner, A., Schmoll, H., Schwahn, C. & Kocher, T. (2004) Observations on experimental marginal periodontitis in rats. *Journal of Periodontal Research* **39**, 101–106.
- Lalla, E., Lamster, I., Feit, M., Huang, L. & Schmidt, A. (1998) A murine model of accelerated periodontal disease in diabetes. *Journal of Periodontal Research* **33**, 387–399.
- Lalla, E., Lamster, I. B., Feit, M., Huang, L., Spessot, A., Qu, W., Kislinger, T., Lu, Y., Stern, D. M. & Schmidt, A. M. (2000) Blockade of RAGE suppresses periodontitis-associated bone loss in diabetic mice. *The Journal of Clinical Investigation* **105**, 1117–1124.
- Lalla, E., Lamster, I. B., Hofmann, M. A., Bucciarelli, L., Jerud, A. P., Tucker, S., Lu, Y., Papapanou, P. N. & Schmidt, A. M. (2003) Oral infection with a periodontal pathogen accelerates early atherosclerosis in apolipoprotein E-null mice. *Arteriosclerosis Thrombosis and Vascular Biology* **23**, 1405–1411.
- Leone, C. W., Bokhadhor, H., Kuo, D., Desta, T., Yang, J., Siqueira, M. F., Amar, S. & Graves, D. T. (2006) Immunization enhances inflammation and tissue destruction in response to *Porphyromonas gingivalis*. *Infection and Immunity* **74**, 2286–2292.
- Li, H., Suzuki, J., Bayna, E., Zhang, F., Dalle, M., Clark, A., Engler, R. & Lew, W. (2002a) Lipopolysaccharide induces apoptosis in adult rat ventricular myocytes via cardiac AT(1) receptors. *American Journal of Physiology – Heart and Circulatory Physiology* **282**, H461–467.
- Li, L., Khansari, A., Shapira, L., Graves, D. & Amar, S. (2002b) Contribution of interleukin-11 and prostaglandin(s) in lipopolysaccharide-induced bone resorption in vivo. *Infection and Immunity* **70**, 3915–3922.
- Liu, R., Bal, H. S., Desta, T., Behl, Y. & Graves, D. T. (2006a) Tumor necrosis factor- α mediates diabetes-enhanced apoptosis of matrix-producing cells and impairs diabetic healing. *American Journal of Pathology* **168**, 757–764.
- Liu, R., Bal, H. S., Desta, T., Krothapalli, N., Alyassi, M., Luan, Q. & Graves, D. (2006b) Diabetes enhances periodontal bone loss through enhanced resorption and diminished bone formation. *Journal of Dental Research* **85**, 510–514.
- Liu, R., Desta, T., He, H. & Graves, D. (2004) Diabetes alters the response to bacteria by enhancing fibroblast apoptosis. *Endocrinology* **145**, 2997–3003.
- Lohinai, Z., Benedek, P., Feher, E., Gyorf, A., Rosivall, L., Fazekas, A., Salzman, A. L. & Szabo, C. (1998) Protective effects of mercaptoethylguanidine, a selective inhibitor of inducible nitric oxide synthase, in ligature-induced periodontitis in the rat. *British Journal of Pharmacology* **123**, 353–360.
- Mac Donald, J. B., Socransky, S. & Sawyer, S. (1959) A survey of the bacterial flora of the periodontium in the rice rat. *Archives Oral Biology* **1**, 1–17.
- Mahamed, D. A., Marleau, A., Alnaeeli, M., Singh, B., Zhang, X., Penninger, J. M. & Teng, Y. T. (2005) G(-) anaerobes-reactive CD4+ T-cells trigger RANKL-mediated enhanced alveolar bone loss in diabetic NOD mice. *Diabetes* **54**, 1477–1486.
- Mahan, M. J., Schlauch, J. M. & Mekalanos, J. J. (1993) Selection of bacterial virulence genes that are specifically induced in host tissues. *Science* **259**, 686–688.
- Melkus, M. W., Estes, J. D., Padgett-Thomas, A., Gatlin, J., Denton, P. W., Othieno, F. A., Wege, A. K., Haase, A. T. & Garcia, J. V. (2006) Humanized mice mount specific adaptive and innate immune responses to EBV and TSST-1. *Nature Medicine* **12**, 1316–1322.
- Mosier, D. E., Gulizia, R. J., Baird, S. M. & Wilson, D. B. (1988) Transfer of a functional human immune system to mice with severe combined immunodeficiency. *Nature* **335**, 256–259.
- Mosier, D. E., Gulizia, R. J., Baird, S. M., Wilson, D. B., Spector, D. H. & Spector, S. A. (1991) Human immunodeficiency virus infection of human-PBL-SCID mice. *Science* **251**, 791–794.
- Naguib, G., Al-Mashat, H., Desta, T. & Graves, D. (2004) Diabetes prolongs the inflammatory response to a bacterial stimulus through cytokine dysregulation. *Journal of Investigative Dermatology* **123**, 87–92.
- Nociti, F. H. Jr., Nogueira-Filho, G. R., Tramontina, V. A., Machado, M. A., Barros, S. P., Sallum, E. A. & Sallum, A. W. (2001) Histometric evaluation of the effect of nicotine administration on periodontal breakdown: an in vivo study. *Journal of Periodontal Research* **36**, 361–366.
- Nowotny, A. & Sanavi, F. (1983) Induction of nonspecific tolerance to endotoxins reduces the alveolar bone resorption in ligature-treated rats. *Infection and Immunity* **39**, 873–878.
- O'Brien-Simpson, N. M., Pathirana, R. D., Paolini, R. A., Chen, Y. Y., Veith, P. D., Tam, V., Ally, N., Pike, R. N. & Reynolds, E. C. (2005) An immune response directed to proteinase and adhesin functional epitopes protects against *Porphyromonas gingivalis*-induced periodontal bone loss. *Journal of Immunology* **175**, 3980–3989.
- Offenbacher, S., Williams, R., Jeffcoat, M., Howell, T., Odle, B., Smith, M., Hall, C., Johnson, H. & Goldhaber, P. (1992) Effects of NSAID on beagle crevicular cyclooxygenase metabolites and periodontal bone loss. *Journal of Periodontal Research* **27**, 207–213.
- Paster, B. J., Olsen, I., Aas, J. A. & Dewhirst, F. E. (2006) The breadth of bacterial diversity in the human periodontal pocket and other oral sites. *Periodontology 2000* **42**, 80–87.
- Pathirana, R. D., O'Brien-Simpson, N. M., Brammar, G. C., Slakeski, N. & Reynolds, E. C. (2007) Kgp and RgpB, but not RgpA, are important for *Porphyromonas gingivalis* virulence in the murine periodontitis model. *Infection and Immunity* **75**, 1436–1442.
- Peterson, R. (1995) The Zucker Diabetic Fatty (ZDF) rat. In: Shafirir, E. (ed). *Advances in the Research of Diabetic Animals*, pp. 1–7. London: Smith-Gordon.
- Pihlstrom, B. L., Michalowicz, B. S. & Johnson, N. W. (2005) Periodontal diseases. *Lancet* **366**, 1809–1820.

- Pouliot, M., Clish, C. B., Petais, N. A., Van Dyke, T. E. & Serhan, C. N. (2000) Lipoxin A(4) analogues inhibit leukocyte recruitment to *Porphyromonas gingivalis*: a role for cyclooxygenase-2 and lipoxins in periodontal disease. *Biochemistry* **39**, 4761–4768.
- Rohane, P. W., Shimada, A., Kim, D. T., Edwards, C. T., Charlton, B., Shultz, L. D. & Fathman, C. G. (1995) Islet-infiltrating lymphocytes from prediabetic NOD mice rapidly transfer diabetes to NOD-scid/scid mice. *Diabetes* **44**, 550–554.
- Rovin, S., Costich, E. R. & Gordon, H. A. (1966) The influence of bacteria and irritation in the initiation of periodontal disease in germfree and conventional rats. *Journal of Periodontal Research* **1**, 193–204.
- Sakamoto, M., Umeda, M. & Benno, Y. (2005) Molecular analysis of human oral microbiota. *Journal of Periodontal Research* **40**, 277–285.
- Samejima, Y., Ebisu, S. & Okada, H. (1990) Effect of infection with *Eikenella corrodens* on the progression of ligature-induced periodontitis in rats. *Journal of Periodontal Research* **25**, 308–315.
- Sandhu, J., Shpitz, B., Gallinger, S. & Hozumi, N. (1994) Human primary immune response in SCID mice engrafted with human peripheral blood lymphocytes. *Journal of Immunology* **152**, 3806–3813.
- Sasaki, H., Hou, L., Belani, A., Wang, C.Y., Uchiyama, T., Müller, R. & Stashenko, P. (2000) IL-10, but not IL-4, suppresses infection-stimulated bone resorption in vivo. *Journal of Immunology* **165**, 3626–3630.
- Scannapieco, F. A., Millar, S. J., Reynolds, H. S., Zambon, J. J. & Levine, M. J. (1987) Effect of anaerobiosis on the surface ultrastructure and surface proteins of *Actinobacillus actinomycetemcomitans* (*Haemophilus actinomycetemcomitans*). *Infection and Immunity* **55**, 2320–2323.
- Schenkein, H. A. (1989) Failure of *Bacteroides gingivalis* W83 to accumulate bound C3 following opsonization with serum. *Journal of Periodontal Research* **24**, 20–27.
- Schreiner, H. C., Sinatra, K., Kaplan, J. B., Furgang, D., Kachlany, S. C., Planet, P. J., Perez, B. A., Figurski, D. H. & Fine, D. H. (2003) Tight-adherence genes of *Actinobacillus actinomycetemcomitans* are required for virulence in a rat model. *Proceedings of the National Academy of Sciences of USA* **100**, 7295–7300.
- Senpuku, H., Nakai, M., Koga, T., Hanada, N. & Nisizawa, T. (1996) Identification of a repeated epitope recognized by human serum antibodies in a surface protein antigen of *Streptococcus mutans*. *Oral Microbiology and Immunology* **11**, 121–128.
- Sharma, A., Honma, K., Evans, R. T., Hruby, D. E. & Genco, R. J. (2001) Oral immunization with recombinant *Streptococcus gordonii* expressing *porphyromonas gingivalis* FimA domains. *Infection and Immunity* **69**, 2928–2934.
- Sharma, A., Inagaki, S., Honma, K., Sfintescu, C., Baker, P. J. & Evans, R. T. (2005) *Tannerella forsythia*-induced alveolar bone loss in mice involves leucine-rich-repeat BspA protein. *Journal of Dental Research* **84**, 462–467.
- Shaw, J. H., Griffiths, D. K. & Auskaps, A. M. (1961) The influence of antibiotics on the periodontal syndrome in the rice rat. *Journal of Dental Research* **40**, 511–519.
- Shenker, B. J., Wannberg, S., Vitale, L., Kieba, I. & Lally, E. T. (1993) Reconstruction of severe combined immunodeficient mice with lymphocytes from patients with localized juvenile periodontitis. *Journal of Periodontal Research* **28**, 487–490.
- Shpitz, B., Chambers, C. A., Singhal, A. B., Hozumi, N., Fernandes, B. J., Roifman, C. M., Weiner, L. M., Roder, J. C. & Gallinger, S. (1994) High level functional engraftment of severe combined immunodeficient mice with human peripheral blood lymphocytes following pretreatment with radiation and anti-asialo GM1. *Journal of Immunological Methods* **169**, 1–15.
- Shultz, L. D., Ishikawa, F. & Greiner, D. L. (2007) Humanized mice in translational biomedical research. *Nature Reviews Immunology* **7**, 118–130.
- Shultz, L. D., Schweitzer, P. A., Christianson, S. W., Gott, B., Schweitzer, I. B., Tennent, B., McKenna, S., Mobraaten, L., Rajan, T. V., Greiner, D. L. et al. (1995) Multiple defects in innate and adaptive immunologic function in NOD/LtSz-scid mice. *Journal of Immunology* **154**, 180–191.
- Simpson, W. & Simmons, D. J. C. (1980) Two *Actinobacillus* species isolated from laboratory rodents. *Lab Animals* **14**, 15–16.
- Sin, Y. M., Sedgwick, A. D., Chea, E. P. & Willoughby, D. A. (1986) Mast cells in newly formed lining tissue during acute inflammation: a six day air pouch model in the mouse. *Annals of the Rheumatic Diseases* **45**, 873–877.
- Socransky, S. S., Haffajee, A. D., Goodson, J. M. & Lindhe, J. (1984) New concepts of destructive periodontal disease. *Journal of Clinical Periodontology* **11**, 21–32.
- Teng, Y. (2002a) Mixed periodontal Th1-Th2 cytokine profile in *Actinobacillus actinomycetemcomitans*-specific osteoprotegerin ligand (or RANK-L)-mediated alveolar bone destruction in vivo. *Infection and Immunity* **70**, 5269–5273.
- Teng, Y. (2003) The role of acquired immunity and periodontal disease progression. *Critical Reviews in Oral Biology and Medicine* **14**, 237–252.
- Teng, Y., Nguyen, H., Gao, X., Kong, Y., Gorczynski, R., Singh, B., Ellen, R. & Penninger, J. (2000) Functional human T-cell immunity and osteoprotegerin ligand control alveolar bone destruction in periodontal infection. *Journal of Clinical Investigations* **106**, R59–R67.
- Teng, Y. T. (2002b) Mixed periodontal Th1-Th2 cytokine profile in *Actinobacillus actinomycetemcomitans*-specific osteoprotegerin ligand (or RANK-L)-mediated alveolar bone destruction in vivo. *Infection and Immunity* **70**, 5269–5273.
- Teng, Y. T. & Hu, W. (2003) Expression cloning of a periodontitis-associated apoptotic effector, cagE homologue, in *Actinobacillus actinomycetemcomitans*. *Biochemical and Biophysical Research Communications* **303**, 1086–1094.
- Teng, Y. T., Mahamed, D. & Singh, B. (2005) Gamma interferon positively modulates *Actinobacillus actinomycetemcomitans*-specific RANKL+ CD4+ Th-cell-mediated alveolar bone destruction in vivo. *Infection and Immunity* **73**, 3453–3461.
- Teng, Y. T., Nguyen, H., Hassanloo, A., Ellen, R. P., Hozumi, N. & Gorczynski, R. M. (1999) Periodontal immune responses of human lymphocytes in *Actinobacillus actinomycetemcomitans*-inoculated NOD/SCID mice engrafted with peripheral blood leukocytes of periodontitis patients. *Journal of Periodontal Research* **34**, 54–61.
- Teng, Y. T. & Zhang, X. (2005) Apoptotic activity and sub-cellular localization of a T4SS-associated CagE-homologue in *Actinobacillus actinomycetemcomitans*. *Microbial Pathogenesis* **38**, 125–132.
- Trinchieri, G. (2003) Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nature Reviews Immunology* **3**, 133–146.
- Van Dyke, T. E., Bartholomew, E., Genco, R. J., Slots, J. & Levine, M. J. (1982) Inhibition of neutrophil chemotaxis by soluble bacterial products. *Journal of Periodontology* **53**, 502–508.
- Wang, M., Shakhathreh, M. A., James, D., Liang, S., Nishiyama, S., Yoshimura, F., Demuth, D. R. & Hajishengallis, G. (2007) Fimbrial proteins of *porphyromonas gingivalis* mediate in vivo virulence and exploit TLR2 and complement receptor 3 to persist in macrophages. *Journal of Immunology* **179**, 2349–2358.
- Weiner, G. S., DeMarco, T. J. & Bissada, N. F. (1979) Long term effect of systemic tetracycline administration on the severity of induced periodontitis in the rat. *Journal of Periodontology* **50**, 619–623.
- Wilensky, A., Gabet, Y., Yumoto, H., Hourihaddad, Y. & Shapira, L. (2005) Three-dimensional quantification of alveolar bone loss in *Porphyromonas gingivalis*-infected mice using micro-computed tomography. *Journal of Periodontology* **76**, 1282–1286.
- Williams, R., Jeffcoat, M., Kaplan, M., Goldhaber, P., Johnson, H. & Wechter, W. (1985) Flurbiprofen: a potent inhibitor of alveolar bone resorption in beagles. *Science* **227**, 640–642.
- Wong, K. H., Arko, R. J., Schalla, W. O. & Steurer, F. J. (1979) Immunological and serological diversity of *Neisseria gonorrhoeae*: identification of new immunotypes and highly protective strains. *Infection and Immunity* **23**, 717–722.
- Yu, J. J., Ruddy, M. J., Wong, G. C., Sfintescu, C., Baker, P. J., Smith, J. B., Evans, R. T. & Gaffen, S. L. (2007) An essential role for IL-17 in preventing pathogen-initiated bone destruction: recruitment of neutrophils to

- inflamed bone requires IL-17 receptor-dependent signals. *Blood* **109**, 3794–3802.
- Zappa, U., Reinking-Zappa, M., Graf, H. & Chase, D. (1992) Cell populations associated with active probing attachment loss. *Journal of Periodontology* **63**, 748–752.
- Zhang, X. & Teng, Y. T. (2006) Interleukin-10 inhibits gram-negative-microbe-specific human receptor activator of NF-kappaB ligand-positive CD4⁺-Th1-cell-associated alveolar bone loss in vivo. *Infection and Immunity* **74**, 4927–4931.
- Zhou, Q., Desta, T., Fenton, M., Graves, D. T. & Amar, S. (2005) Cytokine profiling of macrophages exposed to *Porphyromonas gingivalis*, its lipopolysaccharide, or its FimA protein. *Infection and Immunity* **73**, 935–943.
- Zimmerli, W., Waldvogel, F. A., Vaudaux, P. & Nydegger, U. E. (1982) Pathogenesis of foreign body infection: description and characteristics of an animal model. *Journal of Infectious Disease* **146**, 487–497.
- Zubery, Y., Dunstan, C., Story, B., Kesavalu, L., Ebersole, J., Holt, S. & Boyce, B. (1998) Bone resorption caused by three periodontal pathogens in vivo in mice is mediated in part by prostaglandin. *Infection and Immunity* **66**, 4158–4162.

Address:
 Dana T. Graves
 Department of Periodontology and
 Oral Biology
 Boston University School of Dental Medicine
 700 Albany Street, W-202D
 Boston, MA 02118
 USA
 E-mail: dgraves@bu.edu

Clinical Relevance

Scientific rationale for the study: Animal models are widely used to establish cause and effect relationships. However, no animal model recapitulates all aspects of disease initiation and progression. The purpose of this review is to illustrate hypotheses that can be addressed in different rodent models to provide an insight into the processes that occur in human periodontal diseases.

Principal findings: Different rodent models are well suited to address specific questions that are important in understanding host–bacteria interactions. Insights can be applied to design studies using animal models with specific end points in mind so

that the strength of a particular model is utilized. Thus, more than one model may be needed to provide a better understanding of the different steps in host/microbial interactions that lead to pathogenic developments. The important steps that may be investigated include the successful colonization of a susceptible host, bacterial invasion into vulnerable tissue, host–bacterial interactions that lead to an inflammatory response that may ultimately stimulate tissue destruction and finally a repair process that may mitigate some of the tissue damage that has occurred. With this approach in mind, animal models can provide data that would be impossible to obtain from humans

because in an animal model a particular process can be examined at many points in time with interventions that establish cause and effect relationships. In this manner, a better understanding of discrete steps in disease initiation or progression can be obtained.

Practical implications: Animal models are particularly well suited for studying cause and effect relationships and for initial studies of the safety and efficacy of new treatments. A better understanding of animal models will allow a more judicious match of a model with the hypothesis or therapy to be tested.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.