# Perceptions of Hispanic American Schoolchildren to the Risk Factors for Oral Cancer

Kishore Shetty, DDS, MS, MRCS John Brown BDS, PhD

## ABSTRACT

**Purpose:** The purpose of this study was to assess the knowledge of high schoolchildren on oral cancer risk factors in southern Texas.

**Methods:** A 39-item, self-administered questionnaire was completed by 1,667 students (14-18 years) in 4 high schools chosen by stratified cluster sampling.

**Results:** The effective response rate was 67%. Ninety-five percent of the students were Hispanic American, 55% had not heard about oral cancer, and 60% failed to identify the 2 most common early signs of oral cancer.

**Conclusions:** Lack of awareness of oral cancer risk factors indicate the need for improved health education directed towards minority population, especially in underprivileged areas. (J Dent Child 2007;74:62-6)

KEYWORDS: HISPANIC, ORAL CANCER, HEALTH EDUCATION

A nnually, more than 30,000 people in the United States are diagnosed with oral cancers. Oral cancer alone is responsible for more than 8,000 deaths each year, more than cervical cancer or malignant melanoma.<sup>1,2</sup> The number of deaths may be even higher, as oral cancer is often masked by other, more conspicuous comorbid conditions.<sup>3</sup> If detected early, the prognosis for survival of oral cancer is better than most cancers.<sup>4</sup> Oral cancer is a disease with: (1) known high-risk factors; (2) an asymptomatic phase with identifiable clinical features; (3) an available and efficient screening modality; and (4) effective nondeforming treatment for early lesions.<sup>5-7</sup> Most oral cancer lesions are not diagnosed, however, until they are in an advanced stage.<sup>8,9</sup>

Unlike research of other cancers, few studies have been conducted on the US public regarding oral cancer knowledge, opinions, and practices.<sup>10</sup> Little published information exists on young people's knowledge and attitudes about oral cancer and cancer prevention, especially in the Hispanic American population. The US Department of Health and Human Services (DHHS) has designated southern Texas as a "medically and dentally underserved" area. Such designation means many residents of that particular area lack access to basic health care needs. A report by the Texas Hispanic Information Initiative for Good Health<sup>11</sup> stated that southern Texas residents have important barriers to health care, such as: (1) urban isolation; (2) rural distances; (3) poor public health; and (4) low use of cancer screening services.

One such area in southern Texas is Brownsville, with a population greater than 115,000. Brownsville lies at the mouth of the Rio Grande River on the Gulf of Mexico, it is the most southerly city in the United States, with a highly mobile youthful population. Its twin border city in Mexico is Matamoros. Nearly 40 percent of the population lives below the federal poverty level. Historically, there has been a distinct need of oral health services among Brownsville schoolchildren.<sup>12</sup>

### **METHODS**

The sample population was chosen from the public school system in Brownsville, Texas, which represents a crosssection of different socioeconomic groups. Private and parochial schools enroll only 4% of the total ninth- to 12<sup>th</sup>grade population and were not considered for the study. The authors employed a 2-stage cluster sample design to produce a simple convenience representative sample of students in grades 9 to 12. The first stage-sampling frame

Dr. Shetty is Director, Medically Complex Patient Clinic, University of Texas Health Sciences Center, Houston, TX, and Dr. Brown is Professor, Department of Community Dentistry, University of Texas Health Sciences Center, San Antonio, TX. Correspond with Dr. Shetty at Kishore.Shetty@uth.tmc.edu

consisted of all public schools (Hanna, Simon Rivera, Porter, Lopez and Pace) containing any of grades 9 to 12. Four of the 5 high schools agreed to participate in the study (Hanna, Simon Rivera, Porter and Lopez). At the second sampling stage at each of the 4 schools, intact classes on a selected required subject or a selected period (eg, second period) were randomly selected. All students in the selected classes were eligible to participate in the survey.

The National Health Interview Survey on Teenage Attitudes and Practices and Youth Risk Behavior Surveillance System Questionnaire was used in designing the study questionnaire.<sup>13</sup> Precoded questions relating to the following topics were included: risk factors specific to oral cancer, and early signs and risk sites for oral cancer.

The questions were designed and selected to be appropriate, self explanatory, and easy to answer with little or no assistance. Survey procedures were designed to protect the student's privacy by allowing for anonymous and voluntary participation. Teachers administered the survey, which took approximately 15 minutes to complete. Students recorded their responses directly on the questionnaire. Parental consent procedures were followed before administering the questionnaire, according to the IRB protocol. A positive parental consent was taken from all students who wished to participate in the survey.

## RESULTS

The sampling frame consisted of 2,600 children to whom consent forms and questionnaires were distributed. A total of 1,754 questionnaires were returned (response rate=67%). Eighty seven questionnaires were: incomplete, unanswered, and not considered for data analysis. Thus, results computed are from the 1,667 completed questionnaires (Table 1). Hispanic Americans comprised 95% of the sample population. The sample was evenly distributed regarding gender, with 51% (N=856) males and 49% (N=811) females. The overwhelming majority of Hispanic Americans in Brownsville are of Mexican Americans, and in this report these terms may be used interchangeably.

More than half of the respondents (55%) had not heard about oral cancer (Figure 1). Tobacco use was the only risk factor, which was correctly identified by most of the respondents. 53% did not consider alcohol to be a risk factor. Only 11% correctly responded that sunlight could be a risk factor for lip cancer. About a quarter of the study population incorrectly mentioned drinking coffee as a risk factor for oral cancer; 42% thought poor oral hygiene was a risk factor for oral cancer. Almost 80% of the students mentioned gums as a common risk site for oral cancer, which, in fact, is one of the least common sites. The same holds true for the roof of the mouth. About 20% of the respondents even mentioned teeth as a risk site for cancer.

The most common type of precancerous lesions in the United States occurs along or under the tongue. About 50% of the respondents did not mention tongue (or under the tongue) as a risk site for oral cancer (Figure 2). A white/red patch and a nonhealing ulcer are the most common signs of oral cancer. Nearly two thirds of the population thought a white or red patch in the mouth was not an early sign for oral cancer. About 55% of the respondents reported a nonhealing ulcer in the mouth as not an early risk sign for oral cancer (Figure 3). The fact that 60% of the population identified bleeding from the gums as an early sign of oral cancer suggests the lack of awareness and misconceptions regarding the disease. About 55% of the respondents reported a nonhealing ulcer in the mouth as not an early risk sign for oral cancer.

#### Table 1. Demographics of the Brownsville Oral Cancer Risk Behavior Survey Characteristics Number Percent Sex Female 811 49 Male 856 51 Grade 9th Grade 690 42 10th Grade 421 25 11th Grade 328 20 12<sup>th</sup> Grade 213 13 Age 14 years 100 6 15 years 379 23 16 years 517 31 17 years 379 23 2.87 18 years 17 Race or ethnicity White 41 3 Black 4 1 Hispanic 1585 95 Asian or Pacific Islander 4 1 Native American 6 1 Other 12 1 School А 524 31 В 376 23 С 416 25 D 351 21



Figure 1. Knowledge and perception of oral cancer by gender (by percentage).



*Figure 2. Risk sites for oral cancer in the mouth (by percentage).* 



Figure 3. Early signs for oral cancer in the mouth

# DISCUSSION

This study was limited to a school-based sample and the findings may not generalize to adolescents who are not in school. The ability of students to complete the questionnaire varied. The number of questions left blank increased towards the end of the questionnaire. Some of the students may have found the questions too difficult or lost interest. It must be considered that the beliefs expressed by the participants in part may reflect: (1) their own personal general health and well-being; (2) current media hype; and (3) existing economic conditions. A few individuals had a tendency to mark multiple answers or report ambiguous responses to certain questions. The sample

had a preponderance of Hispanic American subjects (95%) reflecting the Brownsville high school students' community. These considerations necessitate a degree of caution in drawing general conclusions from the data. It is representative of the Brownsville high school population, however, and has implications for Mexican American population, especially those living in underprivileged areas.

This study's results suggest that the high school students are ill informed about the risk factors of oral cancer. Many of them hold misconceptions about the early signs and risk sites in the mouth that can have an impact on preventive behaviors. Across all groups, there was a higher level of knowledge about tobacco use as a risk factor. For decades, it has been known that the use of tobacco products and excessive alcohol is detrimental to health. The use of tobacco products, especially cigarettes, is harmful and a major contributing factor to mortality vis-à-vis heart disease, emphysema, lung cancer, and low birth-weight babies. A variety of educational and informational campaigns have urged tobacco users to stop. Rarely do available educational materials also include information on the use of tobacco products as risk factor for oral cancer. Similarly, there have been major educational efforts regarding alcohol use as a risk factor for cirrhosis of the liver, liver cancer, and fetal alcohol syndrome. However these educational messages seldom identify alcohol use as a risk factor for oral cancers.

There is a great need for more information about oral cancer and cancer prevention in the schools. More than 50% of this study's students wanted more information about oral cancer. Relatively few oral cancer educational materials for the public have been produced. This observation is particularly noteworthy, especially for dental professionals, when compared with the plethora of materials that have been developed on tooth-brushing, flossing, and the need for dental visits.

Studies are lacking that determine what educational materials are available and that assess their content in terms of accuracy, comprehensiveness, reading level, and cultural acceptability. A review of health education textbooks for kindergarten through 12th grade students found that the oral cancer coverage was uneven, misleading, sometimes incorrect; or (4) most often omitted altogether.<sup>14</sup> A lack of content and incorrect content in health education textbooks may contribute to the public's overall lack of knowledge about oral cancers. There is a great need for cancer prevention in the school curriculum. The content should include knowledge about cancer risk factors, particularly in the area of diet, where specific dietary advice is required.<sup>15-17</sup>

One objective in the Healthy People 2010 initiative is to reduce mortality from oral cancers.<sup>18</sup> As with most other cancers, when oral cancer is detected early the prognosis is greatly improved. Without accurate and appropriate information, people can neither make nor be expected to make informed, intelligent decisions about their own health.<sup>19</sup> Preventable diseases, such as oral cancer, impair quality of life and impose financial burdens on society. Many people do not practice preventive behaviors because they have never been taught about them and they do not have skills to seek such information or access to the information. Individuals need to know the risk factors for, signs and symptoms of oral cancers and how to obtain a thorough oral examination.<sup>20</sup> According to the 1995-96 Texas Oral Cancer Risk Behavior Survey, only 33% of Texan adults reported at least one risk behavior for oral cancer (tobacco use and risky alcohol behaviors).<sup>21</sup> Public awareness and education efforts should be increased with emphasis on the oral cancer risks associated with tobacco and alcohol use. Although oral cancer typically strikes older adults, interventions targeting younger persons are more likely to be successful and discourage long-term tobacco and alcohol exposure. Dental health care providers should focus on improving oral cancer awareness during patient treatments.

# **CONCLUSIONS**

Based on this study's results, the authors conclude that oral cancer prevention programs targeting the adolescent high school population should emphasize the providing of factual information about: (1) cancer; (2) its risk factors; and (3) screening. Future studies in this area should explore the role of acculturation and generational differences in predicting alcohol, tobacco, and other substance abuse.

# REFERENCES

1. Boring CC, Montgomery S, Squies TS, Tong T. Cancer statistics. CA Cancer J Clin 1994;44:7-26.

- 2. Blot WJ, McLaughlin JK, Winn DM, et al. Smoking and drinking in relation to oral and pharyngeal cancer. Cancer Res 1988;48:3282-7.
- 3. Farina A, Martinez C, Ramirez A. Hispanic health issues in Texas: A preliminary report. In: Texas Hispanic Information Initiative for Good Health (Texas HIIGH). South Texas Health Research Center and Center for Health Economics and Policy (CHEP). San Antonio, Tex: The University of Texas Health Science Center at San Antonio; 1994.
- 4. FDI Working Group. Nutrition, diet, and oral health. Int Dent J 1994;44:599-612.
- 5. Frazier PJ, Horowitz AM. Oral Health education and promotion in maternal and child health: A position paper. J Public Health Dent 1990;50:390-5.
- 6. Gold RS, Horowitz AM. Oral health information in textbooks [abstract]. Presented at the 121st Scientific Session of the American Public Health Association, Oct 26, 1993, San Francisco, Calif.
- Green LW, Kreuter MW. Health Promotion Planning: An Educational and Environmental Approach. 2nd ed. Mountain View, Calif: Mayfield Publishing Co; 1991.
- Hebert JR, Landon J, Miller DR. Consumption of meat and fruit in relation to oral esophageal cancer: A cross-national study. Nutr Cancer 1993;19:169-79.
- 9. Horowitz AM, Nourjah P, Gift HC. US Adult knowledge of risk factors and signs of oral cancers: 1990. J Am Dent Assoc 1995;126:39-44.
- 10. Mashberg A, Samit A. Early diagnosis of asymptomatic oral and oropharyngeal squamous cancers. CA Cancer J Clin 1995;45:328-51.
- 11. Mashberg A, Samit AM. Early detection, diagnosis, and management of oral and oropharyngeal cancer. Int J Oral Maxillofac Surg 1989;18:99-103.
- 12. Moore C. Education and oral cancer. In: Oral Cancer Interprofessional Symposium. Washington, DC: Government Printing Office; 1966:83-92. Public Health Service publication no.1806.
- National Cancer Institute. Cancer Statistics Review 1973-1987. Washington, DC: Government Printing Office, Department of Health and Human Services, Public Health Service, National Institute of Health; 1989. NIH publication no. 88-2789.
- Public Health Service. Chronic Disease and Health Promotion Reprints from the MMWR: 1990-1992. Youth Risk Behavior Surveillance System. Washington, DC: US Department of Health and Human Services; 1993.
- 15. Silverman S Jr., Shillitoe EJ. Etiology and predisposing factors. In: Silverman S Jr, ed. Oral Cancer. 3rd ed Atlanta, Ga: American Cancer Society; 1990:7-30.
- 16. Stalpers LJA, Van Vierzen PBJ, Bruns JJA, et al. The role of yearly chest radiography in the early detection of lung cancer following oral cancer. Int J Oral Maxillofac Surg 1989;18:99-103.

- 17. Texas Department of Health. Available at: "<u>http://</u> <u>www.dshs.state.tx.us/dental/default.shtm</u>". Accessed April 2007.
- Texas Risk Factor Report. Behavioral Risk Factor Surveillance System—Oral Cancer Risk Behaviors 1995-1996 Survey Data. Texas Dept. of Health, Bureau of Chronic Diseases. Available at: "http://www. dshs.state.tx.us/chs/brfss". Accessed April 2007.
- US Department of Health and Human Services, Public Health Service. Healthy People 2010: National Health Promotion and Disease Prevention Objectives. Washington, DC: DHHS publication no. (PHS) 91-50212. Available at: "http://www.cdc.gov/nchs/about/otheract/hpdata2010/abouthp.htm". Accessed April 2007.
- 20. Winn DM. Diet and nutrition in the etiology of oral cancer. Am J Clin Nutr 1995;61(suppl):437S-45S.

Copyright of Journal of Dentistry for Children is the property of American Academy of Pediatric Dentistry and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.