# Metachronous Bilateral Dentigerous Cysts Associated With Permanent First Molars

Eduardo Rodrigues Fregnani, DDS, PhD Danyel Elias da Cruz Perez, DDS, PhD Paulo André Gonçalves de Carvalho, DDS Fábio Abreu Alves, DDS, PhD

#### ABSTRACT

The occurrence of bilateral dentigerous cysts (DC) has been described as being associated mostly with third molars in adult patients. Very few reports show the bilateral occurrence of dentigerous cysts associated with first molars or premolars in childhood. The purpose of this report was to describe the case of a 5-year-old boy with bilateral mandibular dentigerous cysts associated with permanent first molars. (J Dent Child 2008;75:197-200)

Received March 3, 2007; Last Revision May 21, 2007; Revision Accepted May 23, 2007.

Keywords: Cysts odontogenic, cysts dentigerous, cysts, radicular, metachronous lesions

A ccording to the World Health Organization,<sup>1</sup> dentigerous cysts (DC) are developmental odontogenic cysts associated with the crowns of permanent teeth. Frequently, they do not cause symptoms or pain, but noneruption of the affected permanent tooth is a common clinical finding. A radiolucent area with well-defined sclerotic borders associated with the crown of an unerupted tooth is a typical radiographic finding.<sup>1</sup>

DCs are the second most common odontogenic cysts after radicular cysts and occur most frequently in the mandible during the second or third decades of life, mainly associated with impacted third molar teeth.<sup>2</sup> The occurrence of this lesion associated with primary teeth and permanent first molars is very rare,<sup>3</sup> and it is usually a single lesion. Bilateral DCs are an extremely uncommon finding in children. The purpose of this paper was to describe the case of metachronous bilateral dentigerous cysts in a nonsyndromic child patient.

### **CASE DESCRIPTION**

A 5-year-old boy was referred to the Stomatology Department at A.C. Camargo Cancer Hospital, Sao Paulo, Brazil, complaining of a painless lump of 25 days duration that appeared in the mandible's posterior region. His previous medical history was not contributory.

Upon extraoral exam, slight facial asymmetry was observed due to a firm, well-circumscribed nodule, which could be palpated in the left mandibular body's posterior region (Figure 1). No presence of palpable neck lymph nodes was noted. Intraoral examination revealed discoloration of



Figure 1. During clinical examination, a slight facial asymmetry was observed in the left mandibular body's posterior region.

Dr. Fregnani is a dental researcher, Dr. de Carvalho is a graduate student, and Dr. Alves is chief of department, all in the Department of Stomatology, A.C. Camargo Cancer Hospital, São Paulo, Brazil; Dr. Perez is Associated-Professor, Oral Pathology Division, University of Ribeirão Preto, São Paulo, and is a dental researcher at the Department of Stomatology, A.C. Camargo Cancer Hospital, São Paulo, Brazil. Correspond with Dr. Fregnani at erfreg@terra.com.br



Figure 2. Computed tomography showing a rounded lesion with liquid density sited on the mandible's left side, which expanded the vestibular bone plate and adjacent soft tissues.



Figure 3. Three-dimensional reconstruction confirmed a wellcircumscribed lesion associated with the permanent first molar's crown.



Figure 4. After 2 months, a clearly defined radiolucent lesion involving the permanent right first molar's crown was noted.

the bottom of the fold in the left posterior inferior region adjacent to the primary second molar's distal region, with noneruption of the inferior permanent first molar.

Panoramic radiography evidenced a unilocular lesion on the mandible's left side, with clear limits that dislocated the permanent second molar germ, reabsorbed the primary second molar's distal root, and involved the nonerupted permanent first molar's entire crown, measuring 2 cm in diameter. Computed tomography revealed a rounded lesion with liquid density sited on the mandible's left side, expanding the vestibular bone plate and adjacent soft tissues (Figure 2). Three-dimensional reconstruction using the cuts from computed tomography confirmed a bounded lesion associated with the permanent mandibular first molar's crown (Figure 3). During nitrous-oxide sedation to perform the computed tomography, aspiration punction was performed and 3 ml of liquid was withdrawn. According to clinical and image features, a DC was considered as the main diagnostic hypothesis. Thus, an incisional biopsy of the lesion and marsupialization (decompression) were performed simultaneously to allow eruption of the involved tooth. Microscopically, the diagnosis of DC was confirmed.

After 2 months of clinical and radiographic followup, the patient complained of a swelling on the right side of his face. On clinical examination, the presence of a nodule located in the right mandibular body's posterior region was observed. Radiographically, a well-circumscribed radiolucent lesion 2 cm in extent was noted involving the permanent right first molar's crown (Figure 4). On the left side, there was still a residual lesion associated with the permanent first molar's crown, similar to that observed on the right side. Thus, under general anesthesia, the patient was submitted to enucleation of the bilateral cystic lesions, maintaining the involved permanent teeth. Histopathologically in

both lesions, cystic cavities lined by cubic epithelium were observed, confirming the diagnosis of bilateral DCs. Clinical re-evaluation showed no associated syndromes.

After 15 days, it was clinically possible to note the initial eruption of both first molars. After 24 months, complete eruption could be observed, without clinical and radio-graphic signs of recurrence (Figure 5).

#### DISCUSSION

DCs usually occur singly, and are mainly associated with impacted permanent mandibular third molar teeth. Moreover, they occur mainly in patients during the second and third decades of life.<sup>1</sup> Their bilateral presentation is considered rare and can be sometimes associated with syndromes, such as cleidocranial dysplasia and Maroteaux-Lamy syndrome.<sup>4</sup> De Biase et al<sup>5</sup> reported a case of bilateral mandibular cysts in an 8-year-old boy treated with cyclosporin (CsA) after



Figure 5. Clinical and radiographic follow–up showing complete eruption of the mandibular first molars without signs of recurrence.

renal transplantation. The genesis of mandibular cysts might be associated with the combined post-transplantation use of CsA and a calcium channel blocker. The occurrence of DCs in childhood can also be considered uncommon.<sup>6</sup> In the present case, the patient was 5 years old, and there were no associated syndromes.

Most previous articles reporting bilateral DCs were cases

associated with impacted third molars. To our knowledge, there are only about 24 cases of bilateral dentigerous cysts reported in the dental literature. Very few reports showed the occurrence in other sites, such as central incisors, second premolar, and first molars.<sup>3,7-12</sup> Nevertheless, only 8 cases have been shown to be associated with mandibular first molars, as in the present case. Table 1 summarizes the data of bilateral mandibular first molars reported in the dental literature.

It is plausible that bilateral dentigerous cysts are either under-recognized or under-reported, particularly in the third molar area, because not all pericoronal tissues associated with these impacted teeth are sent for histopathological analysis. In our opinion, as the noneruption of first molar teeth is not a common finding, most patients would probably ask their dentist for diagnosis and treatment. This may reflect the true rarity of this condition associated with permanent first molars.

An interesting aspect observed in the present case was how fast the second cyst developed on the right side. When the image exams were obtained for evaluation of the cyst on the left side (first lesion), it was not visible—being detectable only after 2 months, when the patient complained of a swelling in the region. Although the second lesion had developed in only 2 months and the presence of bilateral DCs was rare, the radiographic findings were very suggestive of a DC. Thus, the hypothesis of diagnosis for the second lesion was also DC.

There are 2 common surgical treatments for this condition: (1) marsupialization; and (2) enucleation. Marsupialization is indicated in larger dentigerous cysts, allows lesion decompression, and decreases the cystic cavity. Sometimes, later enucleation can be performed with a less extensive surgical procedure. Marsupialization and enucleation were the treatments of choice for our patient, which agrees with other authors.<sup>17,18</sup> There are few reports in the dental literature reporting spontaneous regression of dentigerous cysts, although they were not confirmed histopathologically.<sup>19,20</sup>

In summary, bilateral dentigerous cysts affecting the permanent first molars are extremely rare and can arise at different times (metachronous). In addition, the presence of bilateral dentigerous cysts does not mean that a patient is syndromic. Correct investigation should be conducted, however, because this possibility does exist.

Year	Gender	Race*	Age (ys)	Treatment
2001	Male	С	8	Enucleation
1998	Female	N/A	3	Enucleation
1996	Male	N/A	7	Enucleation
1989	Male	Bl	5	Enucleation
1989	Male	С	15	Enucleation
1980	Female	С	7	Enucleation
1970	Male	N/A	9	Enucleation
1955	Male	N/A	7	Enucleation
	Year 2001 1998 1996 1989 1989 1980 1970 1955	Year Gender   2001 Male   1998 Female   1996 Male   1998 Male   1989 Male   1989 Male   1989 Male   1989 Male   1980 Female   1990 Male   1980 Female   1990 Male   1980 Female   1990 Male	Year Gender Race*   2001 Male C   1998 Female N/A   1996 Male N/A   1996 Male C   1996 Male C   1996 Male C   1989 Male C   1980 Female C   1980 Female N/A   1980 Female N/A   1955 Male N/A	Year Gender Race* Age (ys)   2001 Male C 8   1998 Female N/A 3   1996 Male N/A 7   1996 Male Bl 5   1989 Male C 15   1980 Female C 7   1980 Female C 7   1980 Female C 7   1980 Female C 7   1980 Female N/A 9   1970 Male N/A 7   1975 Male N/A 7

\* BI=black; C=Caucasian; N/A=not available.

#### Table 1. Summary of Previously Reported Bilateral Mandibular First Molars

## REFERENCES

- Kramer IR, Pindborg JJ, Shear M. World Health Organization: Histological Typing of Odontogenic Tumors. 2<sup>nd</sup> ed. Berlin, Germany: Springer-Verlag; 1992.
- Daley TD, Pringle GA. Relative incidence of odontogenic tumors and oral and jaw cysts in a Canadian population. Oral Surg Oral Med Oral Pathol 1994; 77:276-80.
- 3. Ko KS, Dover DG, Jordan RC. Bilateral dentigerous cysts—report of an unusual case and review of the literature. J Can Dent Assoc 1999;65:49-51.
- Gorlin RJ, Cohen MM, Hennekam RCM. Syndromes of the Head and Neck. 4<sup>th</sup> ed. New York, NY: Oxford University Press; 2001.
- De Biase A, Ottolenghi L, Polimeni A, Benvenuto A, Lubrano R, Magliocca FM. Bilateral mandibular cysts associated with cyclosporine use: A case report. Pediatr Nephrol 2001;16:993-5.
- 6. O'Neil DW, Mosby EL, Lowe JW. Bilateral mandibular dentigerous cyst in a five-year-old child: Report of a case. J Dent Child 1989;56:383-4.
- 7. Carr MM, Anderson RD, Clarke KD. Multiple dentigerous cysts in childhood. J Otolaryngol 1996;25: 267-70.
- 8. Henefer EP. Bilateral dentigerous cysts of the maxilla. Report of a case. Oral Surg Oral Med Oral Pathol 1964;17:296-8.
- 9. Crinzi RA. Bilateral dentigerous cysts of the mandible. Oral Surg Oral Med Oral Pathol 1982;54:367.
- 10. Stanback JS III. The management of bilateral cysts of the mandible. Oral Surg Oral Med Oral Pathol 1970;30:587-91.

- Ustuner E, Fitoz S, Atasoy C, Erden I, Akyar S. Bilateral maxillary dentigerous cysts: A case report. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2003;95: 632-5.
- 12. Freitas DQ, Tempest LM, Sicoli E, Lopes-Neto FC. Bilateral dentigerous cysts: Review of the literature and report of an unusual case. Dentomaxillofac Radiol 2006;35:464-8.
- 13. Sands T, Tocchio C. Multiple dentigerous cysts in a child. Oral Health 1998;88:27-9.
- 14. Eidinger GB. Bilateral dentigerous cysts in the child patient. Report of a case and review of the literature. Univ Tor Dent J 1989;2:20-3.
- 15. Swerdloff M, Alexander SA, Ceen RF, Ferguson FS. Bilateral mandibular dentigerous cysts in a seven-year-old child. J Pedod 1980;5:77-84.
- 16. Tam JC. Bilateral dentigerous cysts involving first molars. J Am Dent Assoc 1955;50:197-8.
- 17. Motamedi MH, Talesh KT. Management of extensive dentigerous cysts. Br Dent J 2005;198:203-6.
- 18. Bodner L. Cystic lesions of the jaws in children. Int J Pediatr Otorhinolaryngol 2002;62:25-9.
- 19. Irving SP. Spontaneous regression of a dentigerous cyst in a middle-aged adult. Oral Surg Oral Med Oral Pathol 1984;57:604.
- 20. Shah N, Thuau H, Beale I. Spontaneous regression of bilateral dentigerous cysts associated with impacted mandibular third molars. Br Dent J 2002;192:75-6.

Copyright of Journal of Dentistry for Children is the property of American Academy of Pediatric Dentistry and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.