

Mutual induction of noncollagenous bone proteins at the interface between epithelial cells and fibroblasts from human periodontal ligament

M. Shimonishi¹, J. Hatakeyama²,
Y. Sasano², N. Takahashi³,
M. Komatsu⁴, M. Kikuchi¹

¹Division of Comprehensive Dentistry, Tohoku University Dental Hospital, Sendai, Japan,

²Division of Craniofacial Development and Regeneration, Department of Craniofacial Engineering and Regeneration, Tohoku University Graduate School of Dentistry, Sendai, Japan,

³Division of Oral Ecology and Biochemistry, Department of Oral Biology Tohoku University Graduate School of Dentistry, Sendai, Japan, and

⁴Division of Operative Dentistry, Department of Restorative Dentistry, Tohoku University Graduate School of Dentistry, Sendai, Japan

Shimonishi M, Hatakeyama J, Sasano Y, Takahashi N, Komatsu M, Kikuchi M. Mutual induction of noncollagenous bone proteins at the interface between epithelial cells and fibroblasts from human periodontal ligament. *J Periodont Res* 2008; 43: 64–75. © 2007 The Authors. Journal compilation © 2007 Blackwell Munksgaard

Background and Objective: Epithelial–mesenchymal interactions are responsible for cell differentiation during periodontal regeneration. The present study was undertaken to examine the expression of alkaline phosphatase and noncollagenous bone proteins, such as osteopontin, osteocalcin and bone sialoprotein, with respect to interaction between the cells of the epithelial rests of Malassez and fibroblasts from human periodontal ligament.

Material and Methods: Explants of human periodontal ligament tissues produced outgrowths containing both putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts in a modified serum-free medium. Putative epithelial rests of Malassez cells cultured alone, and human periodontal ligament fibroblasts cultured alone, were used as controls. The expression levels of amelogenin were analyzed by *in situ* hybridization. The expression and distribution of alkaline phosphatase and noncollagenous bone proteins in both cell populations at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts were analyzed by immunohistochemistry, *in situ* hybridization and reverse transcription-polymerase chain reaction.

Results: Amelogenin mRNA was detected at high levels only in putative epithelial rests of Malassez cells at the interface. Alkaline phosphatase and bone sialoprotein mRNAs were detected significantly at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblast cells. In particular, bone sialoprotein and its mRNA were expressed significantly in human periodontal ligament fibroblasts at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblast cells. The expressions of osteopontin and its mRNA were not different between putative epithelial rests of malassez cells and human periodontal ligament fibroblasts at the interface. Osteocalcin and its mRNA were expressed strongly in putative epithelial rests of

Mitsuru Shimonishi, DDS, PhD, Division of Comprehensive Dentistry, Tohoku University Dental Hospital, 4–1, Seiryō-machi, Aoba-ku, Sendai, 980–8575, Japan
Tel: +81 22 717 8434
Fax: +81 22 717 8434
e-mail: shimo@ddh.tohoku.ac.jp

Key words: cultured epithelial rests of Malassez; *in vitro* model; noncollagenous bone proteins; periodontal maintenance

Accepted for publication January 16, 2007

Malassez cells at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts.

Conclusion: These findings indicate that the epithelial–mesenchymal interaction modulates the expression of alkaline phosphatase, osteocalcin and bone sialoprotein in putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts, suggesting that epithelial–mesenchymal interactions play a role in the maintenance of periodontal ligament.

Human periodontal ligament contains the epithelial rests of Malassez, which appear as cellular clusters in cord- or net-like formations or as isolated islands near the cementum. Hertwig's epithelial root sheath cells represent an extension of the epithelium that participates in coronal amelogenesis and which is retained in association with the developing root. Immediately as the outer layer of the dentin matrix starts to calcify, the epithelial cells of the root sheath separate from the dentin surface, and breaks occur in the continuity of its previously continuous double-layered sheet of cells to form the epithelial rests of Malassez. As a result, the epithelial rests of Malassez are odontogenic epithelial cells derived from Hertwig's epithelial root sheath. Amelogenin is the major enamel protein produced by ameloblasts at the differentiation and secretory stages (1). We demonstrated previously that intense immunoreactivity for amelogenin mRNA is observed in putative epithelial rests of Malassez cells, whereas gingival epithelial cells and human periodontal ligament fibroblasts do not show amelogenin mRNA expression (2). Amelogenin is therefore a useful marker for cultured epithelial rests of Malassez cells.

The formation, turnover and repair of mineralized tissues, such as bone, calcified cartilage and teeth, are all complex processes requiring differentiated cell function and extracellular matrix assembly and degradation during normal development as well as during tissue repair after injury. It has been determined that several of the mineral-binding proteins are capable not only of guiding extracellular calcification, but also of mediating cell dynamics, such as cell adhesion, spreading and migration, in mineral-

ized tissues (3–5). The activity of alkaline phosphatase is required to generate the inorganic phosphate needed for hydroxyapatite crystallization. Extracellular inorganic pyrophosphate is a potent inhibitor of hydroxyapatite formation. Alkaline phosphatase hydrolyzes extracellular inorganic pyrophosphate (6–9). Osteopontin, osteocalcin and bone sialoprotein are the major noncollagenous proteins that are secreted by osteoblastic cells and deposited into the bone matrix (5). Osteopontin is highly phosphorylated and sulfated but is unlikely to be a primary nucleator of hydroxyapatite formation because it was found to appear before bone sialoprotein as a marker of an early stage of bone formation during the regenerative processes (10). Moreover, it has been demonstrated that osteopontin inhibits mineralization (3,10–13). Indeed, osteopontin is also expressed by a number of nonmineralizing tissues, as well as by transformed cells (14,15) and activated lymphocytes and macrophages (3,16), kidney epithelial cells, luminal epithelial cells of several organs (17) and smooth muscle cells in athelerosclerotic lesions of the aorta (18). Osteocalcin is known to localize in bone, cementum and dentin, and in the formative cells of these hard tissues (5,18). Its function, however, is still not clear, although it has been implicated to play a role in delaying nucleation and in preventing excessive crystal growth (19). Bone sialoprotein appears to be unique to mineralized connective tissue. In addition, bone sialoprotein has the potential to mediate the initial formation of hydroxyapatite crystals (20–22).

It appears that Hertwig's epithelial root sheath cells are responsible for initiating the spatial and temporal

differentiation of odontoblasts and cementoblasts into matrix-producing cells. The intermediate layer of cementum may be produced by Hertwig's epithelial root sheath cells (23–29). Moreover, Bosshardt *et al.* suggested that cementoblasts originate from Hertwig's epithelial root sheath (29,30), but species differences make the origin and differentiation of cementoblasts inconsistent between different studies, and therefore the understanding of cementogenesis is still incomplete. Mouri *et al.* demonstrated expression of alkaline phosphatase and osteopontin mRNA in cultured epithelial rests of Malassez cells from human periodontal ligament (31). Rincon *et al.* also demonstrated the production of osteopontin mRNA, but not of alkaline phosphatase mRNA, by cultured porcine epithelial rests of Malassez cells (32,33). Moreover, Hasegawa *et al.* reported that epithelial rests of Malassez cells are immunoreactive for osteopontin and ameloblastin during early cementum repair after the experimental induction of root resorption *in vivo* (26). It is conceivable that epithelial rests of Malassez cells, as well as Hertwig's epithelial root sheath cells, may have a direct or indirect role in the formation of cementum.

Some authors have suggested that epithelial rests of Malassez play a role in the maintenance of the periodontal space. Epithelial rests of Malassez cells were found to be stimulated to a more proliferating state as a result of orthodontic tooth movement, inflammatory conditions, or wound healing (34–39). A regrowth of epithelial rests of Malassez cells may be linked to reconstitution of the periodontal ligament and repair of the resorptive defect in the tooth surface. We demonstrated

previously that epithelial rests of Malassez cells have a high proliferation rate at the interface between epithelial rests of Malassez cells and human periodontal ligament fibroblasts and that the synthesis of type IV collagen and laminin is induced by direct interaction between epithelial rests of Malassez cells and human periodontal ligament fibroblasts (40). Considering the homeostasis of the periodontium, much seems to need to be done regarding the aspects of epithelial-mesenchymal interactions.

The present study was undertaken to determine whether epithelial rests of Malassez cells and human periodontal ligament fibroblasts at the interface *in vitro* have the ability to regulate alkaline phosphatase and the noncollagenous proteins, osteopontin, osteocalcin and bone sialoprotein, supporting their involvement in the maintenance of periodontal ligament.

Material and methods

Cell culture

Freshly extracted third molars from 24 patients between 17 and 25 years of age were obtained from the Oral Surgery Department, Tohoku University Graduate School of Dentistry. Informed consent was obtained from the patients prior to extractions. The human ethics board of Tohoku University Graduate School of Dentistry specifically granted permission for our project to work with human subjects. After washing the teeth several times with alpha minimal essential medium (COSMO BIO Co. Ltd, Tokyo, Japan), supplemented with 10% fetal bovine serum and antibiotics (60 µg/mL of kanamycin, 20 units/mL of penicillin G, 10 µg/mL of amphotericin B), human periodontal ligament explants attached to the mid-third of each root were removed carefully from the root with a scalpel. The explants were plated into 35-mm culture dishes using the supplemented alpha minimal essential medium and produced outgrowths that were primarily composed of fibroblasts. After 1 wk the explants were cultured in a modified serum-free medium [3 : 1 (v/v) MCDB153 med-

ium; Sigma Chemical Co., St Louis, MO, USA] supplemented with 5 µg/mL of insulin (Sigma), 0.5 µg/mL of hydrocortisone (Sigma), 10 µg/mL of transferrin (Sigma), 14.1 µg/mL of phosphorylethanolamine (Sigma), 10 ng/mL of epidermal growth factor (Sigma): alpha minimal essential medium (COSMO BIO Co. Ltd) including 40 µg/mL of bovine pituitary extract (Kyokuto, Tokyo, Japan) and antibiotics (40). Cultures were incubated at 37°C in a humidified atmosphere of 95% air and 5% CO₂. This resulted in outgrowths of epithelial cells as well as of fibroblasts.

Differential adhesion to the surface in the presence of 0.075 mg/mL of protease solution (Sigma) was used to produce cultures that were predominantly of one cell type [i.e. either fibroblasts (5–10 min) or epithelial cells (15–20 min)] from cultures that contained both cell types (41). Putative epithelial rests of Malassez cells cultured alone, and human periodontal ligament fibroblasts cultured alone, were used as controls.

Immunohistochemistry

The cells were fixed with 4% paraformaldehyde at room temperature for 10 min, and 3% H₂O₂ was used to inhibit endogenous peroxidase. The cells were then incubated with 5% normal goat serum for 30 min to block nonspecific binding, after which they were reacted at 4°C overnight with monoclonal mouse antihuman cytokeratin AE1/AE3 (1 : 50) (Dako, Carpinteria, CA, USA) to confirm the presence of epithelial cells. In addition, the cells were incubated at 4°C overnight with primary antibodies to monoclonal mouse antihuman osteopontin (0.1 µg/mL) (IBL, Gunma, Japan), polyclonal rabbit antihuman osteocalcin (1 : 250) (Biomedical Technology Inc., Stoughton, MA, USA) and polyclonal rabbit antihuman bone sialoprotein (LF-84) (1 : 400) (courtesy of Dr L. W. Fisher, National Institutes for Dental and Craniofacial Research, Bethesda, MD, USA) (42–45). After rinsing in phosphate-buffered saline, the cells were incubated with biotinylated immuno-

globulin at room temperature for 30 min and stained by the avidin-biotinylated peroxidase complex using an ExtrAvidin[®] peroxidase staining kit (Sigma) and the 3-amino-9-ethyl-carbazole chromogen kit (Sigma). Mayer's hematoxylin solution was applied for counterstaining. Phosphate-buffered saline, instead of the primary antibody, and rabbit serum and phosphate-buffered saline, instead of the primary antiserum, were used for control staining.

Proliferating, DNA-synthesizing cells in cocultures were identified by nuclear incorporation of 100 µmol/L of 5-bromo-2'-deoxyuridine for 24 h. Labeled nuclei were detected with monoclonal mouse antibody to 5-bromo-2'-deoxyuridine (Boehringer Mannheim Biochemica, Mannheim, Germany) for 30 min at 37°C. After washing three times in phosphate-buffered saline, the cells were incubated with horseradish peroxidase-conjugated goat antimouse immunoglobulin (Chemicon International Inc., Temecula, CA, USA) for 30 min at 37°C and stained with 3,3'-diaminobenzidine and H₂O₂.

Determination of alkaline phosphatase activity

The cells were fixed with 4% paraformaldehyde at room temperature for 10 min. To determine the localization of alkaline phosphatase in cultured cells, the cells were stained histochemically for alkaline phosphatase according to a modified method of the Azo-dye coupling method (46).

In situ hybridization

The oligonucleotide probes used for the *in situ* hybridization were synthesized by Nihon Gene Research Laboratories Inc. (Sendai, Japan). The sequences are shown in Table 1. A biotin label was added at the 3' end. A computer-assisted search (GenBank) of the antisense sequences shown in Table 1, as well as that of sense sequences, revealed no significant homology with any known sequences other than those of the amelogenin chain, the alkaline phosphatase chain,

Table 1. Oligonucleotide probes used for the in situ hybridization

Oligo name	Sequence (5'-3')	Mer	Label	Reference
Amelogenin	CAT GGG TTC GTA ACC ATA GGA AGG	24	3' Biotin	(47)
Alkaline phosphatase	ACA TGA TGA CAT TCT TAG CCA CGT	24	3' Biotin	(48)
Osteopontin	ATG GCT TTC GTT GGA CTT ACT TGG	24	3' Biotin	(49)
Osteocalcin	GCG AGG AGT GTG TGA GGG CTC ATG	24	3' Biotin	(50)
Bone sialoprotein	ATT TTT CAT TGA GAA AGC ACA GGC	24	3' Biotin	(51)

the osteopontin chain, the osteocalcin chain and the bone sialoprotein chain.

In situ hybridization was carried out using the *In Situ* Hybridization Detection Kit for Biotin Labeled Probes (Sigma). Briefly, the cells were fixed with 4% paraformaldehyde at room temperature for 10 min. They were then immersed in phosphate-buffered saline containing RNase inhibitor at room temperature and 3% H₂O₂ was used to inhibit endogenous peroxidase. The specimens were hybridized with biotin-labeled probes, present in the hybridization solution, in the humid incubation chamber overnight at 37°C. After washing in phosphate-buffered saline, the specimens were reacted with blocking solution (5% bovine serum albumin, 500 µg/mL of normal sheep IgG, 100 µg/mL of salmon testicular DNA and 100 µg/mL of yeast tRNA in phosphate-buffered saline) at room temperature for 15 min. The cells were then incubated with ExtrAvidin® peroxidase solution (Sigma) at 37°C for 20 min and reacted with biotin-conjugated anti-avidin immunoglobulin (Sigma) in a humid chamber at 37°C for 30 min. After washing three times in phosphate-buffered saline, the sites of peroxidase activity were visualized using a solution containing 3,3'-diaminobenzidine and H₂O₂ and counterstained by Mayer's hematoxylin solution. Sense oligonucleotide probes were used for control staining.

Reverse transcription polymerase chain reaction (RT-PCR)

The interface area cells, which are present between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts, were obtained using silicone cylinders (Fig. 1A,B). Briefly, silicone cylinders (5 mm in diameter) pasted with white




Fig. 1. (A) Schematic diagram showing the preparation of the samples at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts, except for only putative epithelial rests of Malassez cells or only human periodontal ligament fibroblasts away from the interface in the same dishes. The silicone cylinders (5 mm in diameter) pasted with white petrolatum on the bottom, were placed on the dishes. (B) Phase-contrast photomicrograph taken of primary cultured putative epithelial rests of Malassez cells (ERM) and of human periodontal ligament fibroblasts (FB) at the interface (I).

petrolatum on the bottom, were placed on the dishes after aspirating the medium and washing twice with phosphate-buffered saline. Fifty microlitres of Buffer RLT (buffer containing guanidine thiocyanate and β-mercaptoethanol) (Qiagen Pty Ltd, Victoria, Australia) was added to the silicone cylinders. The cells disrupted with Buffer RLT were collected in the microcentrifuge tubes. Putative epithelial rests of Malassez cells cultured alone and human periodontal ligament fibroblasts cultured alone were sampled as controls. Total cellular RNA was isolated from cultured cells, according to the manufacturer's instructions, using an RNeasy® Mini

Kit (Qiagen Pty Ltd), and 0.05 µg/µL of total RNA was used as a template for RT-PCR. One-step RT-PCR was performed using SuperScript™ one-step RT-PCR with the Platinum® Taq kit (Invitrogen Corp., Carlsbad, CA, USA) according to the manufacturer's instructions and a Programmable Thermal Controller PTC-100 (MJ Research, Watertown, MA, USA). To quantify the expression of alkaline phosphatase and noncollagenous bone proteins (osteopontin, osteocalcin and bone sialoprotein), semiquantitative RT-PCR relative to glyceraldehyde-3-phosphate dehydrogenase was performed. Amplimers designed for alkaline phosphatase, osteopontin, osteocalcin,

Table 2. Primer pairs used for reverse transcription-polymerase chain reaction amplification

Gene (fragment)	Primer sequences	Denaturation/annealing/ extension (°C)	Cycle	Reference
Alkaline phosphatase (475 bp)	5'-ACGTGGCTAAGAATGTCATC-3' 5'-CTGGTAGGCGATGTCCTTA-3'	94/58/72	40	(52)
Osteopontin (126 bp)	5'-CCAAGTAAGTCCAACGAAAG-3' 5'-GGTGATGTCCTCGTCTGTA-3'	94/58/72	40	(53)
Osteocalcin (230 bp)	5'-GGCAGCGAGGTAGTGAAGA-3' 5'-CTGGAGAGGAGCAGAACTG-3'	94/58/72	40	(54)
Bone sialoprotein (248 bp)	5'-CAACAGCACAGAGGCAGAA-3' 5'-CGTACTCCCCCTCGTATTC-3'	94/58/72	40	(51)
GAPDH (485 bp)	5'-TGTTTGTGATGGGTGTGAA-3' 5-ATGGGAGTTGCTGTTGAAG-3'	94/58/72	40	(55)

GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

bone sialoprotein and glyceraldehyde-3-phosphate dehydrogenase, and related information, are provided in Table 2. The thermal profile for human alkaline phosphatase, osteopontin, osteocalcin, bone sialoprotein and glyceraldehyde-3-phosphate dehydrogenase amplification was 40 cycles, starting with denaturation for 1 min at 94°C followed by 1 min of annealing at 58°C and 1 min of extension at 72°C. The PCR products were subjected to electrophoresis, and digital images were obtained and analyzed with the use of IMAGE J (NIH image software, Bethesda, MD, USA). The data were consistent, as confirmed by three independent experiments. Statistical analysis of the results was carried out using the Bonferroni/Dunn post-hoc test, and *p*-values of < 0.05 were considered significant.

Results

Immunohistochemistry and determination of alkaline phosphatase activity

Putative epithelial rests of Malassez cells stained positive for broad-spectrum antibodies to cytokeratins (AE1/AE3), indicating their epithelial origin, whereas human periodontal ligament fibroblasts did not show cytokeratin expression at the interface in the same dishes (Fig. 2A). Putative epithelial rests of Malassez cells incorporated 5-bromo-2'-deoxyuridine more extensively than human periodontal ligament fibroblasts derived from the same periodontal ligament explant, indicating that putative epithelial rests of

Malassez cells have a higher proliferation rate than human periodontal ligament fibroblasts (Fig. 2C).

Alkaline phosphatase activity was expressed strongly at the interface (Fig. 3A). Putative epithelial rests of Malassez cells cultured alone and human periodontal ligament fibroblasts cultured alone also stained strongly for alkaline phosphatase activity (Fig. 3B,C). Intense immunoreactivity for osteopontin was observed in both putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts at the interface (Fig. 3D). Osteopontin was also detected in putative epithelial rests of Malassez cells cultured alone and in human periodontal ligament fibroblasts cultured alone (Fig. 3E,F). Strong immunoreactivity for osteocalcin was observed in putative epithelial rests of Malassez cells but not in human periodontal ligament fibroblasts at the interface (Fig. 3G). In contrast, bone sialoprotein was detectable in human periodontal ligament fibroblasts but not in putative epithelial rests of Malassez cells at the interface (Fig. 3J). Osteocalcin and bone sialoprotein proteins were not detected in putative epithelial rests of Malassez cells cultured alone (Fig. 3H,K), but were present in human periodontal ligament fibroblasts cultured alone (Fig. 3I,L).

In situ hybridization to localize mRNAs for amelogenin, alkaline phosphatase and noncollagenous bone proteins

To localize the mRNA expression of amelogenin, alkaline phosphatase and

noncollagenous bone proteins, the cells were hybridized *in situ* with biotin-labeled antisense oligo-DNA probes. Putative epithelial rests of Malassez cells showed strongly positive signals for amelogenin mRNA. However, the amelogenin mRNA signal was not detectable in human periodontal ligament fibroblasts (Fig. 2B). These results supported that putative epithelial rests of Malassez cells are different from human periodontal ligament fibroblasts and are derived from the odontogenic epithelial origin.

In situ hybridization analyses also showed considerable differences between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts and reflected the histochemical and immunohistochemical data. The mRNAs of alkaline phosphatase, osteopontin and osteocalcin were expressed strongly in putative epithelial rests of Malassez cells at the interface, whereas bone sialoprotein mRNA expression was not detectable (Fig. 4A,D,G,J). On the other hand, mRNAs of alkaline phosphatase, osteopontin and bone sialoprotein were expressed strong in human periodontal ligament fibroblasts at the interface, whereas osteocalcin mRNA expression was not detectable (Fig. 4A,D,G,J). Putative epithelial rests of Malassez cells cultured alone showed strong positive signals for alkaline phosphatase and osteopontin mRNAs (Fig. 4B,E). mRNA for osteocalcin was detectable only very weakly (Fig. 4H) and mRNA for bone sialoprotein was not expressed at all in putative




Fig. 2. (A) Photomicrograph showing strong immunoreactivity for cytokeratin AE1/AE3 in putative epithelial rests of Malassez cells. Immunostaining for cytokeratin in human periodontal ligament fibroblasts is not present. (B) Photomicrograph showing strong immunoreactivity for amelogenin mRNA in putative epithelial rests of Malassez cells. Immunostaining for amelogenin mRNA in human periodontal ligament fibroblasts is not present. (C) Photomicrograph showing assessment of proliferation by 5-bromo-2'-deoxyuridine labeling in putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts. (D) Photomicrograph showing negative-control immunostaining in putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts. (E) Photomicrograph showing the negative control hybridized *in situ* with biotin-labeled sense oligo-DNA probes for amelogenin mRNA in putative epithelial rests of Malassez cells and in human periodontal ligament fibroblasts. ERM, epithelial rests of Malassez cells; FB, human periodontal ligament fibroblasts.

epithelial rests of Malassez cells cultured alone (Fig. 4K). On the other hand, mRNAs for osteopontin, osteocalcin and bone sialoprotein were detectable in human periodontal ligament fibroblasts cultured alone (Fig. 4F,I,L), and a strong signal for alkaline phosphatase mRNA was detected in human periodontal ligament fibroblasts cultured alone (Fig. 4C).

RT-PCR

The expression of four genes (mRNAs of alkaline phosphatase, osteopontin, osteocalcin and bone sialoprotein) was investigated in three cell populations, both putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts at the interface, putative epithelial rests of Malassez cells cultured alone and human perio-

dontal ligament fibroblasts cultured alone, using RT-PCR (Fig. 5). The mRNA intensities relative to that of glyceraldehyde-3-phosphate dehydrogenase are presented in Fig. 6. The expression of mRNAs for alkaline phosphatase and bone sialoprotein were significantly stronger at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts ($p < 0.05$). On the other hand, there were no significant differences in the relative intensities of mRNAs for osteopontin and osteocalcin among the three cell populations. The mRNA for osteocalcin was detectable only very weakly, and the mRNA for bone sialoprotein was not expressed at all in putative epithelial rests of Malassez cells cultured alone. A strong signal for alkaline phosphatase mRNA expression was detected in human periodontal ligament fibroblasts cultured alone, and mRNAs for osteopontin, osteocalcin and bone sialoprotein were detectable in human periodontal ligament fibroblasts cultured alone.

Discussion

Amelogenin is the major enamel protein produced by ameloblasts at the differentiation and secretory stages (3). Amelogenin has been observed to be localized in the enamel matrix, in normal ameloblasts, in odontogenic tumor cells, in Hertwig's epithelial root sheath cells and their extracellular matrices *in vivo*, in mantle dentin and in odontoblasts (3,30,56–60). In the present study, amelogenin mRNA expression distinguished putative epithelial rests of Malassez cells from human periodontal ligament fibroblasts at the interface, and putative epithelial rests of Malassez cells were the odontogenic epithelial cells derived from Hertwig's epithelial root sheath (Fig. 2B). Bosshardt *et al.* suggest that Hertwig's epithelial root sheath cells occasionally assume a lingering ameloblastic activity at the start of root formation in the pig, but they do not support the hypothesis of a causal relationship between enamel matrix proteins and cementogenesis (30). Moreover, they indicated that ectopic enamel deposits




Fig. 3. (A–C) Photomicrographs showing strong alkaline phosphatase activity in interface area cells, in putative epithelial rests of Malassez cells cultured alone and in putative human periodontal ligament fibroblasts cultured alone. (D–F) Photomicrographs showing strong immunoreactivity for osteopontin in interface area cells. Osteopontin was also detected in putative epithelial rests of Malassez cells cultured alone and in putative human periodontal ligament fibroblasts cultured alone. (G–I) Photomicrographs showing strong immunoreactivity for osteocalcin in putative epithelial rests of Malassez cells at the interface area but very weak immunoreactivity for osteocalcin in putative human periodontal ligament fibroblasts cultured alone. Osteocalcin was not detected in putative epithelial rests of Malassez cells cultured alone. (J–L) Photomicrographs showing no immunoreactivity for bone sialoprotein in putative epithelial rests of Malassez of the interface area and in putative epithelial rests of Malassez cells cultured alone, and weak immunoreactivity for bone sialoprotein in human periodontal ligament fibroblasts of the interface area and putative human periodontal ligament fibroblasts cultured alone. ERM, epithelial rests of Malassez cells; FB, human periodontal ligament fibroblasts; IEF, interface area; PE, putative epithelial rests of Malassez cells cultured alone; PF, putative human periodontal ligament fibroblasts cultured alone.

on the root retain a high amount of amelogenin, whereas cementicles contain bone sialoprotein and osteopontin, typically found in bone and cementum (61). Hertwig's epithelial root sheath cells may be involved in both enamel formation and cementum formation, and they may participate in either enamel formation or cementum formation in different situations.

It has been suggested that epithelial rests of Malassez may protect the root surface from resorption, prevent ankylosis and consequently maintain the integrity of the periodontal ligament. Some studies have postulated that proliferating epithelial rests of Malassez cells may be involved in reconstitution of the periodontium (38,39,62). We showed, in the present study, that putative epithelial rests of Malassez cells have a higher proliferation rate, according to the incorporation of 5-bromo-2'-deoxyuridine extensively at the interface (Fig. 2C). Moreover, in our previous study, the synthesis of type IV collagen and laminin including the basement membrane was induced by the interaction at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts (40). The initiation of cementogenesis is at least temporally linked to the disintegration process of Hertwig's epithelial root sheath. It is natural that the basement membrane is broken down in this disintegration. The disruption of a basement membrane may be correlated to the expression of osteopontin and bone sialoprotein. Osteopontin and bone sialoprotein are members of the small integrin-binding ligand, N-linked glycoprotein (SIBLING) family. The SIBLING family can specifically bind pro-matrix metalloproteinases (pro-MMPs) and activate MMPs. MMPs play a crucial role in extracellular matrix degradation during tissue remodeling. Moreover, epithelial growth *in vitro* and during wound healing *in vivo* is intrinsically linked to the expression of MMPs (63,64).

Alkaline phosphatase and noncollagenous bone proteins are considered to have multiple functions related to the formation, turnover and repair of collagen-based mineralized tissues. Previ-




Fig. 4. (A–C) Photomicrographs showing strong immunoreactivity for alkaline phosphatase mRNA in interface area cells, in putative epithelial rests of Malassez cells cultured alone and in putative human periodontal ligament fibroblasts cultured alone. (D–F) Photomicrographs showing strong immunoreactivity for osteopontin mRNA in interface area cells. Osteopontin mRNA was also detected in putative epithelial rests of Malassez cells cultured alone and in putative human periodontal ligament fibroblasts cultured alone. (G–I) Photomicrographs showing strong immunoreactivity for osteocalcin mRNA in putative epithelial rests of Malassez cells of the interface area but very weak immunoreactivity for osteocalcin mRNA in putative human periodontal ligament fibroblasts cultured alone. Osteocalcin mRNA was not detected in putative epithelial rests of Malassez cells cultured alone. (J–L) Photomicrographs showing no immunoreactivity for bone sialoprotein mRNA in putative epithelial rests of Malassez cells of the interface area and in putative epithelial rests of Malassez cells cultured alone, but weak immunoreactivity for bone sialoprotein mRNA in human periodontal ligament fibroblasts of the interface area and in putative human periodontal ligament fibroblasts cultured alone.

ous studies have shown that alkaline phosphatase and noncollagenous bone proteins are present in periodontal ligament, mantle dentin, acellular and cellular cementum, and cementoblasts (10,20,28,52,65). An understanding of the distribution of these proteins within the human periodontium is of fundamental importance. When we set out to determine the level of mRNA expression of alkaline phosphatase and noncollagenous bone proteins in the samples extracted at the interface, only alkaline phosphatase mRNA expression was significantly higher at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts. Although alkaline phosphatase and osteopontin were expressed strongly in putative epithelial rests of Malassez cells cultured alone and in human periodontal ligament fibroblasts cultured alone, the interaction has a strong influence on alkaline phosphatase expression. In fact, it can be easy that alkaline phosphatase expresses under various conditions, e.g., in mineralization medium which includes ascorbic acid, β -glycerophosphate and dexamethasone (66–68). There were no significant differences in osteopontin mRNA expression among the three cell populations. Osteopontin may not be directly associated with the interaction between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts, although it can be associated with mineralization and cell adhesion in each cell (3,10–12,26,31). Tissue nonspecific alkaline phosphatase and osteopontin can be considered as differentiation markers because they are not exclusive to bone and cementum and they can be detected during the differentiation stages on proliferating precursor cells (3,10,52).

On the other hand, osteocalcin and bone sialoprotein have a very restricted tissue distribution and are almost exclusively produced by the hard tissue formation cells. The level of osteocalcin was increased at the interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblast cells, whereas osteocalcin mRNA was not detected in putative epithelial rests of Malassez




Fig. 5. Reverse transcription-polymerase chain reaction analysis of the mRNA expression of alkaline phosphatase, osteopontin, osteocalcin and bone sialoprotein in cultures of interface area cells, of putative epithelial rests of Malassez cells cultured alone and of putative human periodontal ligament fibroblasts cultured alone. The mRNAs were extracted from the samples including both putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts at the interface (interface area cells), putative epithelial rests of Malassez cells cultured alone, or human periodontal ligament fibroblasts cultured alone. Reverse transcription-polymerase chain reaction products were separated by electrophoresis on a 2% agarose gel and stained with ethidium bromide. GAPDH, glyceraldehyde-3-phosphate dehydrogenase; IEF, the cells including both putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts at the interface; PE, putative epithelial rests of Malassez cells cultured alone, PF, putative human periodontal ligament fibroblasts cultured alone.

cells cultured alone. Moreover, immunohistochemical and *in situ* hybridization data indicated that stronger expression and a wider distribution of osteocalcin were found in putative epithelial rests of Malassez cells at the interface compared with putative epithelial rests of Malassez cells cultured alone, but osteocalcin was not detectable in human periodontal ligament fibroblasts at the interface. The interface between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts showed a stronger expression of bone sialoprotein mRNA compared with human periodontal ligament fibroblasts cultured alone, whereas immunohistochemical and *in situ* hybridization data indicated that bone sialoprotein was not detectable in putative epithelial rests of Malassez cells. When the RT-PCR findings of increased levels of osteocalcin and bone sialoprotein mRNA can reflect the immunohistochemical and *in situ*

hybridization data, interaction between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts facilitate osteocalcin expression in putative epithelial rests of Malassez cells and bone sialoprotein expression in human periodontal ligament fibroblasts.

A hypothesis advanced by Slavkin & Boyde proposes that Hertwig's epithelial root sheath cells synthesize proteins that are related to enamel polypeptides, and that these Hertwig's epithelial root sheath-derived proteins are instructive for ectomesenchymal determination, leading to cementoblast differentiation and acellular cementum (23). It has also been suggested that the intermediate layer of the cementum seen in rodent molars is produced by Hertwig's epithelial root sheath cells (23–30). We also indicated that putative epithelial rests of Malassez cells have the ability to produce amelogenin, alkaline phosphatase and

the noncollagenous proteins, osteopontin and osteocalcin, supporting their involvement in cementum formation. Bone sialoprotein was detectable in human periodontal ligament fibroblasts, but not in epithelial rests of Malassez cells. Previous reports indicated that bone sialoprotein is detectable in cementum and alveolar bone, but not in periodontal ligament *in vivo*, whereas alkaline phosphatase and osteopontin are detectable in normal periodontal ligament tissue (52,68). The origin of cementogenesis is still controversial. It appears that the extracellular matrix and the constructed cementum are formed by both epithelial rests of Malassez cells and human periodontal ligament fibroblasts, but it is difficult to identify the role of cementoblasts with certainty.

In contrast, immunohistochemical and *in situ* hybridization data indicated that osteocalcin was not detectable in human periodontal ligament fibroblasts in close proximity to epithelial rests of Malassez cells at the interface, while the RT-PCR showed that weak signs of osteocalcin mRNA could be detected in human periodontal ligament fibroblasts cultured alone. These findings suggested that the interaction between epithelial rests of Malassez cells and human periodontal ligament fibroblasts suppressed the production of osteocalcin in human periodontal ligament fibroblasts at the interface. Osteocalcin is one of the extracellular matrix proteins of cementum and bone that has been implicated to have a role in calcification. However, Ducy *et al.* reported that the finding of increased bone density in osteocalcin knockout mutant mice implicates osteocalcin as a potential inhibitor of bone formation (19). Aubin *et al.* showed that over a time course of osteoblast differentiation *in vitro*, the expression of type I collagen was found to be high and then to decrease; the level of alkaline phosphatase increased but decreased when mineralization was well progressed; osteopontin appeared before certain other matrix proteins, including bone sialoprotein and osteocalcin; bone sialoprotein was first detected in differentiated osteoblasts forming





Alkaline phosphatase**Osteopontin****Osteocalcin****Bone sialoprotein**

Fig. 6. Graphical representation of relative band intensities standardized by glyceraldehyde-3-phosphate dehydrogenase (mean \pm standard deviation, $n = 3$; significant difference: *, $p < 0.05$). GAPDH, glyceraldehyde-3-phosphate dehydrogenase; IEF, interface area cells, comprising both putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts; PE, putative epithelial rests of Malassez cells cultured alone, PF, putative human periodontal ligament fibroblasts cultured alone.

bone; and osteocalcin appeared with mineralization (5). We showed, in the present study, that bone sialoprotein was detectable, but osteocalcin was not, in human periodontal ligament fibroblasts at the interface. Although human periodontal ligament fibroblasts have a potentially recognizable stage of differentiation, they may not divide and differentiate into the mature cells forming the hard tissues because of the invasion of the proliferating epithelial rests of Malassez cells at the interface.

Cell culture favors growth and selects for the most rapidly proliferating cells. Cell culture data should be

cautiously extrapolated to the *in vitro* situation, particularly in providing evidence for cellular origins. However, this should not preclude the fact that valuable information can be obtained from *in vitro* studies, because the results from *in vitro* studies reflect various situations *in vivo*. Moreover, cells derived from periodontal ligament should produce the same proteins and mRNAs as those expressed in periodontal ligament tissues.

In conclusion, the data presented here indicated that the interaction between putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts modulates the

expression of alkaline phosphatase, osteocalcin and bone sialoprotein in putative epithelial rests of Malassez cells and human periodontal ligament fibroblasts *in vitro*. Considering the homeostasis of the periodontium, we suggest that epithelial-mesenchymal interactions are highly associated with mineralization in periodontal ligament and play a role in the maintenance of the periodontal ligament.

Acknowledgements

We gratefully acknowledge Shigeki Matsumoto for kindly supplying human extracted third molars. This work was supported by a Grant-in-Aid for Encouragement of Young Scientists (B) (No. 15791088) from the Japan Society for the Promotion of Science, Japan.

References

1. Uchida T, Tanabe T, Fukae M. Immunocytochemical localization of amelogenins in the deciduous tooth germs of the human fetus. *Arch Histol Cytol* 1989;**52**: 543–552.
2. Shimonishi M, Hatakeyama J, Sasano Y *et al*. *In vitro* differentiation of epithelial cells cultured from human periodontal ligament. *J Periodont Res* 2007. Epub ahead of print, doi: 10.1111/j.1600-0765.2007.00974.x.
3. McKee MD, Nanci A. Secretion of osteopontin by macrophages and its accumulation at tissue surfaces during wound healing in mineralized tissues: a potential requirement for macrophage adhesion and phagocytosis. *Anat Rec* 1996;**245**:394–409.
4. Pitaru S, Narayanan SA, Olson S *et al*. Specific cementum attachment protein enhances selectively the attachment and migration of periodontal cells to root surfaces. *J Periodont Res* 1995;**30**:360–368.
5. Aubin JE, Liu F, Malaval L, Gupta AK. Osteoblast and chondroblast differentiation. *Bone* 1995;**17**:77S–83S.
6. Beertsen W, Van Den Bos T, Everts V. The possible role of alkaline phosphatase in acellular cementum formation. *J Biol Buccale* 1990;**18**:203–205.
7. Beertsen W, Van Den Bos T. Alkaline phosphatase induces the deposition of calcified layers in relation to dentin: an *in vitro* study to mimic the formation of afibrillar acellular cementum. *J Dent Res* 1991;**70**:176–181.
8. Hessele L, Johnson KA, Anderson HC *et al*. Tissue-nonspecific alkaline phos-

- phatase and plasma cell membrane glycoprotein-1 are central antagonistic regulators of bone mineralization. *Proc Natl Acad Sci USA* 2002;**99**:9445–9449.
9. Foster BL, Nociti FH Jr, Swanson EC *et al*. Regulation of cementoblast gene expression by inorganic phosphate *in vitro*. *Calcif Tissue Int* 2006;**78**:103–112.
 10. Lekic P, Sodek J, McCulloch CAG. Osteopontin and bone sialoprotein expression in regenerating rat periodontal ligament and alveolar bone. *Anat Rec* 1996;**244**:50–58.
 11. Worcester EM, Blumenthal SS, Beshensky AM, Lewand DL. The calcium oxalate crystal growth inhibitor protein produced by mouse kidney cortical cells in culture is osteopontin. *J Bone Miner Res* 1992; **7**:1029–1036.
 12. Boskey AL, Maresca M, Ullrich W, Doty SB, Butler WT, Prince CW. Osteopontin–hydroxyapatite interactions *in vitro*: inhibition of hydroxyapatite formation and growth in a gelatin-gel. *Bone Miner* 1993;**22**:147–159.
 13. Hunter GK, Kyle CL, Goldberg HA. Modulation of crystal formation by bone phosphoprotein: structural specificity of the osteopontin-mediated inhibition of hydroxyapatite formation. *Biochem J* 1994;**300**:723–728.
 14. Craig AM, Nemir M, Mukherjee BB, Chambers AF, Denhardt DT. Osteopontin, a transformation-associated cell adhesion phosphoprotein, is induced by 12-O-tetradecanoyl-phorbol 13-acetate in mouse epidermis. *J Biol Chem* 1989;**264**: 9682–9689.
 15. Senger DR, Peruzzi CA, Papadopoulos A. Elevated expression of secreted phosphoprotein I (osteopontin, 2ar) as a consequence of neoplastic transformation. *Anticancer Res* 1989;**9**:1291–1300.
 16. Patarca R, Freeman GJ, Singh RP *et al*. Structural and functional studies of the early T lymphocyte activation (Eta-1) gene. *J Exp Med* 1989;**170**:145–161.
 17. Butler WT. The nature and significance of osteopontin. *Connect Tissue Res* 1989;**23**:123–136.
 18. Giachelli CM, Schwartz SM, Liaw L. Molecular and cellular biology of osteopontin: potential role in cardiovascular disease. *Trends Cardiovasc Med* 1995;**5**: 88–95.
 19. Ducy P, Desbois C, Boyce B *et al*. Increased bone formation in osteocalcin-deficient mice. *Nature* 1996;**382**:448–452.
 20. MacNeil LR, Berry J, Strayhorn C, Somerman MJ. Expression of bone sialoprotein mRNA by cells lining the mouse tooth root during cementogenesis. *Arch Oral Biol* 1996;**41**:827–835.
 21. Chen J, Shapiro HS, Sodek J. Developmental expression of bone sialoprotein mRNA in rat mineralized connective tissues. *J Bone Miner Res* 1992;**7**:987–997.
 22. Kasugai S, Todescan R Jr, Nagata T, Yao K-L, Butler WT, Sodek J. Expression of bone matrix proteins associated with mineralized tissue formation by adult rat bone marrow cells *in vitro*: inductive effects of dexamethasone on the osteoblastic phenotype. *J Cell Physiol* 1991; **147**:111–120.
 23. Slavkin HC, Boyde A. Cementum: an epithelial secretory product? *J Dent Res* 1974;**53**:157.
 24. Fong CD, Hammarström L. Expression of amelogenin and amelogenin in epithelial sheath remnants of fully formed rat molars. *Oral Surg Oral Med Oral Pathol* 2000;**90**:218–223.
 25. Hammarström L. Enamel matrix, cementum development and regeneration. *J Clin Periodontol* 1997;**24**:658–668.
 26. Hasegawa N, Kawaguchi H, Ogawa T, Uchida T, Kurihara H. Immunohistochemical characteristics of epithelial cell rests of Malassez during cementum repair. *J Periodont Res* 2003; **38**:51–56.
 27. Tenorio D, Cruchley A, Hughes FJ. Immunocytochemical investigation of the rat cementoblast phenotype. *J Periodont Res* 1993;**28**:411–419.
 28. McKee MD, Zalzal S, Nanci A. Extracellular matrix in tooth cementum and mantle dentin: localization of osteopontin and other noncollagenous proteins, plasma proteins, and glycoconjugates by electron microscopy. *Anat Rec* 1996;**245**: 293–312.
 29. Bosshardt DD, Shroeder HE. Cementogenesis reviewed: a comparison between human premolar and rodent molars. *Anat Rec* 1996;**245**:267–292.
 30. Bosshardt DD, Nanci A. Hertwig's epithelial root sheath, enamel matrix proteins, and initiation of cementogenesis in porcine teeth. *J Clin Periodont* 2004;**31**: 184–192.
 31. Mouri Y, Shiba H, Mizuno M, Noguchi T, Ogawa T, Kurihara H. Differential gene expression of bone-related proteins in epithelial and fibroblastic cells derived from human periodontal ligament. *Cell Biol Int* 2003;**27**:519–524.
 32. Rincon JC, Xiao Y, Young WG, Bartold PM. Production of osteopontin by cultured porcine epithelial cell rests of Malassez. *J Periodont Res* 2005;**40**:417–426.
 33. Rincon JC, Young WG, Bartold PM. The epithelial cell rests of Malassez – a role in periodontal regeneration? *J Periodont Res* 2006;**41**:245–252.
 34. Loe H, Waerhaug J. Experimental replantation of teeth in dogs and monkeys. *Arch Oral Biol* 1961;**3**:176–184.
 35. Spouge JD. A new look at the rests of Malassez. A review of their embryological origin, anatomy, and possible role in periodontal health and disease. *J Periodontol* 1980;**51**:437–444.
 36. Lindskog S, Blomlof L, Hammarström L. Evidence for a role of odontogenic epithelium in maintaining the periodontal space. *J Clin Periodont* 1988;**15**:371–373.
 37. Wallace JA, Vergona K. Epithelial rests' function in replantation: is splinting necessary in replantation? *Oral Surg Oral Med Oral Pathol* 1990;**70**:644–649.
 38. Brice GL, Sampson WJ, Sims MR. An ultrastructural evaluation of the relationship between epithelial rests of Malassez and orthodontic root resorption and repair. *Aust Orthod J* 1991;**12**:90–94.
 39. Gilhuus-Moe O, Kvam E. Behavior of the epithelial remnants of Malassez following experimental movement of rat molars. *Acta Odont Scand* 1972;**30**:139–149.
 40. Shimonishi M, Sato J, Takahashi N, Komatsu M. Expression of type IV collagen and laminin at the interface between epithelial cells and fibroblasts from human periodontal ligament. *Eur J Oral Sci* 2005;**113**:34–40.
 41. Brunette DM, Melcher AH, Moe HK. Culture and origin of epithelium-like and fibroblast like cells from porcine periodontal ligament explants and cell suspensions. *Arch Oral Biol* 1976;**21**:393–400.
 42. Fisher LW, Stubbs JT III, Young MF. Antisera and cDNA probes to human and certain animal model bone matrix non-collagenous proteins. *Acta Orthop Scand* 1995;**66**:61–65.
 43. Mintz KP, Grzesik WJ, Midura RJ, Robey PG, Termine JD, Fisher LW. Purification and fragmentation of non-denatured bone sialoprotein: evidence for a cryptic, RGD-resistant cell attachment domain. *J Bone Miner Res* 1993;**8**:985–995.
 44. Bianco P, Riminucci M, Silvestrini G *et al*. Localization of bone sialoprotein (BSP) to Golgi and post-Golgi secretory structures in osteoblasts and to discrete sites in early bone matrix. *J Histochem Cytochem* 1993;**41**:193–203.
 45. Ogbureke KUE, Fisher LW. Expression of SIBLINGs and their partner MMPs in salivary glands. *J Dent Res* 2004;**83**:664–670.
 46. Burstone MS. Histochemical observations on enzymatic processes in bones and teeth. *Ann NY Acad Sci* 1960;**85**:431.
 47. Paine ML, White SN, Luo W, Fong H, Sarikaya M, Snead ML. Regulated gene expression dictates enamel structure and tooth function. *Matrix Biol* 2001;**20**:273–292.
 48. Weiss MJ, Ray K, Henthorn PS, Lamb B, Kadesch T, Harris H. Structure of human liver/bone/kidney alkaline phosphatase gene. *J Biol Chem* 1988;**263**:120002–121010.

49. Kiefer MC, Bauer DM, Barr PJ. The cDNA and derived amino acid sequence for human osteopontin. *Nucleic Acids Res* 1989;**17**:3306.
50. Kiefer MC, Saphire AAC, Bauer DM, Barr PJ. The cDNA and derived amino acid sequences for human and bovine bone Gla protein. *Nucleic Acids Res* 1990;**18**:1909.
51. Fisher LW, McBride OW, Termine JD, Young MF. Human bone sialoprotein: deduced protein sequence and chromosomal localization. *J Biol Chem* 1990;**265**:2347–2351.
52. Ivanovski S, Li H, Haase HR, Bartold PM. Expression of bone associated macromolecules by gingival and periodontal ligament fibroblasts. *J Periodont Res* 2001;**36**:131–141.
53. Young MF, Kerr JM, Termine JD *et al*. cDNA cloning, mRNA distribution and heterogeneity, chromosomal location, and RFLP analysis of human osteopontin (OPN). *Genomics* 1990;**7**:491–502.
54. Turhani D, Cvikl B, Watzinger E *et al*. *In vitro* growth and differentiation of osteoblast-like cells on hydroxyapatite ceramic granule calcified from red algae. *J Oral Maxillofac Surg* 2005;**63**:793–799.
55. Onodera K, Takahashi I, Sasano Y *et al*. Stepwise mechanical stretching inhibits chondrogenesis through cell-matrix adhesion mediated by integrins in embryonic rat limb-bud mesenchymal cells. *Eur J Cell Biol* 2005;**84**:45–58.
56. Hamamoto Y, Nakajima T, Ozawa H, Uchida T. Production of amelogenin by enamel epithelium of Hertwig's root sheath. *Oral Surg Oral Med Oral Pathol* 1996;**81**:703–709.
57. Kumamoto H, Yoshida M, Ooya K. Immunohistochemical detection of amelogenin and cytokeratin 19 in epithelial odontogenic tumors. *Oral Dis* 2001;**7**:171–176.
58. Abiko Y, Murata M, Ito Y *et al*. Immunohistochemical localization of amelogenin in human odontogenic tumors, using a polyclonal antibody against bovine amelogenin. *Med Electron Microsc* 2001;**34**:185–189.
59. Shimokawa H, Sobel ME, Sasaki M, Termine JD, Young MF. Heterogeneity of amelogenin mRNA in the bovine tooth germ. *J Biol Chem* 1987;**25**:4042–4047.
60. Papagerakis P, MacDougal M, Hatton D, Bailleul-Forestier I, Oboeuf M, Berdal A. Expression of amelogenin in odontoblasts. *Bone* 2003;**32**:228–240.
61. Bosshardt DD, Nanci A. Immunocytochemical characterization of ectopic enamel deposits and cementicles in human teeth. *Eur J Oral Sci* 2003;**111**:51–59.
62. Brunette DM. Mechanical stretching increases the number of epithelial cells synthesizing DNA in culture. *J Cell Sci* 1984;**69**:35–45.
63. Saarialho-Kere U, Kovacs SO, Pentland AP, Olerud JE, Welgus HG, Parks WC. Cell-matrix interactions modulate interstitial collagenase expression by human keratinocytes activity involved wound healing. *J Clin Invest* 1993;**92**:2858–2866.
64. Salo T, Käkälä M, Kylmäniemi M, Autio-Harmainen H, Larjava H. Expression of matrix metalloproteinase-2 and -9 during early wound healing. *Lab Invest* 1994;**70**:176–182.
65. Ramakrishnan PR, Lin W-L, Sodek J, Cho MI. Synthesis of noncollagenous extracellular matrix proteins during development of mineralized nodules by rat periodontal ligament cells *in vitro*. *Calcif Tissue Int* 1995;**57**:52–59.
66. Haase HR, Ivanovski S, Waters MJ, Bartold PM. Growth hormone regulates osteogenic marker mRNA expression in human periodontal fibroblasts and alveolar bone-derived cells. *J Periodont Res* 2003;**38**:366–374.
67. Mauney JR, Jaquiere C, Volloch V, Heberer M, Martin I, Kaplan DL. *In vitro* and *in vivo* evaluation of differentially demineralized cancellous bone scaffolds combined with human bone marrow stromal cells for tissue engineering. *Bio-materials* 2005;**26**:3173–3185.
68. Nohutcu RM, McCauley LK, Shigeyama Y, Somerman MJ. Expression of mineral-associated proteins by periodontal ligament cells: *in vitro* vs. *in vivo*. *J Periodont Res* 1996;**31**:369–372.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.