

Prevalence of Xerostomia in Population-based Samples: A Systematic Review

M.F. Orellana, DDS, PhD; M.O. Lagravère, DDS, MSc; D.G.J. Boychuk, DDS; P.W. Major, DDS, MSc, FRCD(c); C. Flores-Mir, DDS, Cert Ortho, DSc

Abstract

The consequences of xerostomia on oral health have been studied for decades; however, the actual prevalence of this disorder on the general population remains controversial. The purpose of this systematic review was to determine the prevalence of xerostomia in population-based samples. Electronic databases were screened for relevant articles and reference lists of pertinent articles were also hand-searched. Thirteen articles meeting the final inclusion criteria were identified. Based on the definition of xerostomia used in this review, only publications including a subjective diagnosis of the disorder were included. All of the self-reported diagnoses were achieved through a questionnaire either by mail, telephone, interview or self-administered by the patient. There was a variation across papers regarding number and content of the questions as well as guidelines for the diagnosis of xerostomia (i.e. answer yes to one or more than one question). The prevalence of xerostomia in the selected articles ranged from 0.9% to 64.8%. The majority of these studies were performed in Scandinavia. In 9 of the articles, the studied samples were 50 years and older. None of the studies evaluated the prevalence xerostomia among individuals younger than 18 years. Based on these observations, it can be concluded that there is a need for population-based studies on prevalence of xerostomia in regions other than Scandinavia. A standardized protocol to diagnose xerostomia needs to be developed. Patient-perceived treatment needs and impact on quality of life should be included to have a complete picture of public health implications of the disease.

Key Words: xerostomia, prevalence, systematic review

Introduction

Controversy appears to exist regarding the definition of xerostomia, and other relevant terms such as dry mouth and hyposalivation have been used indistinctively in the literature. One commonly accepted clinical definition of xerostomia is the subjective sensation of dry mouth (1). The controversy regarding the meaning of xerostomia arises when dry mouth is indistinctly used for two conditions that are not always linked: xerostomia, defined as an individual subjective feeling of dry mouth and salivary gland hypofunction (SGH), a reduced unstimulated or stimulated salivary flow (2).

The presence of xerostomia may indicate that the salivary output is decreased or altered, placing patients at a higher risk for oral complications. Diverse symptoms and consequences have been associated with xerostomia. Symptoms such as halitosis, soreness, oral burning, and difficulty with swallowing and altered taste sensation have been linked with xerostomia (3). In more serious cases, a scalded sensation in the tongue, pharynx and esophagus has been reported. A significant increase in dental caries, periodontal diseases, and denture discomfort/loss of retention and oral infections like candidiasis might also be expected (4-8).

It is clear that xerostomia represents a serious problem with medical and dental implications. It has a damaging effect on the sufferer in particular and society in general. It affects people at an emotional and social level and deteriorates their overall quality of life (9). The devastating effects of xerostomia have been widely perceived by health professionals and researchers, as manifested by the extensive publications on risk factors, consequences and treatment of xerostomia (10-15).

Presumably, it would be the task of the dental professional to first recognize the symptoms and signs of this condition. Awareness of the prevalence of xerostomia in the general population is very important for the clinical practitioner. As stated by Nedefors, "If objective tests do not confirm the subjective complaints, often the patient's experience is that the symptoms are not taken seriously, many times even denied by the medical and dental professional" (16).

Inconsistency appears to exist in the prevalence of xerostomia reported in diverse populations. This is mostly due to pre-selection of the samples (17-20) and reports of xerostomia on patients affected by certain diseases and/or under pharmacological treatment (21-25). The aim of this systematic review is to consolidate current knowledge on the prevalence of xerostomia in population-based samples.

Send correspondence and reprint requests to: Carlos Flores-Mir, Director, Craniofacial & Oral Health Evidence Practice Group, Faculty of Medicine and Dentistry, Room 4051A, Dentistry/Pharmacy Centre, University of Alberta, Edmonton, Alberta, Canada T6G 2N8. Phone: (780) 492-7409; Fax: (780) 492-1624. E-mail address: carlosflores@ualberta.ca. Dr. Orellana is an MSc student in the Orthodontic Graduate Program, Department of Dentistry, University of Alberta, Edmonton, Alberta, Canada. Dr. Lagravère is a PhD student in the Orthodontic Graduate Program, Department of Dentistry, University of Alberta, Edmonton, Alberta, Canada. Dr. Boychuk is an MSc student in the Temporomandibular Disorder and Orofacial Pain Unit, Department of Dentistry, University of Alberta, Edmonton, Alberta, Canada. Dr. Major is a Professor and Director of the Orthodontic Graduate Program, Department of Dentistry, University of Alberta, Edmonton, Alberta, Canada. Dr. Flores-Mir is a Clinical Associate Professor, Orthodontic Graduate Program, Department of Dentistry, University of Alberta, Edmonton, Alberta, Canada. Manuscript received 3/5/05; accepted for publication 1/15/06.

Materials and Methods

Selection criteria. *Types of publication* – Only journal articles were included. Letters, editorials, theses, abstracts and other types of publications were excluded. *Types of studies* – Only observational studies (cohorts and cross-sectionals) were included. Clinical trials, case reports, case-control studies, studies of predictors of treatment outcome, and experimental laboratory studies were excluded. *Population* – Only population-based studies encompassing well-defined geopolitical boundaries were selected. Studies on other populations such as occupational groups, clinical populations, individuals living in retirement communities, nursing homes or any extended care facility, dental/medical practices, health centers or education colleges or any sample than could not be considered population-based were excluded. Studies evaluating convenience samples were also considered ineligible. Studies on individuals using medication on a chronic basis or with diagnosed medical conditions such as Sjogren's syndrome, cancer or any other chronic diseases were also excluded, as they could potentially skew the prevalence of xerostomia. *Disease definition* – Xerostomia, self reported dry mouth sensation (1). *Exposure* – Prevalence or frequency of xerostomia.

Search strategy. The selection and specific use of each term inside every database search were made with the help of a senior librarian specializing in health sciences database searches (Table 1). Briefly, epidemiological terms (prevalence, epidemiology, and population studies) were combined with diagnostic terms (xerostomia, dry mouth and hypo-salivation). Publications were retrieved by a computerized search of the following databases: MEDLINE (1966 to February week 3 of 2005); MEDLINE in process (February 28, 2005), Lilacs (February 2005) Pubmed (1966 to week 4 of February 2005), EMBASE (1996 to week 9, 2005), Web of Science (February 28, 2005), and all EBM reviews (Cochrane Database of Systematic Reviews, ASP Journal Club,

DARE and CCTR) (to the first Quarter of 2005) databases.

Eligibility of potential studies was determined by reading the title and abstract of each article identified by the different search engines. Based on the abstract information, three authors independently selected the articles to be retrieved. At this stage, any abstract that seemed to report xerostomia in a population-based population was selected. Any discrepancies were settled through discussion. When the abstract failed to provide sufficient information, a reprint of the full paper was obtained.

The final selection was independently completed by all the authors, followed by careful examination of each complete article. The selection criteria were fully applied at this stage. Reference lists of the selected articles were hand-searched for additional relevant publications that may have been missed in the database searches. In cases where specific data were necessary for the discussion, and were not specified in the article, efforts were made to contact the authors to obtain the required extra information.

Results

Forty-four abstracts satisfied the abstract inclusion criteria. After careful evaluation of the articles, thirteen (21, 26-37) were finally considered for this review (Table 2). A flow chart of the selection process can be found on Figure 1, with the inclusion or exclusion criteria described in the Methods Section above being used. Several articles were excluded because they did not satisfy the population criterion (18, 19, 38-60). Additional data on the diagnosis of dry mouth was required for three studies (37, 61, 62). Contact was attempted with the corresponding authors through electronic mail asking for the specific information. Only one of these authors (37) provided the required information. Since additional information was unavailable for the other two studies (61, 62), they were not included in the final discussion.

In some cases, the literature search retrieved more than one study report-

ing results from the same sample. Only the more detailed and completed report (28, 31, 35, 66) was considered to avoid duplication of results (9, 47, 63-65) which would potentially bias the conclusions.

All of the self-reported diagnoses were achieved through a questionnaire either by mail (21, 30, 34, 36), telephone (32, 33); in the form of an oral interview (26, 28, 29, 31); or written interview (27, 37). There was a variation across papers regarding number and content of the questions as well as guidelines for xerostomic diagnoses (i.e. answer yes to one or more than one question). Seven of the studies also performed an additional objective measurement of salivary gland function (21, 26, 27, 29, 31, 32, 35).

Six studies (46%) were conducted in Scandinavia. Of the 6 studies, 4 were performed in Sweden and 2 in Finland. Most of the data on the prevalence of xerostomia was retrieved from articles studying xerostomia (11 publications, 85%) (26, 27, 28, 29, 30, 31, 32, 33, 35, 55, 67). Two articles provided this information while evaluating the prevalence of oral mucosal lesions (34, 37) and two while assessing oral symptoms in menopause (21, 36).

All the studies evaluated the prevalence of xerostomia grouped by age and gender. The fact that the data were presented in different age group intervals made a direct comparison between studies difficult. In general, prevalence varied according to gender, age and sample origin. In 9 (69%) of the selected articles (21, 26, 27, 28, 29, 31, 33, 36, 37), the population sample consisted of individuals 50 years and older. None of the studies evaluated the prevalence of xerostomia in individuals younger than 18 years old. One of the highest rates of xerostomia in older adults was reported in Helsinki, Finland (29) in 1994 where 46% of 75-, 80-, and 85-year-old individuals complained of dry mouth. The lowest rate was a 0.9% in older German individuals (34).

All the selected articles reported a greater prevalence of xerostomia in

TABLE I
Database search strategy and sensitivity of the electronic databases used

Database	Keywords	Results	Selected	% or total selected abstracts (13)*
PubMed	1) xerostomia (2) hyposalivation; (3) dry mouth; (4) 1 or 2 or 3; (5) prevalenc*; (6) epidemiol*; (7) population stud*; (8) 5 or 6 or 7; (9) 4 and 8; (10) limit human	855	9	69.2%
Medline	(1) xerostomia.mp (2) hyposalivation.mp; (3) dry mouth.mp (4) 1 or 2 or 3; (5) prevalenc\$.mp; (6) epidemiol\$.mp; (7) population stud\$.mp; (8) 5 or 6 or 7; (9) 4 and 8; (10) limit human	254	10	76.9%
Medline In-Process & Other Non-Indexed Citations	(1) xerostomia.mp (2) hyposalivation.mp; (3) dry mouth.mp (4) 1 or 2 or 3; (5) prevalenc\$.mp; (6) epidemiol\$.mp; (7) population stud\$.mp; (8) 5 or 6 or 7; (9) 4 and 8; (10) limit human	10	0	0
Web of Science	TS=(xerost* OR hyposalivation OR dry mouth*) AND TS=(epidemiol* OR prevalenc* or population stud*) DocType=Article; Language=All languages; Database(s)=SCI-EXPANDED.	245	10	76.9%
Embase	(1) xerostomia.mp (2) hyposalivation.mp; (3) dry mouth.mp (4) 1 or 2 or 3; (5) prevalenc\$.mp; (6) epidemiol\$.mp; (7) population stud\$.mp; (8) 5 or 6 or 7; (9) 4 and 8	439	2	15.4%
All EBM reviews	(1) xerostomia.mp (2) hyposalivation.mp; (3) dry mouth.mp (4) 1 or 2 or 3; (5) prevalenc\$.mp; (6) epidemiol\$.mp; (7) population stud\$.mp; (8) 5 or 6 or 7; (9) 4 and 8; (10) limit human	73	1	7.7%
Lilacs	xerostomia AND prevalence	2	0	0

* Percentages do not add up to 100% as the same reference could be found in several databases

females. One study (36) reported prevalence in a female sample only.

All but one (37) of the selected articles was in English.

The diversity of sample size (ranging from 259 to 3,313 individuals) and differences in the assessment of xerostomia, as well as in the result presentation, made these articles unsuitable for a meta-analysis. Therefore, the results presented here are merely descriptive.

Discussion

This systematic review evaluated the results of 13 studies on the prevalence of xerostomia in population-based samples. An electronic literature search was conducted using 6 databases to gather articles in all languages. None of the databases included more than three quarters of the total selected abstracts; therefore, doing a Medline search alone in a systematic review of this topic was not sufficient to reveal all the related abstracts. Every effort was made to locate published population-based reports on prevalence of xerostomia.

Although xerostomia and hyposalivation refer to two different entities, as was discussed in the introduction to this review, hyposalivation was included as one of the search terms in an attempt to prevent the omission of relevant information. Significant disagreement in the use of these terms is evident in the leading search engines (MEDLINE and PubMed). Nederfors *et al.* (16) noted that the database MEDLINE, under the MESH heading, refers to xerostomia as "decreased salivary flow". Likewise, when the term "hyposalivation" is entered, the reference "see xerostomia" is made. The National Library of Medicine database (PubMed) presents the same limitation observed in MEDLINE. The definition of xerostomia in epidemiological studies was extremely diverse. There is a lack of consensus in the literature regarding the definition and diagnosis of xerostomia.

The definition of xerostomia as the sensation of dry mouth that could exist in the presence of a normal or abnormal salivary flow and rate is sup-

ported by Narhi *et al.* (29), who reported no significant differences in salivary flow rates between individuals complaining of oral dryness and controls. Furthermore, only 5.7% of a sample population of 684 individual had low salivary flow rates and xerostomia. (33) Nevertheless, a decrease in salivary flow may be considered the most common etiologic factor in xerostomia (68-70). Complementary diagnostic procedures such as salivary flow rate and oral examination have been completed in several studies even though xerostomia might not correlate with actual physical evidence of oral dryness. Additionally those objective tests which do not confirm patients' subjective complaints could potentially result in patients perceiving that their symptoms are not taken seriously (16). Therefore, the development of tests considering the patient-perceived need for xerostomia treatment is required.

The majority of epidemiological studies located in this review have been conducted in Scandinavia, par-

TABLE 2
Description of xerostomia prevalence in the general population

Author	Country	Total	Range	Sample Prevalence		Assessment of Xerostomia	
				Age Gender	%	Subjective	Objective
Osterberg (26)	Sweden	968	70	Female Male	25 16	Interview. One question.	Unstimulated and stimulated whole saliva flow rate
Nederfors (30)	Sweden	3313	20-80	Female Male	28.3 23.1	Self-administered questionnaire. Does your mouth usually feel dry? and dryness related symptoms and behavior.	N/A
Jansson (36)	Sweden	1159	53-54	Female	16	Self-administered questionnaire. Different extragenital Symptoms.	N/A
Bergdahl (35)	Sweden	1427	20-69	Female Male	28.2 14.9	Self-administered questionnaire.	Unstimulated and stimulated whole saliva flow rate
Narhi (29)	Finland	341	75-85	Female Male	46* Total sample	Personal Interview. Yes or No answers to dry mouth related symptoms.	Unstimulated and stimulated whole saliva flow rate.
Anttila (21)	Finland	780	55	Female Male	33.3 25.8	Self-administered questionnaire. plus personal interview	Unstimulated and stimulated whole saliva flow rate
Reichart (34)	Germany	2023	65-74	Female Male	1.2 0.4	Personal Interview according to WHO** and ICD-DA***	N/A
Pujol (32)	Spain	268	18-over 65	Female Male	13 6.2	Telephone Interview.	N/A
Locker (28)	Canada	907	50-over 65	Female Male	20.7 13.8	Personal Interview as part of inventory in oral symptoms and complaints.	N/A
Hochberg (31)	USA	2482	65-84	Female Male	20.1 13.2	Personal Interview. Does your mouth usually feel dry? Do you wake up at night feeling so dry in your mouth that you need to drink fluid?	Unstimulated and stimulated whole saliva production by modified saxon test.
Thomson (33)	Australia	684	65-100	Female Male	24 17	Computer assisted telephone interview.	Unstimulated whole saliva flow rate
Ben-Aryeh (27)	Israel	259	over 60	Female Male	27.7* Total Sample	Self-administered questionnaire containing a short explanation of xerostomia	Unstimulated and stimulated whole saliva flow rate
Espinoza (37)	Chile	889	65-over 75	Female Male	44* total sample	Personal Interview according to WHO.	N/A

* In these studies, prevalence of xerostomia was not divided by gender.

** WHO: Recording of data based on World Health Organization.

*** ICD-DA: International Classification of Diseases in Dentistry


ticularly Sweden and Finland. Prevalence data from one country may not be characteristic of the global population since unique cultural and etiological factors may play a role. Therefore, further studies are needed based on populations other than Scandinavian.

The studies presented here show a very wide range in the prevalence of xerostomia, from 0.9% (34) to 46% (29). This variation between reports might be explained as a consequence of differences in the diagnostic pro-

cess. It is interesting to note that if the study (34) with the lowest prevalence is not considered, the prevalence values are less dispersed (20-46% for females and 13-26% for males). Again, the diagnostic process may be a significant reason for this discrepancy. Being a subjective sensation, it is personal and varies depending on the individual reporting the symptom as well as manner in which the questions are worded. As suggested by Sreebny, asking "Do you suffer from dryness in the morning or evening?" yields one

kind of answer and asking, "Do you suffer from continuous dryness?" produces a different one (71). Also, a "reporting behavior" or the way in which people report symptoms in general may have an impact. Underlying physical/medical conditions, medications being taken, and exposure to risk factors could also potentially affect the reported prevalence. This should not be the case in the present systematic review because these conditions were considered as exclusion criteria. Development of

FIGURE 1
Flow diagram of the literature search



uniform protocols for collecting and reporting the data would be very important in studying the worldwide prevalence of xerostomia.

All of the articles in the present review also demonstrated increasing prevalence of subjective perception of dry mouth as people age. Pujol *et al.* (32) reported a difference of more than 10 points between the 35- to 64-year-old group (7.3%) and the over-65 group (18.2%). Most of the articles evaluated the prevalence of subjective dry mouth on older populations (50 years and older). This condition is believed to be fairly common among older adults, and this may explain the large number of studies focusing on this population. On the other hand, there are very limited data on prevalence of xerostomia in children. Only 4 studies on children were identified. (72-75). As all of them only measured salivary flow rates, these results were not included in the present review. Due to the shortage of population-based studies on the prevalence of xerostomia in younger groups, comments and/or conclusions about implications of xerostomia in adolescents and children cannot be drawn

from this review. The need for more population-based studies about xerostomia in younger adults and especially teenagers and children is somewhat controversial. Although there is no solid evidence of younger non-medically compromised populations being affected by xerostomia, it is important to acknowledge that most of the reported data are from individuals 50 years and older. The scarceness of reports on younger populations makes it difficult to conclude that xerostomia increases with age.

Almost all of the selected publications discriminated their sample by gender; only one study (36) involved only females. Although all the studies did not report the actual prevalence of xerostomia in males and females, those reporting showed greater rates in woman than men. Bergdahl *et al.* (35) reported a difference of 13 points between genders. On average, female patients reported an 8% higher prevalence of subjective dry mouth. Based on these reports, it is reasonable to believe that the differences in prevalence between genders are real and that women have a higher prevalence of xerostomia than men.

Conclusions

- ♦ A standardized definition and protocol to diagnose xerostomia needs to be developed to facilitate comparison between studies and communication between researchers.

- ♦ Studies evaluating younger age groups should be conducted to understand the impact throughout the lifespan.

- ♦ Most of the current literature focuses on prevalence of xerostomia in the Scandinavian population. More population-based studies in other regions are needed.

Acknowledgments

Special thanks to Linda Seale for her professional assistance in the database search.

References

1. Neville BW, Damm DDA, C.M., Bouquet JE. Oral & Maxillofacial Pathology. 2nd edition ed. Anonymous Philadelphia, PA: W.B. Saunders; 2002.
2. Fox PC, van der Ven PF, Sonies BC, Weiffenbach JM, Baum BJ. Xerostomia: evaluation of a symptom with increasing significance. J Am Dent Assoc. 1985 Apr;110(4):519-25.
3. Nally F. Dry mouth and halitosis. Practitioner. 1990 Jun 8;234(1490):603.
4. Narhi TO, Meurman JH, Ainamo A. Xerostomia and hyposalivation: causes, consequences and treatment in the elderly. Drugs Aging. 1999 Aug;15(2):103-16.
5. Billings RJ. An epidemiologic perspective of saliva flow rates as indicators of susceptibility to oral disease. Crit Rev Oral Biol Med. 1993;4(3-4):351-6.
6. Anusavice KJ. Dental caries: risk assessment and treatment solutions for an elderly population. Compend Contin Educ Dent. 2002 Oct; 23 (10 Suppl): 12-20.
7. Niedermeier WH, Kramer R. Salivary secretion and denture retention. J Prosthet Dent. 1992 Feb;67(2):211-6.
8. Papas AS, Joshi A, MacDonald SL, Maravelis-Splagounias L, Pretara-Spanedda P, Curro FA. Caries prevalence in xerostomic individuals. J Can Dent Assoc. 1993 Feb;59(2):171,4, 177-9.
9. Locker D. Xerostomia in older adults: a longitudinal study. Gerodontology. 1995 Jul;12(1):18-25.
10. Crockett DN. Xerostomia: the missing diagnosis? Aust Dent J. 1993 Apr;38(2):114-8.
11. Astor FC, Hanft KL, Ciocon JO. Xerostomia: a prevalent condition in the elderly. Ear Nose Throat J. 1999 Jul;78(7):476-9.

12. Guggenheimer J, Moore PA. Xerostomia: etiology, recognition and treatment. *J Am Dent Assoc.* 2003 Jan;134(1):61-9; quiz 118-9.
13. Fox PC. Acquired salivary dysfunction. Drugs and radiation. *Ann N Y Acad Sci.* 1998 Apr 15;842:132-7.
14. Navazesh M, ADA Council on Scientific Affairs and Division of Science. How can oral health care providers determine if patients have dry mouth? *J Am Dent Assoc.* 2003 May;134(5):613-20; quiz 633.
15. Ship JA, Pillemer SR, Baum BJ. Xerostomia and the geriatric patient. *J Am Geriatr Soc.* 2002 Mar;50(3):535-43.
16. Nederfors T. Xerostomia and hyposalivation. *Adv Dent Res.* 2000 Dec;14:48-56.
17. O'Grady NP. Incidence of dry mouth complaint in Cork Dental Hospital patient population. *Stoma (Lisb).* 1990 Aug-Oct;2(17):55-6, 58.
18. Gilbert GH, Heft MW, Duncan RP. Mouth dryness as reported by older Floridians. *Community Dent Oral Epidemiol.* 1993 Dec;21(6):390-7.
19. Billings RJ, Proskin HM, Moss ME. Xerostomia and associated factors in a community-dwelling adult population. *Community Dent Oral Epidemiol.* 1996 Oct;24(5):312-6.
20. Locker D. Dental status, xerostomia and the oral health-related quality of life of an elderly institutionalized population. *Spec Care Dentist.* 2003;23(3):86-93.
21. Anttila SS, Knuuttila ML, Sakki TK. Depressive symptoms as an underlying factor of the sensation of dry mouth. *Psychosom Med.* 1998 Mar-Apr;60(2):215-8.
22. Coll J, Anglada J, Tomas S, Reth P, Goday A, Millan M, et al. High prevalence of subclinical Sjogren's syndrome features in patients with autoimmune thyroid disease. *J Rheumatol.* 1997 Sep;24(9):1719-24.
23. Bergdahl M, Bergdahl J. Low unstimulated salivary flow and subjective oral dryness: association with medication, anxiety, depression, and stress. *J Dent Res.* 2000 Sep;79(9):1652-8.
24. Jensen SB, Pedersen AM, Reibel J, Nauntofte B. Xerostomia and hypofunction of the salivary glands in cancer therapy. *Support Care Cancer.* 2003 Apr;11(4):207-.
25. Thomas E, Hay EM, Hajeer A, Silman AJ. Sjogren's syndrome: a community-based study of prevalence and impact. *Br J Rheumatol.* 1998 Oct;37(10):1069-76.
26. Osterberg T, Landahl S, Hedegard B. Salivary flow, saliva, pH and buffering capacity in 70-year-old men and women. Correlation to dental health, dryness in the mouth, disease and drug treatment. *J Oral Rehabil.* 1984 Mar;11(2):157-70.
27. Ben-Aryeh H, Miron D, Berdicevsky I, Szargel R, Gutman D. Xerostomia in the elderly: prevalence, diagnosis, complications and treatment. *Gerodontology.* 1985 Winter;4(2):77-82.
28. Locker D. Subjective reports of oral dryness in an older adult population. *Community Dent Oral Epidemiol.* 1993 Jun;21(3):165-8.
29. Narhi TO. Prevalence of subjective feelings of dry mouth in the elderly. *J Dent Res.* 1994 Jan;73(1):20-5.
30. Nederfors T, Isaksson R, Mornstad H, Dahlof C. Prevalence of perceived symptoms of dry mouth in an adult Swedish population—relation to age, sex and pharmacotherapy. *Community Dent Oral Epidemiol.* 1997 Jun;25(3):211-6.
31. Hochberg MC, Tielsch J, Munoz B, Bandeen-Roche K, West SK, Schein OD. Prevalence of symptoms of dry mouth and their relationship to saliva production in community dwelling elderly: the SEE project. *Salisbury Eye Evaluation. J Rheumatol.* 1998 Mar;25(3):486-91.
32. Pujol T, Coma M, Pujol M, Postigo P. [Prevalence of xerostomia in the general population]. *Aten Primaria.* 1998 Mar 15;21(4):225-8. Spanish
33. Thomson WM, Chalmers JM, Spencer AJ, Williams SM. The Xerostomia Inventory: a multi-item approach to measuring dry mouth. *Community Dent Health.* 1999 Mar;16(1):12-7.
34. Reichart PA. Oral mucosal lesions in a representative cross-sectional study of aging Germans. *Community Dent Oral Epidemiol.* 2000 Oct;28(5):390-8.
35. Bergdahl M. Salivary flow and oral complaints in adult dental patients. *Community Dent Oral Epidemiol.* 2000 Feb;28(1):59-66.
36. Jansson C, Johansson S, Lindh-Astrand L, Hoffmann M, Hammar M. The prevalence of symptoms possibly related to the climacteric in pre- and postmenopausal women in Linköping, Sweden. *Maturitas.* 2003 Jun 30;45(2):129-.
37. Espinoza I, Rojas R, Aranda W, Gamonal J. Prevalence of oral mucosal lesions in elderly people in Santiago, Chile. *J Oral Pathol Med.* 2003 Nov;32(10):571-5.
38. Mersel A. Oral health status and dental needs in a geriatric institutionalized population in Paris. *Gerodontology.* 1989 Summer;8(2):47-51.
39. Handelman SL, Baric JM, Saunders RH, Espeland MA. Hyposalivatory drug use, whole stimulated salivary flow, and mouth dryness in older, long-term care residents. *Spec Care Dentist.* 1989 Jan-Feb;9(1):12-8.
40. Johnson G, Barenthin I, Westphal P. Mouth dryness among patients in longterm hospitals. *Gerodontology.* 1984 Autumn;3(3):197-203.
41. Loesche WJ, Abrams J, Terpenning MS, Bretz WA, Dominguez BL, Grossman NS, et al. Dental findings in geriatric populations with diverse medical backgrounds. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1995 Jul;80(1):43-54.
42. Locker D, Matear D, Stephens M, Lawrence H, Payne B. Comparison of the GOHAI and OHIP-14 as measures of the oral health-related quality of life of the elderly. *Community Dent Oral Epidemiol.* 2001 Oct;29(5):373-81.
43. Loesche WJ, Bromberg J, Terpenning MS, Bretz WA, Dominguez BL, Grossman NS, et al. Xerostomia, xerogenic medications and food avoidances in selected geriatric groups. *J Am Geriatr Soc.* 1995 Apr;43(4):401-7.
44. Rhodus NL, Brown J. The association of xerostomia and inadequate intake in older adults. *J Am Diet Assoc.* 1990 Dec;90(12):1688-92.
45. Sreebny LM, Valdin A. Xerostomia. Part I: Relationship to other oral symptoms and salivary gland hypofunction. *Oral Surg Oral Med Oral Pathol.* 1988 Oct;66(4):451-8.
46. Sreebny LM, Valdin A, Yu A. Xerostomia. Part II: Relationship to nonoral symptoms, drugs, and diseases. *Oral Surg Oral Med Oral Pathol.* 1989 Oct;68(4):419-27.
47. Thomson WM, Brown RH, Williams SM. Medication and perception of dry mouth in a population of institutionalised elderly people. *N Z Med J.* 1993 Jun 9;106(957):219-21.
48. van der Putten GJ, Brand HS, Bots CP, van Nieuw Amerongen A. [Prevalence of xerostomia and hyposalivation in the nursing home and the relation with number of prescribed medication]. *Tijdschr Gerontol Geriatr.* 2003 Feb;34(1):30-6. Dutch
49. Wyatt CC. Elderly Canadians residing in long-term care hospitals: Part I. Medical and dental status. *J Can Dent Assoc.* 2002 Jun;68(6):353-8.
50. Tarkkila L, Linna M, Tiitinen A, Lindqvist C, Meurman JH. Oral symptoms at menopause—the role of hormone replacement therapy. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2001 Sep;92(3):276-80.
51. Field EA, Fear S, Higham SM, Ireland RS, Rostron J, Willetts RM, et al. Age and medication are significant risk factors for xerostomia in an English population, attending general dental practice. *Gerodontology.* 2001 Jul;18(1):21-4.
52. Trombelli L, Mandrioli S, Zangari F, Saletti C, Calura G. [Oral symptoms in the climacteric]. A prevalence study. *Minerva Stomatol.* 1992 Nov; 41(11): 507-13. Italian
53. Pemberton MN, Thornhill MH. British general practitioners' experience and knowledge of xerostomia - A retrospective survey. *Int J Health Promotion Education.* 2001;39:86-7.
54. Ikebe K, Nokubi T, Sajima H, Kobayashi S, Hata K, Ono T, et al. Perception of dry mouth in a sample of community-dwelling older adults in Japan. *Spec Care Dentist.* 2001 Mar-Apr;21(2):52-9.

55. Fox PC, Busch KA, Baum BJ. Subjective reports of xerostomia and objective measures of salivary gland performance. *J Am Dent Assoc.* 1987 Oct;115(4):581-4.
56. Navazesh M, Mulligan R, Komaroff E, Redford M, Greenspan D, Phelan J. The prevalence of xerostomia and salivary gland hypofunction in a cohort of HIV-positive and at-risk women. *J Dent Res.* 2000 Jul;79(7):1502-7.
57. Thomson WM, Williams SM. Further testing of the xerostomia inventory. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2000 Jan;89(1):46-50.
58. Almas K, Al-Hawish A, Al-Khamis W. Oral hygiene practices, smoking habit, and self-perceived oral malodor among dental students. *J Contemp Dent Pract.* 2003 Nov 15;4(4):77-90.
59. Persson RE, Persson GR, Kiyak HA, Powell LV. Oral health and medical status in dentate low-income older persons. *Spec Care Dentist.* 1998 Mar-Apr;18(2):70-7.
60. Sandberg GE, Sundberg HE, Fjellstrom CA, Wikblad KF. Type 2 diabetes and oral health: a comparison between diabetic and non-diabetic subjects. *Diabetes Res Clin Pract.* 2000 Sep;50(1):27-34.
61. Jacobsson LT, Axell TE, Hansen BU, Henricsson VJ, Larsson A, Lieberkind K, et al. Dry eyes or mouth—an epidemiological study in Swedish adults, with special reference to primary Sjogren's syndrome. *J Autoimmun.* 1989 Aug;2(4):521-7.
62. Hakeberg M, Berggren U, Hagglin C, Ahlqvist M. Reported burning mouth symptoms among middle-aged and elderly women. *Eur J Oral Sci.* 1997 Dec;105(6):539-43.
63. Bergdahl M, Bergdahl J. Burning mouth syndrome: prevalence and associated factors. *J Oral Pathol Med.* 1999 Sep;28(8):350-4.
64. Bergdahl M, Bergdahl J. Perceived taste disturbance in adults: prevalence and association with oral and psychological factors and medication. *Clin Oral Investig.* 2002 Sep;6(3):145-9.
65. Schein OD, Hochberg MC, Munoz B, Tielsch JM, Bandeen-Roche K, Provost T, et al. Dry eye and dry mouth in the elderly: a population-based assessment. *Arch Intern Med.* 1999 Jun 28;159(12):1359-63.
66. Thomson WM, Chalmers JM, Spencer AJ, Ketabi M. The occurrence of xerostomia and salivary gland hypofunction in a population-based sample of older South Australians. *Spec Care Dentist.* 1999 Jan-Feb;19(1):20-3.
67. Sreebny LM, Valdin A. Xerostomia. A neglected symptom. *Arch Intern Med.* 1987 Jul;147(7):1333-7.
68. Sreebny LM. Xerostomia: diagnosis, management and clinical complications. In: Edgar WM, O'Mullane DM, editors. *Saliva and oral health.* 2nd edition ed. London, England: British Dental Association; 1996.
69. Sreebny LM, Schwartz SS. A reference guide to drugs and dry mouth—2nd edition. *Gerodontology.* 1997 Jul;14(1):33-47.
70. Wolff M, Kleinberg I. Oral mucosal wetness in hypo- and normosalivators. *Arch Oral Biol.* 1998 Jun;43(6):455-62.
71. Sreebny LM. Saliva in health and disease: an appraisal and update. *Int Dent J.* 2000 Jun;50(3):140-61.
72. Andersson R. The flow rate, pH and buffer effect of mixed saliva in school-children. *Odontol Revy.* 1972;23(4):421-8.
73. Bretz WA, do Valle EV, Jacobson JJ, Marchi F, Mendes S, Nor JE, et al. Unstimulated salivary flow rates of young children. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2001 May;91(5):541-5.
74. Watanabe S, Dawes C. Salivary flow rates and salivary film thickness in five-year-old children. *J Dent Res.* 1990 May;69(5):1150-3.
75. Watanabe S, Ohnishi M, Imai K, Kawano E, Igarashi S. Estimation of the total saliva volume produced per day in five-year-old children. *Arch Oral Biol.* 1995 Aug;40(8):781-2.

Copyright of Journal of Public Health Dentistry is the property of American Association of Public Health Dentistry and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.