

# Comparative genomic hybridization and transcriptome analysis with a pan-genome microarray reveal distinctions between JP2 and non-JP2 genotypes of *Aggregatibacter actinomycetemcomitans*

Y. Huang<sup>1</sup>, W. Kittichotirat<sup>2,3</sup>, M.P.A. Mayer<sup>4</sup>, R. Hall<sup>2</sup>, R. Bumgarner<sup>2</sup> and C. Chen<sup>1</sup>

1 Division of Periodontology, Diagnostic Sciences and Dental Hygiene, Ostrow School of Dentistry of USC, Los Angeles, CA, USA

2 Department of Microbiology, University of Washington, Seattle, WA, USA

3 Systems Biology and Bioinformatics Research Group, Pilot Plant Development and Training Institute, King Mongkut's University of Technology Thonburi, Bangkhuntien, Bangkok, Thailand

4 Departament of Microbiology, Institute of Biomedical Sciences, Universidade de São Paulo, São Paulo, Brazil

**Correspondence:** Casey Chen, Division of Periodontology, Diagnostic Sciences and Dental Hygiene, Herman Ostrow School of Dentistry of USC, 925 W 34th Street, Room 4107, Los Angeles, CA 90089-0641, USA Tel.: +1 213 740 1075; fax: +1 213 740 6778; E-mail: ccchen@hsc.usc.edu

Keywords: aggressive periodontitis; gene expression; genomic islands; horizontal gene transfer Accepted 30 August 2012 DOI: 10.1111/omi.12005

## SUMMARY

It was postulated that the highly virulent JP2 genotype of Aggregatibacter actinomycetemcomitans may possess a constellation of distinct virulence determinants not found in non-JP2 genotypes. This study compared the genome content and the transcriptome of the serotype b JP2 genotype and the closely related serotype b non-JP2 genotype of A. actinomycetemcomitans. A custom-designed pan-genomic microarray of A. actinomycetemcomitans was constructed and validated against a panel of 11 sequenced reference strains. The microarray was subsequently used for comparative genomic hybridization of serotype b strains of JP2 (six strains) and non-JP2 (six strains) genotypes, and for transcriptome analysis of strains of JP2 (three strains) and non-JP2 (two strains). Two JP2-specific and two non-JP2-specific genomic islands were identified. In one instance, distinct genomic islands were found to be inserted into the same locus among strains of different genotypes. Transcriptome analysis identified five operons, including the leukotoxin operon, to have at least two genes with an expression ratio of 2 or greater between genotypes. Two of the differentially expressed operons were members of the membrane-bound nitrate reductase system (*nap* operon) and the Tol-Pal system of gram-negative bacterial species. This study is the first to demonstrate the differences in the full genome content and gene expression between *A. actinomycetemcomitans* strains of JP2 and non-JP2 genotypes. The information is essential for designing hypothesis-driven experiments to examine the pathogenic mechanisms of *A. actinomycetemcomitans*.

# INTRODUCTION

The gram-negative facultative species *Aggregatibacter actinomycetemcomitans* is a major etiologic agent of localized aggressive periodontitis and other forms of

periodontitis, but can also be found in the oral cavity of healthy individuals (Slots, 1999; Chen *et al.*, 2010b). The species comprises discrete clonal lineages represented by different serotypes (Kaplan *et al.*, 2002; Rylev & Kilian, 2008). Substantial variation in genomic content and arrangement has been found among strains (Kittichotirat *et al.*, 2010, 2011). For example, the gene content of a given strain may differ as much as 19.5% from another strain (Kittichotirat *et al.*, 2011). The patterns of genomic variation among strains correlate with the major clonal lineages of *A. actinomycetemcomitans*, suggesting that much of the variation was fixed in the populations long ago.

Strains of A. actinomycetemcomitans can be distinguished based on a variety of genotyping methods such as arbitrarily primed polymerase chain reaction (PCR) and restriction fragment length polymorphism (DiRienzo et al., 1994a; Asikainen et al., 1995; Paju et al., 2000). These typing methods were commonly used to examine the association of specific genotypes of A. actinomycetemcomitans with distinct periodontal disease conditions, leading to substantial evidence for variable virulence potentials among strains. For example, serotype b strains with a 530base-pair (bp) deletion in the promoter region of the leukotoxin operon (designated as JP2 clone) were found to be associated with aggressive periodontitis in patients of African descent (Haubek et al., 1997). In prospective studies, infection by strains of the JP2 clone posed a greater risk for aggressive periodontitis or conversion from periodontal health to disease than non-JP2 clones (DiRienzo et al., 1994b; Haubek et al., 2008). Although it is tempting to attribute the high virulence of the JP2 clone to its high leukotoxicity, there is no clear evidence that this is the only determinant. It is possible that other virulence determinants also play a role in the pathogenesis of periodontal infection by the JP2 clone.

In the literature, the term 'non-JP2' has been loosely applied to all strains that do not have the characteristic 530-bp deletion in the promoter of leukotoxin operon, irrespective of their serotypes. However, it is clear that strains of different serotypes represent genetically distinct groups (Kittichotirat *et al.*, 2011). To focus on JP2-specific genomic features, we here use the terms 'JP2 genotype' and 'non-JP2 genotype' to refer to serotype b strains that are distinguished by the structure of the promoter of leukotoxin operon. This study tested the hypothesis that JP2 and non-JP2 genotypes differ in their genome content and gene expression patterns. A customized pan-genome microarray designed based on genome sequences of 18 strains of *A. actinomycetemcomitans* was used to compare the genome content and the transcriptomes of strains with JP2 and non-JP2 genotypes. The results have helped to narrow the field of interest to a few, but potentially important, differences between the genomes and expression patterns of JP2 and non-JP2 strains. The information we have obtained is essential for hypothesis-driven experimental design to dissect the virulence mechanisms of *A. actinomycetemcomitans*.

# METHODS

## Bacterial strains and genomic DNA isolation

The A. actinomycetemcomitans strains were grown on tryptic soy agar plates with 0.6% yeast extract for 2 days at 37°C in an atmosphere supplemented with 5% CO<sub>2</sub>, and were harvested by washing the bacteria off the plates with phosphate-buffered saline. Genomic DNA was isolated using the Qiagen DNeasy Blood & Tissue Kit (Qiagen, Valencia, CA) according to the manufacturer's manual. Serotypes were determined by PCR as previously described (Kaplan et al., 2002). JP2 and non-JP2 classification was determined by PCR analysis of the Itx promoter using the 5'-TCCATATTAAATCTCCTTGT-3' primers and 5'-AACCTGATAACAGTATT-3' (Brogan et al., 1994) to detect the characteristic 530-bp deletion. Serotype b strains with the characteristic 530-bp deletion were designated JP2 genotype, in contrast to strains of non-JP2 genotype that had the long Itx promoter. Of the 14 previously sequenced strains 11 were used for comparative genomic hybridization (CGH) by microarray as part of the validation process. Subsequently, CGH was performed for eight additional serotype b strains to examine their genome content (see Table 1 for a summary of strains).

# Pan-genome microarray of *A. actinomycetemcomitans*

Information from 14 previously examined genomes of *A. actinomycetemcomitans* (Kittichotirat *et al.*, 2011) and four strains that were sequenced *de novo* (see

Strain	Genome	Race/ethnicity	Age	Periodontal diagnosis <sup>1</sup>	Serotype/ genotype <sup>2</sup>	GenBank accession number
123C	Draft	Caucasian	48	CP	b/non-JP2	AEJQ0000000
SCC1398	Draft	Caucasian	25	LAP	b/non-JP2	AEJP0000000
ANH9381	Complete	Caucasian	NA	Н	b/non-JP2	CP003099
ATCC29524	N/A	NA	NA	NA	b/non-JP2	N/A
194	N/A	NA	NA	AP	b/non-JP2	N/A
G104-2	N/A	NA	28	CP	b/non-JP2	N/A
S067	N/A	NA	NA	AP	b/JP2	N/A
A26	N/A	African	19	AP	b/JP2	N/A
G111-1	N/A	Caucasian	21	AP	b/JP2	N/A
G121-2	N/A	African	38	AP	b/JP2	N/A
D28S-1	N/A	African-American	21	LAP	b/JP2	N/A
D41S-1	N/A	Hispanic	9	AP	b/JP2	N/A
HK1651	Complete	African	18	LAP	b/JP2	N/A
D7S-1	Complete	African-American	29	GAP	а	CP003496
D11S-1	Complete	African-American	16	GAP	С	CP001733
163B	Draft	Caucasian	50	Н	d	AEJL0000000
SCC393	Draft	Caucasian	40	CP	е	AEJN0000000
D18P-1	Draft	Asian-American	20	GAP	f	AEJO0000000
SCC2302	Draft	Caucasian	33	G	С	AEJR0000000
D17P-2	Draft	Asian-American	24	LAP	С	ADOB0000000

 
 Table 1 Aggregatibacter actinomycetemcomitans strains used for comparative genomic hybridization and transcriptome analysis with the pangenome microarray

<sup>1</sup>CP, chronic periodontitis; LAP, localized aggressive periodontitis; AP, aggressive periodontitis; GAP, generalized aggressive periodontitis; H, healthy; G, gingivitis.

<sup>2</sup>Genotype was based on the analysis of the promoter structure of leukotoxin operon by polymerase chain reaction. The promoter structure in strains other than serotype b was confirmed to be identical to that found in strains of non-JP2 genotype.

Supplementary material Table S1 for sequencing results) was used to design a pan-genome microarray. The de novo sequenced strains S23A Gen-Bank accession number AJMH0000000), SCC4092 (AJMF0000000), AAS4A (AJMG0000000) and A160 (AJME0000000) were recovered from the same individuals as the previously studied strains 123C, SCC1398, SCC2302 and SCC393, respectively. Genomic variation of strains within individuals is a subject of another study and will not be described here. The 18 genomes were searched for genes and the genes were grouped based on sequence homology to identify unique gene clusters as described previously (Kittichotirat et al., 2011). A total of 42,668 predicted genes were grouped into 3426 homologous gene clusters and the result can be found at http:// expression.washington.edu/genetable/script/gene table viewer. This set of 3426 homologous gene clusters represents the pan-genome of A. actinomycetemcomitans, which is much larger than the total number of genes in any single genome of this species. In the text below, we refer to each gene in the A. actino*mycetemcomitans* pan-genome by using the p-cluster ID as shown in our homologous gene cluster result.

The longest sequence in each gene cluster was selected as a representative sequence and used for microarray probe design. A total of 13,960 probes (60 bp in length) for 3121 homologous gene clusters were initially generated using EARRAY (https://earray. chem.agilent.com/earray/). The tool ARRAYOLIGOSELEC-TOR (Bozdech et al., 2003) was then used to design 20 probes for five additional clusters. The remaining 300 gene clusters, for which no probes were generated by either software, were often (71%) small genes (<300 bp) with no known function (annotated as hypothetical protein). All the designed probes were then synthesized in situ by Agilent Technologies (Santa Clara, CA) to create an  $8 \times 15$ K microarray with the randomized feature layout option selected. As this array format contains more than 15K spots, some of the probes were duplicated to fill in the whole array. The final array design consisted of 15,208 probes representing 3126 gene clusters of A. actinomycetemcomitans and 536 control probes.

The performance of individual probes in hybridization was not expected to be uniform across all strains because of sequence variation in orthologous genes across the strains. Therefore, in silico analyses were performed to filter out the probes that spanned highly variable regions. The filtering process started by comparing each probe sequence with all orthologous gene sequences that were found in each gene cluster using the NCBI BLAST package. A probe was considered acceptable if it matched all members of the target homologous gene cluster with at least 80% sequence identity over the probe length (60 bp) and did not match genes belonging to a different gene cluster. In addition, homologous gene clusters that were not represented by at least three probes were excluded from subsequent analysis. This filtering process resulted in a set of 10,934 probes representing 2676 gene clusters, including 1762 core genes (i.e. shared by all strains) and 914 accessory genes (i.e. present in one or more but not all strains).

## Comparative genomic hybridization by microarray

The genomic DNA was labeled and hybridized to the pan-genome microarray of A. actinomycetemcomitans according to the protocol recommended by Agilent Oligonucleotide Array-Based CGH for Genomic DNA (Agilent Technologies, Palo Alto, CA). Briefly, 0.5 µg genomic DNA was labeled with cyanine 3. DNA denaturation and fragmentation were performed using a thermocycler by mixing 2.5 µl Random Primers (Agilent) and 13 µl genomic DNA at 94°C for 10 min. The mixture was cooled to 4°C and 9.5 µl labeling master mix (Agilent) was added to make a total volume of 25 µl. DNA labeling was carried out using a thermocycler following the program: 37°C for 2 h, 65° C for 10 min, and holding at 4°C. Labeled genomic DNA was purified using individual Amicon 30-kDA filters (Millipore, Billerica, MA) following the manufacturer's protocol. The yield of labeled DNA was determined by measuring the absorbance at 260 nm (A<sub>260</sub>; DNA) and at 550 nm (A<sub>550</sub>; cyanine 3) using a Nano-Drop (ND-1000). Then, 40 µl hybridization sample mixture was loaded onto the glass slide containing the array and incubated at 65°C for 24 h with rotation. The arrays were then washed with Agilent CGH wash buffer and scanned using an Agilent scanner 6000C with the manufacturer's recommended settings for CGH arrays (scan area  $61 \times 21.6$  mm, 5µm resolution, 100% photo-multiplier tube (PMT) Red and Green channel (R&G)). Data were extracted from the scanner using Agilent FEATURE EXTRACTOR v10.5 software using protocol CGH\_105\_Dec08.

The signal data obtained from the Agilent FEATURE EXTRACTOR software were processed to extract the gProcessedSignal value, which is a background subtracted signal, for each probe. Since the labeling process involved incorporating cyanine 3-dUTP into the genomic sample, it is possible that probes that are rich in nucleotide A have a higher signal relative to probes with lower A content and may not accurately reflect the true abundance of the target sequence. For this reason, the signal values were normalized by dividing them by the total number of A nucleotides in the probe sequence. Finally, this value was transformed using log<sub>2</sub>. Processed signal values from probes that are targeting the same gene cluster were consolidated into a single value by averaging the signals. A specific cut-off point was then selected for declaring gene absence or presence. The comparative genomic hybridization data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (Edgar et al., 2002) and are accessible through GEO Series accession number GSE39143 (http://www.ncbi.nlm.nih.gov/geo/query/ acc.cgi?acc=GSE39143).

The performance of gene detection by CGH was first evaluated using four completed genome sequences and seven draft genomes. The sensitivity for gene detection by CGH was calculated as (number of genes in the genome detected by CGH)/[number of genes identified in the genome by whole genome sequence (WGS)]. The specificity for gene detection was calculated as (number of genes in the genome not detected by CGH)/(number of genes not found in the genome by WGS).

# PCR analysis of gene absence/presence

Six genes found to be distinct between JP2 and non-JP2 genotypes by CGH were confirmed by PCR analysis. A gene present in all strains was used as a positive control. The primers were designed based on published genomic data (Kittichotirat *et al.*, 2011) (see Supplementary material Table S2 for primers sequences). The 25- $\mu$ l PCR mixture included 50– 100 ng genomic DNA, 0.3  $\mu$ M concentration of each

primer, 2.5  $\mu$ l of 2 mM dNTPs, and 1 unit *Taq* DNA polymerase in 1 × *Taq* DNA polymerase buffer. The PCR amplification was performed with the following thermocycling profile: 2 min at 94°C for denaturation followed by 30 cycles of denaturation, annealing and extension at 94°C for 30 s, 64°C to 68°C for 1 min, and 72°C for 1 min, respectively; and then a final extension of 8 min at 72°C. The resultant amplicons were analysed in 1% agarose gels.

# Hierarchical clustering analysis of *A. actinomycetemcomitans* strains by genomic content

The software MEV (Saeed *et al.*, 2006) was used to perform a hierarchical clustering of binary data representing the present/absent profiles of all genes (as determined by CGH) using the Euclidean distance metric. The JSTREE tool (http://lh3lh3.users.source-forge.net/jstree.shtml) was then used to generate the cladogram.

# **Transcriptome analysis**

Three strains of JP2 genotype (HK1651, D28S-1 and D41S-1) and two strains of non-JP2 genotype (SCC1398 and ANH9381) were examined (Table 1). The transcriptome analysis was performed for each strain in biological duplicates. A starter culture was prepared by inoculating the bacteria as a single cell suspension (Karched et al., 2007) in tryptic soy broth with 0.6% yeast extract and incubated overnight at 37°C in an atmosphere supplemented with 5% CO<sub>2</sub>. The starter culture [optical density at 650 nm (OD<sub>650</sub>) of 0.11-0.59) was then diluted with fresh broth to  $\sim OD_{650} = 0.1$  and incubated for a further 4 h. The OD<sub>650</sub> was determined again to assure the continued growth of the bacteria (OD<sub>650</sub> in the range of 0.12-0.35). Aliquots of the bacterial cultures were also used to determine colony-forming units ml<sup>-1</sup> and checked for contamination.

The bacterial RNA was isolated using RiboPure<sup>TM</sup>-Bacteria Kit (Life Technology, Carlsbad, CA). The resultant RNAs were checked for DNA contamination by PCR using 16S rRNA primers. The RNAs were then stored at  $-70^{\circ}$ C until use. MessageAmp<sup>TM</sup> II-Bacteria kit (Life Technology) was used to label the RNA (500 ng for each sample) following the protocol recommended by the manufacturer, with a slight mod-

ification in the IVT reaction in which a combination of 2.6 µl T7 CTP and 1.4 µl of cyanine-3-labeled CTP were used for the master mix. The labeled samples were hybridized to the custom Agilent slides following the recommended protocol. A change to the protocol was made to use a nitrogen gun for drying the arrays after washing (instead of centrifugation or drying solution). RNA Yield and Quality was assessed by Nanodrop-1000 spectrometer and Agilent bioanalyser. When dry, the slides were immediately scanned using the Agilent Scanner 6000C according to the manufacturer's recommended settings for one-color gene expression. For the  $8 \times 15K$  custom arrays, we scanned at 5-µm resolution, dual pass at 100 and 10% PMT in one channel only (green, or cy-3). The data was then Feature-extracted using Agilent's FEATURE EXTRACTOR software v10.5 running the extraction protocol GE1\_105\_Dec08.

Expression data obtained from the Agilent FEATURE EXTRACTOR software were processed using the following steps. First, the gProcessedSignal value for all arrays, a background subtracted signal, was extracted for each probe. Quantile normalization was then applied across all the array data being compared to adjust for differences in the probe intensity distribution using the LIMMA package (Smyth & Speed, 2003). As we were interested in comparison of expression levels of the same genes among different A. actinomycetemcomitans strains, no adjustment was made for the signals based on the numbers of G nucleotides in the probe sequences. Finally, the normalized signal values from probes targeting the same gene cluster were consolidated into a single median value. A total of 1952 genes shared between JP2 and non-JP2 genotypes were analysed for their expression. The data for each replicate were treated individually in statistical analysis. Differentially expressed genes between genotypes were identified by *t*-test in MEV at P < 0.05. The expression data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (Edgar et al., 2002) and are accessible through GEO Series accession number GSE38943 (http://www.ncbi.nlm.nih.gov/geo/guery/acc.cgi?acc=G SE38943).

# Identification of gene operons

Operons in the genome of *A. actinomycetemcomitans* strain HK1651 were identified via the Database of

prOkaryotic OpeRons (DOOR) tool, which predicts bacterial gene operons using a classifier algorithm based on features such as intergenic distance, neighborhood conservation, phylogenetic distance, information from short DNA motifs, similarity score between GO terms of gene pairs, and length ratio between a pair of genes (Mao *et al.*, 2009).

## Quantitative real-time (qRT) PCR

Five of the same RNA samples used in transcriptome analysis (one sample per strain) were also used to assess the transcript levels of selected genes (p-cluster 00978 toIR, p-cluster 00011 lktA, p-cluster 01456 napD, p-cluster 00459 metF, p-cluster 00078 nrfE) by qRT-PCR using BioRad iCycler iQ<sup>®</sup> Real-Time PCR Detection System. A constitutively expressed housekeeping gene clpX (p-cluster 00289) was used as a reference to compare the expression levels (Ramsey & Whiteley, 2009). For each sample, 1 µg RNA in a 20-µl reaction mixture was reverse transcribed into first-strand cDNA using a SuperScript VILO kit (Life Technology). Reactions without reverse transcriptase or RNA template were included as controls. The firststrand cDNA synthesis was performed at 25°C for 10 min, 42°C for 60 min, 85°C for 5 min. The 20- $\mu l$ volume containing the cDNA was then diluted to 200 µl using sterile water. For qRT-PCR, a volume of  $2 \mu$ l of the diluted cDNA from each sample was used following the protocol described by the manufacturer. Briefly, the reaction mixture included 2.5 µl of each primer (3 µм), 12.5 µl of 2× Supermix, 2 µl cDNA, and water to 25 µl. The thermocycling profile consisted of four cycles as follows: Cycle 1:  $(1 \times)$  Step 1: 95.0°C for 3 min; Cycle 2: (40×) Step 1: 95.0°C for 10 s, Step 2: 55.0°C for 30 s; Cycle 3: (1×) Step 1: 95.0°C for 1 min; Cycle 4: (1×) Step 1: 55.0°C for 1 min. For the melting curves, the final DNA products were denatured at 95°C for 1 min and then incubated at 5°C below the annealing temperature for 1 min before the temperature was increased to 95°C at a ramp rate of 0.5°C per 10 s. For each sample, both target gene and reference gene were performed in triplicate. Additional controls include samples without cDNA for each target gene. Data analysis was performed based on the protocol provided by BioRad. The transcript levels determined by gRT-PCR were expressed as ratios to the sample with the lowest transcript level for each assessed gene.

# RESULTS

# Criteria for calling gene presence or absence by CGH

A total of 38 hybridizations for 19 strains (two replicates for each strain) were obtained. The distribution of the processed signals for each gene cluster for each *A. actinomycetemcomitans* strain showed a bimodal pattern, suggesting that the two peaks represented signals for the absent or the present genes (Fig. 1). The signal cut-off for determining gene presence or absence in CGH was taken to be the minimum count found between the two peaks. The cut-off point was determined individually for each CGH analysis (see Supplementary material Table S3 for complete dataset of the distribution of hybridization signals and cut-off points).

# Assessing the performance of gene detection by CGH

The results from two hybridizations failed to form bimodal distribution patterns (for strains D11S-1 and I63B) and were not analysed. The WGS information



**Figure 1** Distribution of processed hybridization signals for gene clusters in *Aggregatibacter actinomycetemcomitans* D7S-1. Processed hybridization signals from an array hybridized with D7S-1 genomic DNA were plotted where the *x*-axis represents the processed signal bins (e.g. 1 represents processed signals ranging from >0 to 1) and the *y*-axis represents the count of data points in each bin. The distribution shows a bimodal shape where the right and left peaks represent processed signal values of genes that are present and absent, respectively. A minimum count found between the two peaks (denoted as an asterisk in the figure) was chosen as a cut-off point for gene present/absent status (e.g. a processed signal above this value is considered present and a signal below this value is considered absent.

was used as the standard to test the performance of the pan-genome microarray in gene detection. The sensitivity for gene detection of the four complete genomes (D7S-1, HK1651, ANH9381 and D11S-1) was in the range of 0.948-0.986, whereas the specificity was between 0.867 and 0.976 (Table 2). The relatively low specificity (0.867) for one of the duplicates of D7S-1 was the result of a few false-positive results with hybridization signals just above the break point for gene detection. For the seven draft genomes, the sensitivity was 0.974-0.989 whereas the specificity was 0.900-0.986. The specificity for replicates for strain I23C was relatively low (0.900 and 0.902), which may be attributed to genes not detected by WGS but nevertheless detected by CGH. This interpretation is supported by the relatively poor WGS quality for I23C, with the lowest coverage ( $16 \times$ ), the highest number of contigs (400) and the highest num-

**Table 2** Sensitivity and specificity for gene detection by microarrayamong11 sequencedAggregatibacteractinomycetemcomitansstrains

Sensitivity	Specificity
0.986	0.976
0.986	0.867
0.984	0.986
0.985	0.986
0.948	0.978
0.986	0.981
0.979	0.967
0.989	0.980
0.983	0.977
0.986	0.977
0.981	0.971
0.988	0.977
0.984	0.979
0.985	0.981
0.983	0.900
0.983	0.902
0.986	0.977
0.984	0.977
0.977	0.912
0.974	0.986
	Sensitivity 0.986 0.986 0.984 0.985 0.948 0.985 0.948 0.986 0.979 0.989 0.983 0.986 0.981 0.984 0.985 0.984 0.985 0.983 0.983 0.983 0.983 0.983 0.986 0.984 0.977 0.974

Sensitivity = (number of genes in the genome detected by CGH)/ (number of genes identified in the genome by WGS); Specificity = (number of genes in the genome not detected by CGH)/(number of genes not found in the genome by WGS) where CGH is comparative genomic hybridization and WGS is whole genome sequence. ber of Q39 bases (33,125) in comparison with the other strains listed in Table 2 (Chen *et al.*, 2009, 2010a; Kittichotirat *et al.*, 2011).

Duplicates of the CGH results for nine of the sequenced strains and eight clinical isolates were further examined to evaluate the consistency in gene detection between duplicates. As shown in Table 3, the percentage of shared genes detected in the replicates ranged from 95.95-100%. In some instances the basis for the differences between duplicates could be identified. For strain D7S-1, the discrepancy between replicates could be attributed to the relatively low specificity (0.867) in one of the replicates (Table 2), whereas for strain ANH9381 the differences in detection between replicates were probably the result of the relatively low sensitivity (0.948) in one of the replicates (Table 2). The information was used to help in selecting one of the duplicates from each strain for further analysis below.

# Clustering analysis of genome content of *A. actinomycetemcomitans* serotype b and c strains

Based on the CGH results, a hierarchical clustering analysis of the genomic content (based on the presence and absence of genes) was performed for strains of serotypes a–f (Fig. 2). Serotypes a, d, e and f formed a major branch separate from serotypes b and c. The six serotype b JP2 genotype strains formed a unique group separated from six other serotype b non-JP2 genotype strains and three serotype c strains.

# Genome content variation between JP2 and non-JP2 genotypes

Among the 2044 gene clusters found in any of the six strains of JP2 genotype, 1968 gene clusters (96%) were shared by all strains. In comparison, a total of 2201 gene clusters were present among the six strains of non-JP2 genotype with 1986 gene clusters (90%) shared by all six non-JP2 strains.

Nine genes were found exclusively in JP2 genotype strains whereas 12 genes were found only in non-JP2 genotype strains (Table 4). The genotype-specificity of five gene clusters (p-cluster03717, p-cluster03907, and p-cluster03273 of JP2 strains, and p-cluster01731 p-cluster02176 of non-JP2 strains) were selected and

Comparative genomics of JP2 and non-JP2

Strain	No. of genes in replicate #1	No. of genes in replicate #2	No. of genes found only in replicate #1	No. of genes found only in replicate #2	No. of genes detected in both replicates (% of genes)
HK1651	2003	2005	0	2	2003 (99.90)
D7S-1	2113	2173	4	64	2109 (96.88)
SCC1398	1991	1992	0	1	1991 (99.95)
SCC393	2164	2171	2	9	2162 (99.49)
D17P-2	2101	2105	0	4	2101 (99.81)
D18P-1	2133	2145	4	16	2129 (99.07)
ANH9381	1957	2033	2	78	1955 (96.07)
SCC2302	1987	1977	0	10	1977 (99.50)
123C	2055	2055	0	0	2055 (100)
A26	2036	2037	0	1	2036 (99.95)
ATCC29524	2035	2032	4	1	2031 (99.75)
194	1992	2037	0	45	1992 (97.79)
G104-2	2062	2059	3	0	2059 (99.85)
S067	1993	2003	5	15	1988 (99.00)
G111-1	2011	2014	0	3	2011 (99.85)
G121-2	2040	2038	2	0	2038 (99.90)
D41S-1	1989	2073	0	84	1989 (95.95)

Table 3 Gene detection between the replicates for each of the 17 Aggregatibacter actinomycetemcomitans strains

The % of genes detected was calculated by (no. of detected genes in both replicates)/(total unique genes identified by both replicates).

confirmed by PCR analysis in the 12 serotype b strains (data not shown). For the purpose of clarity, these 21 JP2-specific and non-JP2-specific genes are further described below in the context of the genomes of the fully sequenced HK1651 and ANH9381 strains.

# Insertion of an island 'iAAI-1' in the genome of HK1651

JP2-specific genes, which are identified by p-cluster03231, p-cluster03273 and p-cluster03132 (see Table 4), were found to be contiguous and located on a 23,394-bp genomic island (here renamed iAAI-1; HK1651 nucleotide coordinates 6573-29966) comprising 20 genes reported previously (Chen et al., 2005; Kittichotirat et al., 2011). The iAAI-1 island was flanked by a truncated malate dehydrogenase and a truncated flavoprotein homolog, followed by an intact cytochrome c protein homolog (Chen et al., 2005). Strain ANH9381 did not harbor iAAI-1 and instead contained a full-length malate dehydrogenase gene and flavoprotein gene in the same locus (Table 4; non-JP2-specific p-cluster1210). The same genetic arrangement (i.e. having an intact malate dehydrogenase and flavoprotein) was also found in the genomes of serotype b non-JP2 genotype strains SCC1398 and I23C (data not shown). Although homologs of the genes on iAAI-1 were detected by CGH in all of the five clinical JP2 strains, the presence of iAAI-1 (as a contiguous island) in these strains was not verified.

# Insertion of an island 'iANH-1' in the genome of ANH9381

The genes p-cluster02176, p-cluster01731, p-cluster01585 and p-cluster09851 were detected in the non-JP2 genotype but not in the JP2 genotype by CGH (Table 4). These four genes were found to be contiguous and located on a genomic island (designated iANH-1) in ANH9381 (Fig. 3). A homolog of the iANH-1 island was also found in the same locus in strain SCC1398 (annotated as genes SCC1398\_0850 to SCC1398\_0858) (Kittichotirat *et al.*, 2011).

# Different islands identified in the same locus in the genomes of HK1651 and ANH9381

JP2-specific p-cluster03717 and p-cluster03907 were located on a 7244-bp genomic island designated as iAAI-3 (HK1651 nucleotide coordinates 1281101– 1288344) flanked by *guaA* and a gene encoding hypothetical protein (AA01895) (Fig. 4). A different genomic island of 9392 bp was identified (here



**Figure 2** Hierarchical clustering analysis of *Aggregatibacter actinomycetemcomitans* strains by genomic content. The MEV software was used to perform the hierarchical clustering of binary data representing the present/absent profiles of genes using Euclidean distance metric. The JSTREE tool was then used to create the cladogram. The serotype and genotype information is provided in parenthesis. Asterisks indicate clinical strains without genome sequence information. The results showed that strains of serotypes a, d, e and f formed a major branch. The six serotype b JP2 genotype strains formed a unique cluster but the six serotype b non-JP2 genotype strains were mixed together with three serotype c strains.

designated as iANH-2; strain ANH9389 nucleotide coordinates 1408870–1418261) in the same locus in ANH9381 (Fig. 3) (Chen *et al.*, 2012). While homologous genes of iANH-2 were identified in I23C, they were located in two contigs that may or may not be a homolog of iANH-2. A distinct 5682-bp genomic island comprising 13 genes that demonstrated partial homology to both iAAI-3 and iANH-2 was found in the same locus in strain I23C. This island was also identified in the previous study (I23C contig00245, nucleotide coordinates 2082–7763) (Kittichotirat *et al.*, 2011).

For the remaining JP2-specific and non-JP2-specific genes, none was found to be on genomic islands. In one instance, the insertion of a p-cluster04245 within *radC* (DNA repair protein) divided the gene into two apparently non-functional pseudogenes in HK1651, whereas ANH9381 (as well as SCC1398 and I23C) has an intact *radC*.

# Transcriptome profiles of log phase *A. actinomycetemcomitans*

The results for two biological duplicates per strain for *A. actinomycetemcomitans* JP2 genotype strains HK1651, D28S-1 and D41S-1 and for non-JP2 genotype strains ANH9381 and SCC1398 are presented in the Supplementary material (Table S4). The *R* values for the linear correlation between duplicates were 0.64 for HK1651, 0.95 for D41S-1, 0.87 for D28S-1, 0.97 for SCC1398 and 0.72 for ANH9381.

One sample from each of the five strains was used to verify expression of each of five selected genes. The results were compared with the same samples used for gene expression analysis with the microarray. The correlations between the results by qRT-PCR and hybridization with the microarray are shown in Fig. 5. Three of the five genes demonstrated excellent correlations between results obtained by Comparative genomics of JP2 and non-JP2

p-Cluster	Gene ID (nucleotide coordinates)	Length (bp)	Function	Island
Genes speci	ic to JP2 genotype			
03231	AA00018 (10690–11817)	1128	Hypothetical protein	iAAI-1
03273	AA00019 (11839–12882)	1044	Phage protein	iAAI-1
03132	AA00020 (12939–14546)	1608	Phage protein	iAAI-1
03717	AA01904 (1284588–1285109)	522	Hypothetical protein	iAAI-3
03907	AA01911 (1287931–1288344)	414	Hypothetical protein	iAAI-3
03979	AA00785 (539732–539361)	372	Phage integrase	Non-Island
05260	AA00666 (457610–457714)	105	Hypothetical protein	Non-Island
04245	AA01318 (892473–892222)	252	Hypothetical protein	Non-Island
03340	AA01449 (971629–970709)	921	Prophage integrase	Non-Island
Genes specit	ic non-JP2 genotype			
01473	ANH-1698 (1601972–1602181)	210	TOBE domain protein	Non-Island
05754	ANH_1697 (1600856–1601890)	1035	Iron chelatin ABC transporter	Non-Island
08744	N/A (816767–816909) <sup>1</sup>	141	Hypothetical protein	Non-Island
01210	ANH9381_0165 (137315-136116)	1200	Flavoproteins	Non-Island
N/A	N/A (1378798–1378947) <sup>1</sup>	150	Hypothetical protein	iANH-1
01585	ANH9381_1475 (1378521-1378814)	294	CRISPR-associated protein Cas2	iANH-1
01731	ANH9381_1474 (1377573-1378517)	945	CRISPR-associated protein Cas1	iANH-1
02176	ANH9381_1473 (1375030-1377450)	2421	ATP-dependent OLD family endonuclease	iANH-1
02878	N/A (1140063–1139875) <sup>1</sup>	189	Antirestriction protein	Non-Island
04532	ANH9381_1521 (1414996-1414814)	183	YcfA family protein	iANH-2
03415	ANH9381_1526 (1418261-1417428)	834	Prophage integrase	iANH-2
04299	ANH9381_2215 (2116174-2115722)	453	Hypothetical protein	Non-Island

Table 4 JP2-specific and Non-JP2-specific genes

<sup>1</sup>N/A, genes were not detected by the annotation programs, but were identified by BLAST search against the genome.



**Figure 3** Genetic maps of the genomic island iANH-1 in ANH9381 and the corresponding region in strain HK1651. All open reading frames are noted and indicated by blue (forward) or green (reverse) pentagons. The gene encoding CRISP-associated Cas3 is truncated at the 3' end by more than 50% of its length in HK1651. In addition, the *glnE* and the genes for Cas3 were transcribed in opposite directions between these strains, suggesting a genomic reversion in conjunction with the insertion/deletion of the island iANH-1. HP: hypothetical protein.



Figure 4 Genetic map of the genomic island iAAI-3 in strain HK1651 and genomic island iANH-2 in strain ANH9381. Red arrows delineate the boundary of the islands. Open reading frames are noted and indicated by blue (forward) or green (reverse) pentagons. HP: hypothetical protein. The islands are flanked by a hypothetical protein and guaA (GMP synthase).

qRT-PCR and microarray analysis (R = 0.89-0.93). For the other two genes, the overall trend in the transcript levels among samples was consistent between the two quantification methods.

The 150 genes with significantly different transcript levels between JP2 and non-JP2 genotypes (P < 0.05 by t-test) and their expression ratios are shown in the Supplementary material (Table S5). We further used the prOkaryotic OpeRons (DOOR) tool to predict the operons of A. actinomycetemcomitans HK1651 (see Supplementary material, Table S6), and identified operons with at least two genes with a twofold difference or greater in expression levels between JP2 and non-JP2 genotypes. Five operons were found to meet the criteria and are listed in Table 5. As expected, the expression levels of the leukotoxin operon were higher in JP2 genotype than in non-JP2 genotype. For the remaining four operons, two were more highly expressed in the JP2 genotype, and the others were more highly expressed in the non-JP2 genotype.

# Analysis of promoters of four differentially expressed operons

Differential gene expression could possibly be attributed to different promoter structures in the two geno-

Gen. DISCUSSION Aggregatibacter

types. Therefore, the upstream regions of the four operons differentially expressed between HK1651 and ANH9381 were analysed for sequence variations. The sgaB/tkt operon and its upstream region in HK1651 and ANH9381 are depicted in Fig. 6. A 223bp insert (including the 93-bp AA02569 sequence encoding a hypothetical protein annotated by Oral-Gen) flanked by 48-bp and 49-bp inverted repeated elements was found in HK1651, whereas only the 49-bp element was found in the same locus in ANH9381. However, the same 223-bp insert was found in the non-JP2 strain SCC1398. Therefore, the 223-bp insertion is not JP2-specific and cannot be the cause of differential expression of the operon. For the other three operons, no differences were found in the promoters between JP2 and non-JP2 genotypes. It was noted that the napG gene of HK1651 is 816 bp in length as found in all other A. actinomycetemcomitans strains, not 678-bp as reported by Oral-

actinomycetemcomitans demonstrates remarkable strain-to-strain variation in genome content. Although the pan-genome of A. actinomycetemcomitans is open-ended (Kittichotirat et al., 2011),



**Figure 5** Correlation of the gene expression levels determined by quantitative real-time polymerase chain reaction (qRT-PCR) and by hybridization with microarray. Identical samples (three strains of JP2 genotype and two strains of non-JP2 genotype) were used in both quantification methods to determine the expression levels of the five selected genes: *lktA*, *tolR*, *nrfE*, *napD* and *metF*. The expression level of the housekeeping gene *clpX* was used to normalize the levels of the target genes in each sample. For qRT-PCR, the sample with the lowest transcript level was arbitrarily set as 1. Linear trend and  $R^2$  value are provided for each graph. Excellent correlations were found for *tolR*, *nrfE* and *napD*. For *lktA* and *metF*, the trend in the expression levels was consistent between the two quantification methods.

16 or fewer new genes are expected with every additional strain included for genome sequencing in the future. The pan-genome microarray described in this study was designed based on the genomes of 18 diverse *A. actinomycetemcomitans* strains. It will provide a good degree of gene coverage and can serve as a cost-effective tool for interrogating the gene contents of *A. actinomycetemcomitans* strains. Also, this microarray is applicable for transcriptome analysis of diverse *A. actinomycetemcomitans* strains. Although alternatives for genomic comparison or transcriptome analysis are available (such as high throughput sequencing), the costs may be higher than using the microarray.

Gene detection with the pan-genome microarray was evaluated with both sequenced strains and clinical isolates of *A. actinomycetemcomitans*. Overall, the performance for gene detection was excellent

and demonstrated high degrees of sensitivity, specificity and reproducibility between biological duplicates. Some of the probes on the pan-genome microarray were excluded in our analysis because they may provide inconsistent hybridization results across the strains as a result of sequence variations. Nearly all the excluded probes were designed for genes that encode small hypothetical proteins and may not represent critical genetic elements of the strains. Nevertheless, we are currently redesigning the pan-genome microarray to further improve its utility.

The results of cluster analysis in this study confirmed our previous finding that *A. actinomycetemcomitans* strains were divided into two major branches. One comprised strains of serotypes a, d, e and f, and the other comprised strains of serotypes b and c. This study further focused on assessing the

Cluster ID	HK1651 Gene ID	Length (bp)	Product description	JP2/nonJP2 ratio
p-cluster00813	AA00401	594	slyD: FKBP-type peptidyl-prolyl cis-trans isomerase	1.721
p-cluster00710	AA00402	717	Periplasmic/secreted protein	2.634 <sup>1</sup>
p-cluster00965	AA00404	432	COG3076: Uncharacterized protein conserved in bacteria	3.276 <sup>1</sup>
p-cluster00993	AA00406	393	mscL: large-conductance mechanosensitive channel	2.416 <sup>1</sup>
p-cluster00218	AA00407	1374	trkA: trk system potassium uptake protein	3.008
p-cluster00233	AA00408	1353	sun, fmu, fmv, rsmB: SUN protein (FMU protein)	3.502 <sup>1</sup>
p-cluster01270	AA00409	954	fmt: methionyl-tRNA formyltransferase	2.662
p-cluster01400	AA00410	510	def: peptide deformylase	2.656 <sup>1</sup>
p-cluster00778	AA00546	633	napC: denitrification system component	0.091 <sup>1</sup>
p-cluster00946	AA00547	447	napB: periplasmic nitrate reductase	0.089 <sup>1</sup>
p-cluster00545	AA00549	879	napH: ferredoxin-type protein	0.06 <sup>1</sup>
p-cluster06328	AA00550	678	napG: ferredoxin-type protein	0.054 <sup>1</sup>
p-cluster01495	AA00552	2484	napA: periplasmic nitrate reductase precursor	0.046
p-cluster01456	AA00553	282	napD: NapD component of periplasmic nitrate reductase	0.034
p-cluster01596	AA00555	162	Hypothetical protein D11S_0211	1.157
p-cluster00484	AA02563	942	tkt: transketolase C-terminal section	0.593
p-cluster01310	AA02564	531	tkt: transketolase N-terminal section	0.348 <sup>1</sup>
p-cluster06816	AA02566	1092	PTS system, IIC component	0.401 <sup>1</sup>
p-cluster09728	AA02568	267	sgaB: sugar phosphotransferase component II B	0.606
n-cluster00192	AA02803	1431	lktD	1 656
p-cluster00056	AA02805	2121	lktB	3.996 <sup>1</sup>
p-cluster00011	AA02806	3165	lktA	2 925 <sup>1</sup>
p-cluster13459	AA02807	504	lktC	6.1 <sup>1</sup>
p-cluster05272	AA02828	105	Hypothetical protein	1.449
p-cluster00938	AA02829	465	pal: peptidoglycan-associated outer membrane	1.183
p-cluster00276	AA02830	1278	tolB: colicin tolerance protein	2.768 <sup>1</sup>
p-cluster01209	AA02832	1206	tolA: outer membrane integrity protein	N/A <sup>2</sup>
p-cluster00978	AA02834	420	toIR: colicin transport protein	2.862 <sup>1</sup>
p-cluster05329	AA02835	96	Hypothetical protein	2.11 <sup>1</sup>
p-cluster00717	AA02836	687	tolQ: colicin transport protein	2.008 <sup>1</sup>
p-cluster00991	AA02837	402	Tol-pal system-associated acyl-CoA	1.731 <sup>1</sup>
	/ / / / / / / / / / / / / / / / / / / /	102	thioesterase	
p-cluster01904	AA02838	276	Hypothetical protein APJL_0311	1.891'
p-cluster05330	AA02839	96	Hypothetical protein	1.028
p-cluster00338	AA02840	1134	cydB: cytochrome D ubiquinol oxidase, subunit II	0.762
p-cluster01161	AA02841	1563	cydA: cytochrome D ubiquinol oxidase, subunit I	0.838

Table 5 Differentially expressed operons between Aggregatibacter actinomycetemcomitans JP2 and non-JP2 genotypes

<sup>1</sup>Significantly different between JP2 and non-JP2 genotypes by *t*-test at P < 0.05.

<sup>2</sup>No analysis was performed because of problems in probe design.



**Figure 6** Genetic map of *sagB/tkt* operon in HK1651 and ANH9381. The dashed lines indicate the homologous regions. Open reading frames are noted and indicated by blue (forward) or green (reverse) pentagons. An insertion of a 223-base-pair (bp) fragment flanked by 48-bp/49-bp repeats (indicated by light blue boxes) is found in HK1651, while only the 49-bp element was found in the comparable region in ANH9381. It was noted that the tkt (transketolase N-terminal section) was truncated in HK1651 (annotated at OralGen as two overlapping genes of AA02564 and AA02565) but intact in ANH8381. Here is the location of each repeat in HK1651 and ANH9381 genomes: >hk1651-repeat-1 (1788949-1788997); TCCACGCTTGGACCGACACAAGCAAAAGCGCGGATGCTTGCGCTATCAT; >hk1651-repeat-2 (1789221-1789268); TCCACGCTTGGACCGGACGCTTGCGCTATCAT; >anh9381-repeat-1 (1922975-1923023); TCCACGCTTGGACCG ACGCTATCAT.

variation in genomes and transcriptomes of JP2 and non-JP2 genotypes. Haubek et al. (2007) suggested that strains of JP2 genotype arose relatively recently as a distinct clone derived from an ancestral strain of serotype b non-JP2. Moreover, serotype b JP2 strains could be distinguished into two groups based on a mutation in the hbpA pseudogene. One group was found in individuals from North Africa of distinct ethnic group (Arabs) and the other was associated with individuals from West Africa (Africans) (Haubek et al., 2007). The results obtained in this study are consistent with this interpretation because it also shows that JP2 and non-JP2 strains constitute separate clusters in the dendrogram based on the analysis of genome content of the strains (i.e. presence and absence of 2676 genes). As expected, the genomic diversity of non-JP2 strains was greater than JP2 strains. To the best of our knowledge, this is the first study to demonstrate the differences between JP2 and non-JP2 genotypes by full genome content analysis. Other than strain HK1651, no sequence information of hbpA pseudogene was available for the other four JP2 strains. However, the four JP2 strains in this study likely belong to the West Africa subgroup of JP2 genotype because they were identified from ethnic groups of African and Hispanic origin in the USA and Brazil, as suggested by Haubek et al. (2007).

One unique aspect of the JP2 genotype is its apparent high virulence (DiRienzo et al., 1994b; Haubek et al., 2008). A premise of this study is that other virulence factors, e.g. genomic islands, may be the basis for the high virulence of the JP2 genotype. The acquisition of genomic islands is thought to play a crucial role in bacterial evolution (Ochman et al., 2000). This study identified several genomic islands specific to JP2 and non-JP2 genotypes. It was also noted that the insertion of genomic islands may interrupt the function of housekeeping genes. Moreover, different genomic islands may be inserted into the same locus in different genomes, suggesting an insertion hot-spot for genomic islands in A. actinomycetemcomitans. Our laboratory is currently assessing the functions of these genomic islands and the effects of their genomic insertion in A. actinomycetemcomitans.

Variable virulence may also arise through differential regulation of virulence determinants. Therefore, transcriptome analysis was performed for *A. actinomycetemcomitans* strains grown to log phase in an enriched liquid medium. The expression levels of five genes determined by microarray hybridization and by qRT-PCR were also compared. The gene expression levels determined by these two methods were relatively consistent except for *lktA*. This could be a result of experimental variables that were difficult to

replicate between experiments. It was noted that the transcriptomes of strain HK1651 demonstrated the least correlation between the biological duplicates compared with other strains. The correlations between gene expression by microarray hybridization and qRT-PCR improved ( $R^2$  values of 0.5732, 0.9924, 0.851. 0.923 and 0.8168 for *lktA*, *tolR*, *nrfE*, *napD* and *metF*, respectively) if the data for strain HK1651 were removed from the analysis.

In our transcriptome analysis, we found that the expression levels of 150 genes (7.7%) were significantly different between JP2 and non-JP2 genotypes (*t*-test, P < 0.05) whereas 98 genes (1952  $\times$  0.05) were expected. We reasoned that the overall expression patterns of operons may be less likely to be subjected to experimental variations than those of individual genes. Therefore, we identified five operons each with two or more genes demonstrating two-fold or greater expression ratios between JP2 and non-JP2 genotypes. The leukotoxin operon was one of the five operons identified, suggesting the validity of our approach. The expression patterns of the other four operons were not known to be genotype-specific. The differences in the expression levels of the nap operon between JP2 and non-JP2 genotypes were particularly pronounced, reaching 10-fold or greater differences between genotypes.

The genotype-specific differential expression pattern of the leukotoxin operon correlated with the promoter structure of the operon as expected (Brogan *et al.*, 1994). A 223-bp insertion was found in the putative promoter the *tkt* operon in strain HK1651 but this insertion was not JP2-specific. No evidence of promoter structure variation was found for the remaining three operons. These results suggest that the observed differential gene expression may be a result of distinct gene regulation mechanisms in JP2 and non-JP2 genotypes.

Testing the role of these four operons in the pathogenesis of *A. actinomycetemcomitans* in periodontitis is beyond the scope of this study. Nevertheless, the results may be used to design experiments to test the influence of these operons on the virulence of *A. actinomycetemcomitans*. The differentially expressed *nap* operon is an interesting choice for further testing. The membrane-bound nitrate reductase has been shown to reduce nitrate (terminal electron acceptor) to nitrite for anaerobic growth in *Escherichia coli* (Potter *et al.*, 1999) and may provide a similar function for *A. actinomycetemcomitans.* Another operon of interest encodes genes of the Tol-Pal system, which is essential for maintaining the integrity of cell envelopes in many gram-negative bacterial species (Godlewska *et al.*, 2009). Pal protein is one of the pathogen-associated molecular patterns and modulates host immune response via Toll-like receptor 2 (Godlewska *et al.*, 2009). In *A. actinomycetemcomitans*, Pal is released as a soluble protein and may enhance the local and systemic immune responses in periodontitis (Paul-Satyaseela *et al.*, 2006; Karched *et al.*, 2008; Oscarsson *et al.*, 2008). Whether the JP2 genotype releases greater amounts of Pal protein than non-JP2 genotype remains to be investigated.

In conclusion, we have created a custom-designed pan-genomic microarray and validated it for applications in comparative genomic analysis and transcriptome analysis of A. actinomycetemcomitans. New genotype-specific genes or genomic islands of unknown functions were identified in strains of JP2 and non-JP2 genotypes. Some genomic islands appeared to be inserted into specific locations in the genomes in strains of JP2 and non-JP2 genotypes. Transcriptome analysis confirmed the upregulation of the leukotoxin operon in the JP2 genotype, and further identified four operons whose expression pat-The terns appeared to be genotype-specific. information can be used to formulate hypothesis-driven experiments to examine the molecular basis of the difference in virulence between JP2 and non-JP2 genotypes of A. actinomycetemcomitans.

# ACKNOWLEDGEMENTS

This study was supported by NIDCR grant R01 DE12212. We are grateful to Bruce Roe, Fares Najar, Sandy Clifton, Tom Ducey, Lisa Lewis and Dave Dyer for the use of unpublished nucleotide sequence data from the *A. actinomycetemcomitans* Genome Sequencing Project at The University of Oklahoma.

# REFERENCES

- Asikainen, S., Chen, C. and Slots, J. (1995) Actinobacillus actinomycetemcomitans genotypes in relation to serotypes and periodontal status. Oral Microbiol Immunol 10: 65–68.
- Bozdech, Z., Zhu, J., Joachimiak, M.P., Cohen, F.E., Pulliam, B. and DeRisi, J.L. (2003) Expression profiling of

#### Comparative genomics of JP2 and non-JP2

the schizont and trophozoite stages of *Plasmodium falciparum* with a long-oligonucleotide microarray. *Genome Biol* **4**: R9.

- Brogan, J.M., Lally, E.T., Poulsen, K., Kilian, M. and Demuth, D.R. (1994) Regulation of *Actinobacillus actinomycetemcomitans* leukotoxin expression: analysis of the promoter regions of leukotoxic and minimally leukotoxic strains. *Infect Immun* 62: 501–508.
- Chen, W., Wang, Y. and Chen, C. (2005) Identification of a genomic island of *Actinobacillus actinomycetemcomitans. J Periodontol* **76**: 2052–2060.
- Chen, C., Kittichotirat, W., Si, Y. and Bumgarner, R. (2009) Genome sequence of *Aggregatibacter actinomycetemcomitans* serotype c strain D11S-1. *J Bacteriol* **191**: 7378–7379.
- Chen, C., Kittichotirat, W., Chen, W., Downey, J.S., Si, Y. and Bumgarner, R. (2010a) Genome sequence of naturally competent *Aggregatibacter actinomycetemcomitans* serotype a strain D7S-1. *J Bacteriol* **192**: 2643–2644.
- Chen, C., Wang, T. and Chen, W. (2010b) Occurrence of *Aggregatibacter actinomycetemcomitans* serotypes in subgingival plaque from United States subjects. *Mol Oral Microbiol* **25**: 207–214.
- Chen, C., Kittichotirat, W., Chen, W., Downey, J.S. and Bumgarner, R. (2012) Genome sequence of a serotype b non-JP2 *Aggregatibacter actinomycetemcomitans* strain, ANH9381, from a periodontally healthy individual. *J Bacteriol* **194**: 1837.
- DiRienzo, J.M., Slots, J., Sixou, M., Sol, M.A., Harmon, R. and McKay, T.L. (1994a) Specific genetic variants of *Actinobacillus actinomycetemcomitans* correlate with disease and health in a regional population of families with localized juvenile periodontitis. *Infect Immun* **62**: 3058–3065.
- DiRienzo, J.M., Slots, J., Sixou, M., Sol, M.A., Harmon, R. and McKay, T.L. (1994b) Specific genetic variants of *Actinobacillus actinomycetemcomitans* correlate with disease and health in a regional population of families with localized juvenile periodontitis. *Infect Immun* **62**: 3058–3065.
- Edgar, R., Domrachev, M. and Lash, A.E. (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* **30**: 207–210.
- Godlewska, R., Wisniewska, K., Pietras, Z. and Jagusztyn-Krynicka, E.K. (2009) Peptidoglycan-associated lipoprotein (Pal) of Gram-negative bacteria: function, structure, role in pathogenesis and potential application in immunoprophylaxis. *FEMS Microbiol Lett* **298**: 1–11.

- Haubek, D., Dirienzo, J.M., Tinoco, E.M. *et al.* (1997)
   Racial tropism of a highly toxic clone of *Actinobacillus actinomycetemcomitans* associated with juvenile periodontitis. *J Clin Microbiol* **35**: 3037–3042.
- Haubek, D., Poulsen, K. and Kilian, M. (2007) Microevolution and patterns of dissemination of the JP2 clone of *Aggregatibacter (Actinobacillus) actinomycetemcomitans. Infect Immun* **75**: 3080–3088.
- Haubek, D., Ennibi, O.K., Poulsen, K., Vaeth, M., Poulsen, S. and Kilian, M. (2008) Risk of aggressive periodontitis in adolescent carriers of the JP2 clone of *Aggregatibacter (Actinobacillus) actinomycetemcomitans* in Morocco: a prospective longitudinal cohort study. *Lancet* **371**: 237 –242.
- Kaplan, J.B., Schreiner, H.C., Furgang, D. and Fine, D.H. (2002) Population structure and genetic diversity of *Actinobacillus actinomycetemcomitans* strains isolated from localized juvenile periodontitis patients. *J Clin Microbiol* **40**: 1181–1187.
- Karched, M., Paul-Satyaseela, M. and Asikainen, S. (2007) A simple viability-maintaining method produces homogenic cell suspensions of autoaggregating wildtype Actinobacillus actinomycetemcomitans. J Microbiol Methods 68: 46–51.
- Karched, M., Ihalin, R., Eneslatt, K. *et al.* (2008) Vesicleindependent extracellular release of a proinflammatory outer membrane lipoprotein in free-soluble form. *BMC Microbiol* **8**: 18.
- Kittichotirat, W., Bumgarner, R. and Chen, C. (2010) Markedly different genome arrangements between serotype a strains and serotypes b or c strains of *Aggregatibacter actinomycetemcomitans. BMC Genomics* **11**: 489.
- Kittichotirat, W., Bumgarner, R.E., Asikainen, S. and Chen, C. (2011) Identification of the pangenome and its components in 14 distinct *Aggregatibacter actinomycetemcomitans* strains by comparative genomic analysis. *PLoS ONE* **6**: e22420.
- Mao, F., Dam, P., Chou, J., Olman, V. and Xu, Y. (2009) DOOR: a database for prokaryotic operons. *Nucleic Acids Res* **37**: D459–D463.
- Ochman, H., Lawrence, J.G. and Groisman, E.A. (2000) Lateral gene transfer and the nature of bacterial innovation. *Nature* **405**: 299–304.
- Oscarsson, J., Karched, M., Thay, B., Chen, C. and Asikainen, S. (2008) Proinflammatory effect in whole blood by free soluble bacterial components released from planktonic and biofilm cells. *BMC Microbiol* **8**: 206.
- Paju, S., Carlson, P., Jousimies-Somer, H. and Asikainen,
   S. (2000) Heterogeneity of *Actinobacillus actinomyce*temcomitans strains in various human infections and

relationships between serotype, genotype, and antimicrobial susceptibility. *J Clin Microbiol* **38**: 79–84.

Paul-Satyaseela, M., Karched, M., Bian, Z. *et al.* (2006)
 Immunoproteomics of *Actinobacillus actinomycetem-comitans* outer-membrane proteins reveal a highly
 immunoreactive peptidoglycan-associated lipoprotein.
 *J Med Microbiol* 55: 931–942.

Potter, L.C., Millington, P., Griffiths, L., Thomas, G.H. and Cole, J.A. (1999) Competition between *Escherichia coli* strains expressing either a periplasmic or a membranebound nitrate reductase: does Nap confer a selective advantage during nitrate-limited growth? *Biochem J* 344: 77–84.

Ramsey, M.M. and Whiteley, M. (2009) Polymicrobial interactions stimulate resistance to host innate immunity through metabolite perception. *Proc Natl Acad Sci U S A* **106**: 1578–1583.

Rylev, M. and Kilian, M. (2008) Prevalence and distribution of principal periodontal pathogens worldwide. *J Clin Periodontol* **35**: 346–361.

Saeed, A.I., Bhagabati, N.K., Braisted, J.C. *et al.* (2006) TM4 microarray software suite. *Methods Enzymol* **411**: 134–193.

Slots, J. (1999) *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* in periodontal disease: introduction. *Periodontol 2000* **20**: 7–13. Smyth, G.K. and Speed, T. (2003) Normalization of cDNA microarray data. *Methods* **31**: 265–273.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

**Table S1.** De novo 454 sequencing of four Aggregatibacter actinomycetemcomitans strains.

 Table S2.
 Primers used for confirmation of genes

 specific to JP2 and non-JP2 genotypes.

 Table S3.
 Comparative genomic hybridization signals distribution and cut-off point for gene detection.

**Table S4.** Transcriptome of log-phase Aggregatibacter actinomycetemcomitans.

**Table S5.** Differentially expressed genes between *Aggregatibacter actinomycetemcomitans* of JP2 and non-JP2 genotypes.

**Table S6.** Predicted operons of Aggregatibacter ac-tinomycetemcomitans strain HK1651.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article. Copyright of Molecular Oral Microbiology is the property of Wiley-Blackwell and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.