

G Shen
U Hägg
MA Darendeliler

Skeletal effects of bite jumping therapy on the mandible – removable vs. fixed functional appliances

Authors' affiliations:

G. Shen, Associate Professor,
Discipline of Orthodontics,
Faculty of Dentistry,
The University of Sydney,
Sydney, Australia

U. Hägg, Professor and Chair,
Orthodontics, Faculty of Dentistry,
The University of Hong Kong,
Hong Kong SAR, China

M.A. Darendeliler, Professor and Chair,
Discipline of Orthodontics,
Faculty of Dentistry,
The University of Sydney,
Sydney, Australia

Correspondence to:

Gang Shen
Discipline of Orthodontics
Faculty of Dentistry
The University of Sydney
Level 2, 2 Chalmers Street
Surry Hill
NSW 2010
Australia
Tel.: +61 2 9351 8314
Fax: +61 29351 8336
E-mail: gshe6437@mail.usyd.edu.au

Dates:

Accepted 7 September 2004

To cite this article:

Orthod Craniofacial Res 8, 2005; 2–10
Shen G, Hägg U, Darendeliler MA:
Skeletal effects of bite jumping therapy on the
mandible – removable vs. fixed functional
appliances

Copyright © Blackwell Munksgaard 2005

Structured Abstract

Authors – Shen G, Hägg U, Darendeliler MA

Objective – Based on an extensive review of the literature, the aim of this study was to explore the mainstream consensus on the controversial topic of whether the bite jumping treatment could enhance mandibular growth.

Design – The data for removable and fixed functional appliances were respectively comprehended and analyzed with regard to their attributes in mandibular growth modification. Furthermore, numerous reported findings were assessed by relating them to some important factors influencing the effects of bite jumping, such as treatment timing, treatment duration and post-treatment follow-up, to allow for a more objective and accurate evaluation.

Results – The key differences between removable and fixed appliances are working hours (intermittent vs. continuous), length of treatment time (long vs. short), optimal treatment timing (before puberty growth vs. at or after puberty spurt), and mode of bite-jumping (considerable vertical opening vs. limited vertical opening). These different features lead to different treatment effects on mandibular and TMJ growth, such as the intensity of possibly increased growth (clinically less significant vs. significant), the direction of enhanced growth (vertical vs. horizontal), and the stability of treatment changes (unstable vs. stable). The short-term or long-term post-treatment relapse mainly relates to the rebound of dental position.

Conclusion – The immediate effects of bite jumping functional appliances on the mandibular growth enhancement are convincing during actual treatment. This extra gain of growth might be sustainable during the short-term and long-term post-treatment period.

Key words: bite jumping; growth modification; mandible

Introduction

Severe overjet is a common malocclusion and its prevalence is reported to be 10% in Chinese (1) and 15% in Caucasians (2). This malocclusion often reflects on a disharmony in the jaw relationship, i.e. the average pattern in Chinese with severe overjet is a protrusive maxilla and a retrusive mandible (3). In Caucasians, it has been reported that increased overjet more often is because of a retrusive mandible only (4,5). Since the mandible is too far back or too small in most patients with severe overjet, one sensible orthodontic treatment option in growing patients is to bring the mandible forward, aiming to enhance growth of the condyle (6,7) and remodeling of the glenoid fossa (8,9). The vertical dimension differs among the patients with severe overjet, from decreased to increased lower facial height. The mandibular and facial growth pattern varies within this group of patients, from favorable vertical-forward condylar growth direction and clockwise rotation of the mandible, to unfavorable sagittal condylar growth direction and counter-clockwise rotation of the mandible (10,11). The growth direction of the posterior cranial base varies and results in an open or closed skull base angle. Subsequently, the glenoid fossa can be in a more posterior or anterior position, which decides the position of the mandible. Therefore, a retrusive mandible can be of 'normal' size with its condyle in a glenoid fossa which is located posteriorly, or of a small size and/or abnormal shape with its condyle in a glenoid fossa which is located more anteriorly. In conclusion, patients with severe overjet are not homogenous in either overall facial or mandibular morphology or growth pattern. This will, at least to some extent, explain the often considerable individual variations in response to various orthodontic treatment mechanisms, such as bite-jumping therapy using functional appliances (12–14).

Since the late 19th century many types of bite jumping functional appliances (BJFA) have been advocated to treat growing patients (15). The fundamental principle for all BJFA is to keep the mandible in protrusive position in an attempt to evoke condylar and then mandibular growth, which in turn consolidates the repositioned mandible. The removable BJFA are mainly of activators, bionator, Clark twin block and Fränkel functional regulator II (FR II) with various modifications (15). For fixed BJFA, the fundamental

design and biomechanics are well represented by Herbst appliance that was originated in as early as 1905 and rejuvenated in the early 1970s by Pancherz. The telescope mechanism creates a continuous mandibular protrusion.

The 'classical functional appliances' are loose fitting appliances (e.g. Monoblock, Andresen activator, Herren activator, Harvold activator, Woodside activator). During the past 30 years, several authors recommended that functional appliances be attached to the dental arches with clasps (e.g. twin block, Bass appliance, Hansa Platte) and/or be held to the maxillary dental arch via occipital headgear (e.g. Headgear activator ad modum Teuscher, Bass appliance, Headgear activator ad modum Van Beek, Hansa Platte). The authentic fixed functional appliances, e.g. Herbst appliance, are banded or splinted (acrylic and/or casting) to the dental arch and therefore somewhat restrict mandibular lateral movement (16). The use of resilient push rod in Jasper–Jumper allows for mandibular excursion but somewhat compromises solid mandibular protrusion (17). The recent renovations and clinical application of fixed BJFA, e.g. Twin Force Bite Corrector (18,19), have shown clinical advantages mainly in two aspects: 1) the coil-spring mechanism by using superior quality of NiTi alloy secures a forcible advancement of the mandible with sufficient flexibility of mandibular functional movement, and 2) the direct and easy placement of the appliance into upper first molar and lower canine significantly simplifies the clinical procedure and reduces patient's discomfort. However, the simplicity in appliance design has somehow weakened the anchorage reinforcement which may arouse the concern over the dental compensation in lower labial segment.

In spite of a widespread application of bite jumping treatment for severe overjet malocclusion, controversy remains on its clinical efficacy in mandibular growth enhancement. The following three factors might account for this controversy: 1) the reports on the success degree of the skeletal effects are very much varying, 2) the parameters adopted to quantify the skeletal changes are often created and determined by the individual researchers and are not of consistence and compatibility between studies (20), and 3) it is extremely difficult, for reasons of ethics and patient's compliance, to design and implement a randomized controlled trial only from which could we seek the

truth. This review, with its focus extensively on clinical studies, was conducted to comprehend and analyze the data in literature based upon the categorization of removable and fixed BJFA, and to explore their respective effects on mandibular growth modification under the circumstances of various treatment timing (early and late), treatment duration and post-treatment follow-up (short- and long-term).

Mode of bite jumping

Proportion in vertical opening and horizontal protrusion

The mandible's working mode depends on the 'construction bite' which decides its position in the sagittal and vertical planes, and also possibly in the transversal plane. In vertical dimension, the magnitude of the construction bite (low, moderate and high) determines the amount of bite opening (i.e. the amount of downward position of the condyle). Andresen's and Häupl's original concept maintains that condylar adaptation is induced by a loose appliance in which mandible is anteriorly displaced with no obvious bite opening. Based on this concept, it has been advocated that the vertical bite opening should be limited within 4 mm, or a minimal magnitude which is not beyond postural rest position (21). Contrary to the concept of minimal vertical opening, some authors (22–24) hold the view that viscoelastic properties of muscle and the stretching of soft tissues are decisive for activator action. This primarily requires dislocating the mandible not only in an anterior positioning, but also creating bite opening up to 10–15 mm, well beyond the postural rest position (24). Between these two extremes, some others (25,26) suggest a modest bite opening of 4–6 mm, in an attempt to achieve the combined isometric muscle contraction and soft tissue stretching.

In the sagittal plane, the mandible might be advanced with varied magnitudes, from minimal (21), moderate (27), incisal edge-to-edge (28,29) to even reverse overjet (30). The significant difference in growth stimulation on condyle/mandible has been claimed between varied magnitudes of horizontal protrusion (31), and insignificant difference has also been reported (32).

Some authors (25,33) contend that the proportion of bite jumping magnitude in vertical and horizontal direction cannot be individually considered,

emphasizing the importance of the correlation or link between vertical opening and horizontal protrusion. That is, the magnitude of horizontal protrusion should be determined upon the consideration of that of vertical opening, and vice versa. When the magnitude of the forward position is large (7 or 8 mm), the vertical opening should be small (2–4 mm) so as not to overstretch the muscles. When the bite opening exceeds 6 mm, mandibular protrusion should be very moderate (3–5 mm) (33).

Activation of bite jumping

A single maximal advancement to edge to-edge relationship has been favored by some authors (34–36), claiming that a major initial advancement could best stimulate mandibular growth. Others (37–40) have advocated progressive small activations. There are indications from animal experimental studies (41) and clinical studies (42,43) that step-by-step advancement of the mandible results in a more forward position of the mandible compared with initial maximal jumping of the mandible. This could be justified by the animal experiments (41,44). In these studies, it was shown that an increase in activity of the lateral pterygoid muscle because of bite jumping was followed by an adaptive growth response at the condyle. The activity of the muscle decreases, however, after 6–8 weeks. By advancing the mandible several times during the treatment (i.e. every other month), the ligaments attached to the condyle were stretched repeatedly, leading to further increase in muscle activity followed by possible new condylar adaptation. By comparing the treatment effects between Bass appliance and Herbst appliance, Bendeus *et al.* (45) also recognize that stepwise activation of mandibular advancement might evoke more skeletal changes than does single activation.

The effects of bite jumping on TMJ and mandibular growth

During normal growth, the mandible is translated downward and forward as the actual growth occurs at the mandibular condyle and along the posterior surface of the ramus. The body of the mandible grows longer by periosteal apposition of bone on its posterior surface, while the ramus grows higher by endochondral replacement at the condyle, together with the

remodeling of glenoid fossa straight down and posteriorly (46). When the mandible is held in a forward position by BJFA, the condyle is brought downward and forward from its original position. The ligaments of the disc attaching to the posterior aspects of both condyle and the glenoid fossa are stretched and affect the tissue involved. In animal experiments, it has been shown that the proliferation of chondrocytes in condylar cartilage increases, and the bone deposition in posterior of glenoid fossa is evident (47,48).

The mesenchymal cells within the articular layer of both condyle and glenoid fossa are main source for bone formation. In an attempt to elucidate the mechanism by which mesenchymal cells proliferate and differentiate in response to mandibular protrusion, Rabie *et al.* (49) identified the temporal sequence of cellular changes in posterior aspect of TMJ in rats. The mesenchymal cells were found to be stretched and oriented in the direction of the pull, which might trigger the biophysiological path of mesenchymal cells differentiating into bone making cells in TMJ.

There have been reports on animal experiments that deny any substantial adaptive changes in TMJ in response to bite jumping (50,51). However, the recent studies at molecular and genetic level have predominantly confirmed an increased endochondral ossification in the condyle of growing rats in response to mandible protrusion, by examining the growth markers such as type X collagen, VEGF and neovascularization – related factors (52–54).

Removable BJFA

The effect of the removable BJFA on mandibular growth is mainly expressed in a vertical direction (55–58). Mandibular growth in vertical dimension is beneficial to the correction of class II malocclusion with an anteriorly rotating growth pattern of the mandible because this tends to open the bite and direct mandibular growth in a more vertical direction (33). The mechanism of the growth stimulation in the vertical component might reflect in the bite opening beyond the vertical dimension of postural rest, which is facilitated by vertical construction bite of the appliance.

Many clinical studies, however, reported that the effect on the mandible with removable BJFA is insignificant and no more than what could be attributed to normal growth; and the treatment changes are only

dento-alveolar (45,58–60). The results from clinical studies on the effect of activator treatment seem to indicate that mandibular growth is not affected by the treatment, i.e. the mandibular growth is no difference from the normal as a result of treatment (61). The clinical success rate seems to be dependent on the selection of the cases, e.g. the success rate for 'suitable activator cases' has been reported to be 50–60% (62). Unsuccessful treatment results in this sample of 'suitable activator cases' were because of 'lack of cooperation' and 'unfavorable growth pattern' (62).

In an attempt to master unfavorable downward growth of the maxilla and enhance condylar growth and remodeling of the glenoid fossa, the amount of the mandibular displacement was increased by making the construction bite considerably higher, well beyond the rest position of the mandible (23,24). It was assumed that by stretching the muscles well beyond rest position of the mandible, sufficient force should be transmitted via the appliance to the maxilla to restrain its downward growth. This attempt to control vertical growth of the maxilla and enhance mandibular growth was unsuccessful (23). Headgear was added to activators for the same reason, resulting in some restrain effect on maxillary forward and downward growth only, and the mandibular effect showed no difference from normal growth (38,45,63).

Fixed BJFA

The key differences in mode of bite jumping between removable and fixed appliances lie in the duration of mandibular protrusion and the magnitude of vertical bite opening. In fixed appliances, the continuation of bite jumping is secured but the dimension in vertical bite opening is limited. These features might lead to skeletal adaptation that differs from that caused by removable functional appliances.

The treatment effects of Herbst appliance on mandibular growth have been reported in the past two decades. These findings might also be comparable with the fixed appliance of other types (i.e. Jasper Jumper), on the grounds that they share similar mechanisms of action; treatment effects produced by the flexible force module presumably are similar to those of the Herbst appliance (17).

The Herbst appliance has been shown to enhance mandibular growth (6,35,64–66). It accelerates the

growth of the condyles and result in a change in the growth direction of the condyles, in most cases, into a more sagittal direction (67,68). Sagittal condylar growth was increased while vertical condylar growth was unaffected by Herbst appliance therapy (69). This differs from the removable appliances which might affect mandibular or condylar growth dominantly in the vertical direction (55,56). In fabrication of removable functional appliance, the bite construction with proper dimension of vertical opening is a necessity to stimulate the condylar growth in vertical direction. The Herbst appliance, on the other hand, brings the mandible forward with limited inferior displacement. This might be attributed to the increased condylar growth dominating in horizontal direction.

The effects of bite jumping related to the timing of treatment

Removable BJFA

The pattern of the mandibular growth curve follows that of the general growth curve (70). Mandibular growth is characterized by an adolescent growth spurt and its peak closely coincides with that of the maxilla and general growth (70). The pubertal growth period is the most favorable time to attack many orthodontic problems with skeletal manifestations. When correction of the malocclusion has been achieved, the patient would reach the late adolescent growth period in which the growth rate would slow down.

For removable appliances, however, the treatment is preferably started in the pre-adolescent period for two reasons: 1) The treatment time with removable appliances is usually as lengthy as 2 years. The commencement of treatment, therefore, should be planned prior to the pubertal growth period, in order to overlap as much treatment time with pubertal growth as possible (71), and 2) Second-phase treatment with fixed appliance is usually needed upon the fulfillment of removable orthopedic intervention, to accomplish the entire correction during the adolescence years (72). Therefore, the treatment timing for orthopedic intervention with removable functional appliances should be ahead of pubertal growth, in order to secure that the late orthodontic treatment with fixed appliances could share part of pubertal growth period.

Dentofacial orthopedic changes were believed to be more dependent on the adaptability of skeletal tissues than the dentofacial growth (73). The dentofacial tissue was more adaptive at a younger age and the tissue adaptability decreased with increase of age. Early treatment with removable functional appliances at pre-adolescent years has been claimed to cause increased growth in mandible (71,73,74).

The commencement of treatment prior to the pubertal growth spurt, however, imposes disadvantages. Early treatment faces the risk of unpredictable post-treatment growth changes (75,76). The long duration of potential jaw growth after the active treatment would pose an uncertainty on the stability of the corrected occlusion (77).

Fixed BJFA

Unlike removable functional appliances which require a lengthy treatment duration, the fixed bite jumping appliances (e.g. Herbst appliance), impose a short treatment time of 6–12 months (78). This leads to the considerable flexibility in the selection of treatment timing. On a short-term basis, the most favorable time to treat patients with Herbst appliance is at, or just after, the peak of the pubertal growth spurt (64,65,79). At this time the influence on mandibular condylar growth is greatest and the risk of undesirable dental effects on the mandible (e.g. proclination of the lower incisors) is comparatively small (64,78). Taking into consideration other factors, such as the long-term stability of treatment results and the efficiency of retention, the ideal period for starting treatment should be when the permanent dentition stage has been reached and at, or just after, peak height velocity of growth. It was claimed that this would promote occlusal stability after treatment and shorten the retention period (77).

However, the amount of enhanced mandibular growth seems to be similar regardless the commencement time of the treatment in growing individuals (79). The major difference in mandibular treatment changes during various growth periods seems to be mainly because of the variations in the basic mandibular growth rate (79). At long-term follow-up there seems to be no difference in length of the mandible between patients treated in different growth periods (80).

The effects of bite jumping related to post-treatment follow-up

There are only a few reports on the short-term and long-term effects of removable appliances. This might be because of the difficulties in recalling the patients for re-evaluation. However, the evaluation for short-term and long-term effects of Herbst appliances has been documented extensively (16,78). The amount of gain in mandibular growth during treatment is small compared with the total growth after treatment; the individual variations of both parameters are large, and the sample size is sometimes too small and not homogenous, which makes it impossible to reveal any possible long-term effects of mandibular bite jumping (16,78).

The treatment effects on mandibular growth a short-term period after treatment with removable functional appliances have been reported to be stable (81) or to decline (82). Nelson *et al.* (60), however, found no growth stimulation on a short-term basis. A limited number of studies on long-term effects of removable BJFA on mandibular growth have almost invariably reported some relapse after 3–20 years after treatment (12,36,83,84). A recently reported long-term (up to 4 years) follow-up study on twin block appliances claimed that much of the significant increase in mandibular length achieved during the active phase of treatment was still present after 3–4 years when the subjects had matured into permanent dentition stage (85).

For fixed functional appliances (e.g. Herbst appliance), the direction and the amount of mandibular growth can be temporarily altered, but return to their pre-treatment pattern after treatment on short-term basis (76,86). The dentoalveolar and skeletal relapse, which occurred primarily during the first 6 months after treatment, was about 30% of the accomplished treatment effect and was mainly of dental origin (13,67). This is further confirmed by a recent randomized controlled trial on headgear Herbst with step-by-step activated mandibular advancement (66). In the study, it was found that a significant enhancement on mandibular growth occurred during the initial treatment phase (6 months), which declined in the late treatment phase (12 months) and after retention phase (18 months). The corrected jaw relationship maintained over this short-term (12 months) and a small

relapse of overjet and molar relationship was the result of dental changes only. On a long-term basis, the clinically significant effects on mandible growth remain despite that mandibular growth appears to return to its earlier pattern after treatment (76). Long-term studies, i.e. 5–10 years follow-up, showed that the mandible effects remained and the reason for relapse was the changes in tooth position (77). In the Herbst-treated patients, the sagittal arch relationship is generally stable and comparable with normal and 'ideal' samples from the Bolton Standards (87). Some studies indicate that mandibular growth is enhanced during active treatment compared with controls, but during the follow-up period the mandibular growth seems to slow down in BJFA patient group. Subsequently in long-term there is no obvious increase in mandibular length even in the samples of patients with an initially favorable response to bite jumping, compared with controls (61).

Consequently, it appears that the unfavorable post-treatment changes occurring after the removal of removable or fixed BJFA, are dominantly dental related. This might be because that the long-term effect of treatment on mandibular growth is difficult to assess, and the increase in sagittal condylar growth and the changes in mandibular morphology seen during treatment could not be verified several years after the therapy had been ceased (42). Any orthopedic appliances seem to have only a temporary impact on the existing skeletofacial growth pattern (i.e. growth rate and growth direction). After the orthopedic intervention period, mandibular growth seems to return to its previous patterns (67). However, these accelerated growth changes are maintained by the additional skeletal growth and dentitional changes (85).

References

1. Fu M, Zhang D, Wang B, Deng Y, Wang F, Ye X. The prevalence of malocclusion in China – an investigation of 25,392 children (in Chinese). *Chin J Stomatol* 2002;**37**:371–3.
2. Mitchell L. *An Introduction to Orthodontics*, 2nd edn. New York: Oxford University Press Inc.; 2001.
3. Lau JWP, Hägg U. Cephalometric morphology of Chinese with Class II Division 1 malocclusion. *Bri Dent J* 1999;**186**:188–90.
4. McNamara JA. Components of Class II malocclusion in children 8–10 years of age. *Angle Orthod* 1981;**51**:177–202.
5. Pancherz H, Zieher K, Hoyer B. Cephalometric characteristics of Class II Division 1 and Class II Division 2 malocclusions: a comparative study in children. *Angle Orthod* 1997;**67**:111–20.

6. Pancherz H. Treatment of Class II malocclusions by jumping the bite with the Herbst appliance: a cephalometric investigation. *Am J Orthod* 1979;**76**:423–42.
7. Vargervik K, Harvold EP. Response to activator treatment in Class II malocclusions. *Am J Orthod* 1985;**88**:242–51.
8. McNamara JA, Carlson DS. Quantitative analysis of temporomandibular joint adaptations to protrusive function. *Am J Orthod* 1979;**76**:593–611.
9. Hinton RJ, Carlson DS. Histological changes in the articular eminence and mandibular fossa during growth of the rhesus monkey (*Macaca mulatta*). *Am J Anat* 1983;**166**:99–116.
10. Björk A. Variations in the growth pattern of the human mandible: longitudinal radiographic study by the implant method. *J Dent Res* 1963;**42**:400–11.
11. Stockli PW, Dietrich UC. Sensation and morphogenesis: experimental and clinical findings following functional forward displacement of the mandible. *Trans Eur Orthod Soc* 1973;435–42.
12. Ahlgren J. A longitudinal clinical and cephalometric study of 50 malocclusion cases treated with activator appliances. *Trans Eur Orthod Soc* 1972;285–93.
13. Pancherz H, Hansen K. Occlusal changes during and after Herbst treatment: a cephalometric investigation. *Eur J Orthod* 1986;**8**:215–28.
14. Wong GW, So LL, Hägg U. A comparative study of sagittal correction with the Herbst appliance in two different ethnic groups. *Eur J Orthod* 1997;**19**:195–204.
15. Graber TM, Rakosi T, Petrovic AG. *Dentofacial Orthopedics with Functional Appliances*, 2nd edn. St Louis: Mosby-Year Book; 1997.
16. Pancherz H. *The Herbst Appliance*. Antonio Reyes Lara: Editorial Aguiram; 1995.
17. Jasper JJ, McNamara JA, Mollenhauer B. The modified Herbst appliance (Jasper Jumper). In: Graber TM, Rakosi T, Petrovic AG, editors. *Dentofacial Orthopedics with Functional Appliances*, 2nd edn. St Louis: Mosby-Year Book; 1997. pp. 367–75.
18. Mahony D. Twin Force Bite Corrector – hyper efficient Class II correction for a busy orthodontic practice. *Int J Orthod* 2003;**14**:9–14.
19. Rothenberg J, Campbell ES, Nanda R. Class II correction with Twin Force Bite Corrector. *J Clin Orthod* 2004;**38**:232–40.
20. Chen JY, Will LA, Niederman R. Analysis of efficacy of functional appliances on mandibular growth. *Am J Orthod Dentofac Orthop* 2002;**122**:470–6.
21. Falck F, Fränkel R. Clinical relevance of step-by-step mandibular advancement in the treatment of mandibular retrusion using the Frankel appliance. *Am J Orthod Dentofac Orthop* 1989;**96**:333–41.
22. Heren D. The activator's mode of action. *Am J Orthod* 1959;**45**:512–27.
23. Harvold EP, Vargervik K. Morphogenetic response to activator treatment. *Am J Orthod* 1971;**60**:478–90.
24. Woodside DG. Some effects of activator treatment on the mandible and midface. *Trans Eur Orthod Soc* 1973;443–47.
25. Witt E. Basic principles of activator and bionator therapy. *Zahnärztliche Praxis* 1971;**22**:1–526.
26. Schmuth GP. Milestones in the development and practical application of functional appliances. *Am J Orthod* 1983;**84**:48–53.
27. Owen AH. The maxillary sagittal appliance: a clinical study. *Am J Orthod Dentofac Orthop* 1987;**91**:271–85.
28. Carels C, van Steenberghe D. Changes in neuromuscular reflexes in the masseter muscles during functional jaw orthopedic treatment in children. *Am J Orthod Dentofac Orthop* 1986;**90**:410–9.
29. Carels C, van der Linden FPGM. Concepts of functional appliances mode of action. *Am J Orthod Dentofac Orthop* 1987;**92**:162–8.
30. Demisch A. Effects of activator therapy on the craniofacial skeleton in Class II, Division 1 malocclusion. *Trans Eur Orthod Soc* 1972;295–310.
31. Heij Op DG, Callaert H, Opdebeek HM. The effect of amount of protrusion built into the Bionator on condylar growth and displacement: a clinical study. *Am J Orthod Dentofac Orthop* 1989;**95**:401–9.
32. DeVincenzo JP, Winn MW. Orthopedic and orthodontic effects resulting from the use of a functional appliance with different amounts of protrusive activation. *Am J Orthod Dentofac Orthop* 1989;**96**:181–90.
33. Rakosi T. The activator. In: Graber TM, Rakosi T, Petrovic AG, editors. *Dentofacial Orthopedics with Functional Appliances*, 2nd edn. St Louis: Mosby; 1997. pp. 161–213.
34. Reey RW, Eastwood A. The passive activator: case selection, treatment response, and corrective mechanics. *Am J Orthod* 1978;**73**:378–409.
35. Pancherz H. The mechanism of Class II correction in Herbst appliance treatment. A cephalometric investigation. *Am J Orthod* 1982;**82**:104–13.
36. Madone G, Ingervall B. Stability of results and function of the masticatory system in patients treated with the Herren type of activator. *Eur J Orthod* 1984;**6**:92–106.
37. McNamara JA, Hugu SA. The Frankel appliance (FRII): model preparation and appliance construction. *Am J Orthod* 1981;**80**:478–95.
38. Bass NM. Dentofacial orthopaedics in the correction of Class II malocclusions. *Bri J Orthod* 1982;**9**:3–31.
39. Fränkel R. Biomechanical aspects of the form/function relationship in craniofacial morphogenesis: a clinical approach. In: McNamara JA, Ribbens KA, Howe RP, editors. *Clinical Alteration of the Growing Face. Monograph No.14, Craniofacial Growth Series*. Ann Arbor: Center for Human Growth and Development, The University of Michigan; 1983. pp. 107–30.
40. Malmgren O, Omblus J. Treatment with an orthopaedic appliance system. *Eur J Orthod* 1985;**7**:205–14.
41. McNamara JA, Connelly TG, McBride MC. Histological studies of temporomandibular joint adaptations. In: McNamara JA, editors. *Determinants of Mandibular Form and Growth*. Ann Arbor: Center for Human Growth and Development, The University of Michigan; 1975. pp. 209–27.
42. Pancherz H, Littmann C. Morphology and position of mandible in Herbst treatment. Cephalometric analysis of changes to end of growth period. *Inf Orthod Kieferorthop* 1989;**21**:493–513.
43. Du X., Hägg U., Rabie AB. Effects of headgear Herbst and mandibular step-by-step advancement versus conventional Herbst appliance and maximal jumping of the mandible. *Eur J Orthod* 2002;**24**:167–74.
44. Petrovic AG, Stutzmann JJ, Oudet CL. Control processes in the postnatal growth of the condylar cartilage of the mandible. In: McNamara JA, editor. *Determinants of Mandibular Form and Growth. Monograph No. 4, Craniofacial Growth Series*. Ann Arbor: Center for Human Growth and Development, The University of Michigan, 1975. pp. 101–54.
45. Bendeus M, Hägg U, Rabie B. Growth and treatment changes in patients treated with a headgear-activator appliance. *Am J Orthod Dentofac Orthop* 2002;**121**:376–84.

46. Proffit WR, Ackerman JL. Orthodontic diagnosis: the development of a problem list. In: Proffit WR, Fields HW, editors. *Contemporary Orthodontics*. St Louis: Mosby Year Book; 1993. pp. 139–86.
47. McNamara JA, Hinton RJ, Hoffman DL. Histological analysis of temporomandibular joint adaptation to protrusive function in young adult rhesus monkeys (Macaca Mulatta). *Am J Orthod* 1982;**82**:288–98.
48. Hinton RJ, McNamara JA. Temporal bone adaptations in response to protrusive function in juvenile and young adult rhesus monkeys (Macaca mulatta). *Eur J Orthod* 1984;**6**:155–74.
49. Rabie AB, Zhao Z, Shen G, Hagg EU, Robinson W. Osteogenesis in the glenoid fossa in response to mandibular advancement. *Am J Orthod Dentofac Orthop* 2001;**119**:390–400.
50. Tonge E, Heath JK, Meikle MC. Anterior mandibular displacement and condylar growth: an experimental study in the rat. *Am J Orthod* 1982;**82**:277–87.
51. Tewson DHTK, Heath JK, Meikle MC. Biochemical and autoradiographical evidence that anterior mandibular displacement in the young growing rat does not stimulate cell proliferation or matrix formation at the mandibular condyle. *Arch Oral Biol* 1988;**33**:99–107.
52. Rabie AB, Hägg U. Factors regulating mandibular condylar growth. *Am J Orthod Dentofac Orthop* 2002;**122**:401–9.
53. Shen G, Rabie AB, Hägg U, Chen RJ. Neovascularization in condylar cartilage in response to mandibular protrusion. *Chin J Dent Res* 2003;**6**:28–38.
54. Rabie AB, Shum L, Chayanupatkul A. VEGF and bone formation in the glenoid fossa during forward mandibular positioning. *Am J Orthod Dentofac Orthop* 2002;**122**:202–9.
55. Luder HU. Skeletal profile changes related to two patterns of activator effects. *Am J Orthod* 1982;**8**:390–6.
56. Creekmore TD, Radney LJ. Frankel appliance therapy: orthopedic or orthodontic? *Am J Orthod* 1983;**83**:89–108.
57. McNamara JA, Bookstein FL, Shaughnessy TG. Skeletal and dental changes following functional regulator therapy on Class II patients. *Am J Orthod* 1985;**88**:91–110.
58. Mills JR. The effect of functional appliances on the skeletal pattern. *Bri J Orthod* 1991;**18**:267–75.
59. Wieslander L, Lagerstrom L. The effect of activator treatment on Class II malocclusions. *Am J Orthod* 1979;**75**:20–6.
60. Nelson C, Harkness M, Herbison P. Mandibular changes during functional appliance treatment. *Am J Orthod Dentofac Orthop* 1993;**104**:153–61.
61. Johnston LE. Growing jaws for fun and profit: a modest proposal. In: McNamara JA, editor. *Growth Modification: What Works, What Doesn't, and Why*. Ann Arbor: Center for Human growth and Development, The University of Michigan; 1999. pp. 63–86.
62. Ahlgren J, Laurin C. Late results of activator treatment: a cephalometric study. *Bri J Orthod* 1976;**3**:181–7.
63. Van Beek H. Overjet correction by a combined headgear and activator. *Eur J Orthod* 1982;**4**:279–90.
64. Pancherz H, Hägg U. Dentofacial orthopedics in relation to somatic maturation. An analysis of 70 consecutive cases treated with the Herbst appliance. *Am J Orthod* 1985;**88**:273–87.
65. Hägg U, Pancherz H. Dentofacial orthopaedics in relation to chronological age, growth period and skeletal development. An analysis of 72 male patients with Class II Division 1 malocclusion treated with the Herbst appliance. *Eur J Orthod* 1988;**10**:169–76.
66. Hägg U, Du X, Rabie AB. Initial and late treatment effects of headgear-Herbst appliance with mandibular step-by-step advancement. *Am J Orthod Dentofac Orthop* 2002;**122**:477–85.
67. Pancherz H, Anehus-Pancherz M. The headgear effect of the Herbst appliance: a cephalometric long-term study. *Am J Orthod Dentofac Orthop* 1993;**103**:510–20.
68. Paulsen HU. Morphological changes of the TMJ condyles of 100 patients treated with the Herbst appliance in the period of puberty to adulthood: a long-term radiographic study. *Eur J Orthod* 1997;**19**:657–68.
69. Pancherz H, Littmann C. Somatic maturity and morphologic changes of the mandible with Herbst treatment. *Inf Orthod Kieferorthop* 1988;**20**:455–70.
70. Björk A, Skieller V. An implant study at the age of puberty. *Am J Orthod* 1972;**62**:339–83.
71. Wieslander L. Early or late cervical traction therapy of Class II malocclusion in the mixed dentition. *Am J Orthod* 1975;**67**:432–9.
72. King GJ, Keeling RD, Hocevar RA, Wheder TT. The timing of treatment for Class II malocclusion in children: a review. *Angle Orthod* 1990;**60**:87–97.
73. Van der Linden FPGM. 1986 *Facial Growth and Facial Orthopedics*. Kinston-upon-Thames, Surrey: Quintessence; 1986.
74. Tulloch JF, Phillips C, Koch G, Proffit WR. The effect of early intervention on skeletal pattern in Class II malocclusion: a randomized clinical trial. *Am J Orthod Dentofac Orthop* 1997;**111**:391–400.
75. Melsen B. Effects of cervical anchorage during and after treatment: an implant study. *Am J Orthod* 1978;**73**:526–40.
76. Pancherz H, Fackel U. The skeletofacial growth pattern pre- and post-dentofacial orthopaedics. A long-term study of Class II malocclusions treated with the Herbst appliance. *Eur J Orthod* 1990;**12**:209–18.
77. Pancherz H. The nature of Class II relapse after Herbst appliance treatment: a cephalometric long-term investigation. *Am J Orthod* 1991;**100**:220–33.
78. Pancherz H. The modern Herbst appliance. In: Graber TM, Rakosi T, Petrovic AG, editors. *Dentofacial Orthopedics with Functional Appliances*, 2nd edn. St Louis: Mosby; 1997. pp. 336–66.
79. Hägg U, Pancherz H, Taranger J. Pubertal growth and orthodontic treatment. In: Carlson DS, Ribbens KA, editors. *Craniofacial Growth During Adolescence. Monograph No.14, Craniofacial Growth Series*. Ann Arbor: Center for Human Growth and Development, The University of Michigan; 1987. pp. 87–115.
80. Hansen K, Pancherz H, Hägg U. Long-term effects of the Herbst appliance in relation to the treatment growth period: a cephalometric study. *Eur J Orthod* 1991;**13**:471–81.
81. Keeling SD, Wheeler TT, King GJ, Garvan CW, Cohen DA. Anteroposterior skeletal and dental changes after early Class II treatment with bionators and headgear. *Am J Orthod Dentofac Orthop* 1998;**113**:40–50.
82. DeVincenzo JP. Changes in mandibular length before, during, and after successful orthopedic correction of Class II malocclusions, using a functional appliance. *Am J Orthod Dentofac Orthop* 1991;**99**:241–57.
83. Pancherz H. Long-term effects of activator (Andresen appliance) treatment. A clinical, biometric, cephalometric roentgenographic and functional analysis. *Odontologisk Revy Suppl* 1976;**35**:1–70.

84. Drage KJ, Hunt NP. Overjet relapse following functional appliance therapy. *Bri J Orthod* 1990;**17**:205–13.
85. Mills CM, McCulloch KJ. Posttreatment changes after successful correction of Class II malocclusions with the twin block appliance. *Am J Orthod Dentofac Orthop* 2000;**118**:24–33.
86. Hägg U. Change in mandibular growth direction by means of a Herbst appliance? A case report. *Am J Orthod Dentofac Orthop* 1992;**102**:456–63.
87. Hansen K, Pancherz H. Logn-term effects of Herbst treatment in relation to normal growth development: a cephalometric study. *Eur J Orthod* 1992;**14**:285–95.

Copyright of Orthodontics & Craniofacial Research is the property of Blackwell Publishing Limited and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.