



L. M. Moreno Uribe
S. F. Miller

Genetics of the dentofacial variation in human malocclusion

Authors' affiliations:

L. M. Moreno Uribe, Department of Orthodontics, College of Dentistry, University of Iowa, Iowa City, IA, USA
L. M. Moreno Uribe, S. F. Miller, Dows Institute for Dental Research, College of Dentistry, University of Iowa, Iowa City, IA, USA

Correspondence to:

L. M. Moreno Uribe
N401 DSB
University of Iowa
Iowa City, IA 52242
USA
E-mail: lina-moreno@uiowa.edu

Moreno Uribe L. M., Miller S. F. Genetics of the dentofacial variation in human malocclusion

Orthod Craniofac Res 2015; **18**(Suppl. 1): 91–99. © 2015 John Wiley & Sons A/S. Published by John Wiley & Sons Ltd

Abstract Malocclusions affect individuals worldwide, resulting in compromised function and esthetics. Understanding the etiological factors contributing to the variation in dentofacial morphology associated with malocclusions is the key to develop novel treatment approaches. Advances in dentofacial phenotyping, which is the comprehensive characterization of hard and soft tissue variation in the craniofacial complex, together with the acquisition of large-scale genomic data have started to unravel genetic mechanisms underlying facial variation. Knowledge on the genetics of human malocclusion is limited even though results attained thus far are encouraging, with promising opportunities for future research. This review summarizes the most common dentofacial variations associated with malocclusions and reviews the current knowledge of the roles of genes in the development of malocclusions. Lastly, this review will describe ways to advance malocclusion research, following examples from the expanding fields of phenomics and genomic medicine, which aim to better patient outcomes.

Key words: candidate genes; cephalometry; craniofacial; genetics; genomic medicine; genotype–phenotype correlation; phenomics

Introduction

Human malocclusion is a disarrangement of teeth and jaws that may lead to distorted facial appearance, limited masticatory function, increased risk for dental trauma, and compromise quality of life (1, 2).

The interactions of genetic and environmental factors may account for the variability in expression of malocclusion. The etiological complexity lies not only in unpredictable expression, but also in the wide spectrum of dentofacial variation present in affected individuals. This complexity explains in part why most

Date:

Accepted 14 October 2014

DOI: 10.1111/ocr.12083

© 2015 John Wiley & Sons A/S.
Published by John Wiley & Sons Ltd

treatment approaches for malocclusion are directed to the symptoms rather than to etiology. However, despite this complexity, the study of malocclusion etiology is fundamental to understanding the biology underlying craniofacial growth and dental relations. Understanding the biology will aid progress toward effective treatment, prevention, thereby decreasing the burden of this condition.

Dentofacial variation in human malocclusion

Given the abundance of hard and soft tissue information contained within lateral cephalometric images, most phenotypic characterization has been in two dimensions. Different analytical methods have been employed, including shape analyses, and principal components and cluster analyses. The following is a brief review of these studies.

Class II

Longitudinal studies indicated that Class II dentoskeletal characteristics can appear during the primary dentition (3). Although catch up growth can occur in some individuals, such discrepancies in general do not self-correct due to differences in growth magnitudes and directions between individuals with Class II and Class I

malocclusions (4). The most recent study of Class II variation evaluated 309 Class II Caucasian adults and resulted in seven principal components explaining 81% of the variation. About half of this variation was depicted by vertical mandibular rotation, incisor angulations and the size of the ramus and body of the mandible (Fig. 1). Moreover, five distinct clusters representing the spectrum of Class II phenotypes were identified, thereby highlighting how variation of a limited number of principal components affect the craniofacial complex (5) (Fig. 2).

Class III

Similarly to the class II malocclusion, class III malocclusion features affect multiple craniofacial structures, appear early in development, worsen with age, and are present in most Class III individuals regardless of ethnicity (6, 7). A recent study in 292 Caucasian adults identified six principal components which accounted for 81% of the phenotypic variation recorded. About 54% of the variation was explained by the anterior-posterior (AP) position of the mandible compared to the cranial base, the size of the maxillomandibular horizontal discrepancy, and the lower incisor AP position (Fig. 3). Cluster analyses identified five distinct phenotypic subgroups (8) (Fig. 4) which were remarkably similar to those found by previous studies (9).




Fig. 1. Individuals in the extreme opposite ends for the first three principal components explaining about 50% of the total variation in Class II malocclusion. PC1 refers to variation in the inclination of the mandibular plane angle. PC2 depicts maxillary incisor angulation. PC3 refers to the mandibular AP and vertical length and posterior facial height.




Fig. 2. Cluster analyses for Class II malocclusion showing dentofacial features of cluster centroids and their main features. Pro, profile; Cb, cranial base; Mx, maxilla; Md, mandible; Vert, vertical; U1, upper incisor; L1, lower incisor; St, soft tissue.




Fig. 3. Individuals in the extreme opposite ends for the first three principal components explaining about 54% of the total variation in Class III malocclusion. PC1 shows the anteroposterior position of the mandible in relationship to the cranial base. PC2 depicts the maxillomandibular horizontal and vertical size discrepancies. PC3 refers to the lower incisor and lower lip variation.




Fig. 4. Cluster analyses for Class III malocclusion showing dentofacial features of cluster centroids and their main features. Pro, profile; Cb, cranial base; Mx, maxilla; Md, mandible; Vert, vertical; U1, upper incisor; L1, lower incisor; St, soft tissue.

The studies described above have demonstrated the utility of data reduction methods applied to multidimensional data to capture meaningful categorical and quantitative malocclusion phenotypes for future studies of malocclusion etiology.

Current knowledge on the genetics of human malocclusion

Multiple data sources suggest that genetic factors contribute to malocclusion susceptibility. Moderate to high heritability proportions (>60%) have been reported for many dental and facial features such as mid and lower facial dimensions, dental spacing, arch dimensions and Bolton type tooth size discrepancies. Conversely, overbite (53%) and overjet (28%) have lower heritability, which suggested a higher susceptibility to environmental factors (10, 11).

Familial aggregation studies suggested that an autosomal dominant model with incomplete penetrance has the greatest validity for Class III pedigrees, including the royal Habsburg family (12) and others from Middle Eastern (13), South American (14, 15), and Eastern European descent (16). In contrast, polygenic inheritance and autosomal dominance models, with incomplete penetrance and variable expressivity, have been suggested for Class II subdivision 1 and 2, respectively (17–19).

The frequent presentation of malocclusion in patients with craniofacial birth defects also supports a strong genetic etiology. About 150 genes/loci are associated with craniofacial conditions presenting malocclusion (OMIM <http://www.ncbi.nlm.nih.gov/omim>). These genes represent molecular pathways to explore in malocclusion etiology.

Similarly, about 275 mouse models of malocclusion have been cataloged, with 235 genes/loci identified in the MGI database (<http://www.informatics.jax.org/>), contributing important information for the etiology of human malocclusions. Recently, mouse studies showed that long-range transcriptional enhancers regulate the expression of genes near and far during

craniofacial development, resulting in subtle differences in craniofacial shape (20). Thus, these enhancers could be a mechanism to explain dentofacial variations underlying malocclusion.

Human genetic mapping approaches for identification of risk loci include linkage and association methods. Linkage analysis is applied to large families or affected sibling pairs and aims at detecting rare variants with large effects. Association analyses applied to case-control or case-parent triad samples are more powerful at detecting common variants with smaller effects (21). Currently, large-scale genomic data generation in the form of genomewide association scans (GWAS) and next-generation sequencing of full exomes and full genomes coupled with meta-analysis tools to identify common and rare genetic variants have significantly increased the scale and scope of complex trait mapping projects (22). However, despite this technological expansion, the identification of susceptibility genes for human malocclusion is just beginning.

Most malocclusion studies to date have focused in Class III malocclusion (Table 1). Although linkage and association findings thus far seem to cluster around chromosome 1 (loci 1p22-p36) and 12 (loci 12q13-q24), a study in 10 large Brazilian pedigrees failed to replicate linkage to chromosome 1 (23) underscoring the genetic heterogeneity of Class III malocclusions in different ethnicities.

Association studies have found positive correlations for mandibular prognathism and genes *EPB41*, *MATN1*, *SSX2IP*, and *PLXNA*, located within the 1p22-p36 locus. Also positive associations have been found for genes *COL2A1*, *MYO1H*, *TGFB3*, and *LTBP2* within the 12q13-q24 locus. Interestingly, variants in *Matn1* (1p35) were associated with mandibular prognathism and anterior cross bites in donkeys (24).

Recently, full exome sequencing of five siblings with maxillary hypoplasia identified a heterozygous missense mutation c.545C>T (p.Ser182Phe) in the gene *DUSP6* (12q21) (16). In a mouse model, *Dusp6* expression correlates with *Fgfrs* domains in the branchial arches and it is stimulated by *Fgf* signaling, a key morphogenetic pathway in the outgrowth of maxillary promi-

Table 1. Summary of mapping studies for Class III malocclusion

Region	Gene/loci	Method	Phenotype	Population	References
1p36		Linkage	Mand. prog.	40 Korean & 50 Japanese sibling pairs	(30)
1p35.3	EPB41	Association	Mand. prog.	211 cases vs. 224 controls Chinese	(31)
1p35.2	MATN1	Association	Mand. prog.	164 cases vs. 132 controls Koreans	(32)
1p22.3	SSX2IP	Association	Mand. prog.	240 cases vs. 360 controls	(33)
1p22.1		Linkage	Max def.	4 Hispanic families, Colombia	(34)
1q32.2	PLXNA2	Association	Mand. prog.	240 cases vs. 360 controls	(33)
3q26.2		Linkage	Max def.	4 Hispanic families, Colombia	(34)
4p16	EVC, EVC2	Linkage	Mand. Prog.	2 Large Chinese pedigrees	(35)
6q25		Linkage	Mand. Prog.	40 Korean & 50 Japanese sibling pairs	(30)
11q22		Linkage	Max def.	4 Hispanic families, Colombia	(34)
12q13.13	COL2A1	Linkage, Association	Max def., Mand. Prog.	4 Hispanic families, Colombia, 211 cases vs. 224 controls Chinese	(34, 36)
12q21.33	DUSP6	Causal/ etiologic	Max def.	1 Estonian family (5 siblings)	(16)
12q23	IGF1	Linkage	Max def.	4 Hispanic families, Colombia	(34)
12q24.11	MYO1H	Association	Mand. Prog.	44 cases vs. 36 controls Pittsburgh, USA, mostly Caucasian	(37)
14q24.3	TGFB3, LTBP2	Linkage	Mand. Prog.	1 Han Chinese large family	(38)
19p13.2		Linkage	Mand. Prog.	40 Korean & 50 Japanese sibling pairs	(30)

nences. Mice null for *Dusp6* exhibit skeletal dwarfism and craniosynostosis (25). It is possible that variants within *DUSP6* could account for Class III malocclusion due to maxillary hypoplasia subsequent to premature fusion of maxillary sutures.

Genetic studies for Class II and Class I malocclusion are more rare. In four Colombian families, individuals with mandibular hypoplasia were homozygous for the rare allele on SNP rs1348322, within the *Noggin* gene (26). This gene is essential for mandibular formation in mice (27). For Class I malocclusion, a reported SNP rs6504340 within the *HOXB* cluster was associated with delayed tooth eruption and occlusion irregularities that required orthodontic therapy (28). Also, significant associations between more than 5 mm of crowding and genes *EDA* (rs3764746 and rs3795170), *XEDAR* (rs372024), and *BMP2* (rs1005464) were reported in Class I Chinese subjects (29).

While informative, the studies above are limited by modest sample sizes, unknown generaliz-

ability of results to populations of other ancestries, and restrictive traits that ignore the complex phenotypic spectrum of malocclusion. Nonetheless, these studies have converged to highlight that genes implicated in bone (*TGFB3*, *LTBP*, *IGF1*, *ENPP1*, *EVC*, and *EVC2*), cartilage development (*Matrilin-1* and *COL2A1*), muscle function (*MYO1H* and *DUSP6*) and tooth morphogenesis (*EDA*, *XEDAR*, and *BMP2*), may be putative candidates for jaw and tooth size discrepancies.

Phenotypic resources for malocclusion studies

To understand causal mechanisms of human disease, a large emphasis has been placed in the field of phenomics, the comprehensive study of the full set of possible phenotypes on an individual (39). Comprehensive phenotypes in combination with large-scale genomic data maximize the efficiency at detecting genotype-phenotype

correlations of clinical importance. Consequently, phenomic initiatives and specimen repositories have been developed for various human conditions. The website dbGAP (<http://www.ncbi.nlm.nih.gov/gap>), for example, is a repository of phenotype–genotype correlation studies, which includes data on dental caries, oral clefts, and limited dental malocclusion variables. Also, the FaceBase consortium (<https://www.facebase.org/>) offers access to a central repository of 3D facial surface images and DNA resources for normative data. Also, dental phenomic initiatives for caries, periodontal disease, and dental morphology are underway (40). With respect to orthodontics, there is limited information available through orthodontic growth studies (e.g., http://www.aaflegacycollection.org/aaof_home.html) since no biological specimens were collected during the periods over which serial lateral cephalometric radiographs were produced. To date, none of the existing databases contains comprehensive dentofacial data for malocclusions. Therefore, phenotype–genotype correlation studies of malocclusion are greatly needed as the knowledge gained from them will aid in our understanding of the mechanisms responsible for human malocclusions and craniofacial anomalies.

The orthodontic community constitutes a great resource for phenomic data and biological specimens because of access to large patient pools with wide ranges of malocclusions and the routine use of diagnostic records. Of these, lateral cephalometric radiographs and clinical photographs are the most abundant and thus are likely to be the primary data source for large-scale genotype–phenotype correlation projects. Although 2D photographs have dimensional errors due to variations in projection and patient positioning, 2D photographs can still be utilized for facial phenotyping through estimates of facial proportions, angles, and shape analyses.

3D facial surface imaging offers more accurate data, without errors due to projection distortion or patient positioning. This imaging method increases the scope of facial variation studies (41). It can be used to detect soft tissue features specific to craniofacial conditions such as cleft

lip and palate (42). However, 3D facial surface images do not include dentoalveolar data, limiting its use in characterizing malocclusion phenotypes unless accompanied by cone-beam computed tomographic imaging (CBCT), or dentoalveolar data generated from scanned dental casts or intraoral scanners.

Despite the challenges of cost, storage, and analyses of deep phenotypic approaches, the advantages of generating malocclusion phenomic data will be numerous. For instance, comprehensive phenotyping may help predict individuals with adverse or favorable treatment responses, thus making practice more rewarding for clinicians. Also, comprehensive phenotyping could reduce problems with missing heritability in genetic studies, as will be discussed below.

Clinical relevance of phenotype–genotype studies of malocclusion

Candidate gene (42–44) and GWAS studies (45–47) of 3D facial soft tissue variation resulted in significant associations between genes *PRDM16*, *PAX3*, *TP63*, *C5orf50*, *Col17A1*, *HMGA2*, *AJUBA*, and *ADK*, and facial width and height. Also, genes and loci associated with oral clefts (*IRF6*, *8q24*, *SNAI1*, *MSX1*, *ABCA4-ARHGAP29*, and *MAFB*) were found to be associated with normal facial variation and facial features within the cleft phenotypic spectrum. Although studies above did not evaluate malocclusion directly, their results help prioritize genes for future projects given these gene's roles in craniofacial and dental development.

Unfortunately, similar to other complex traits such as human height (48, 49) and dental tooth eruption (47), the variants identified above have small size effects in facial variation (<10% of trait variability) and explain only a small proportion of the trait's heritability. This missing heritability has been attributed to poor genotyping coverage of low frequency (0.5–5%) and rare variants (<0.5%) and small effect sizes of common variants. Moreover, the presence of gene–gene and/or gene–environment interaction as mechanisms in the trait's etiology, partial or incomplete phe-

notypic approaches and the combined effect of genetic variants in non-coding regulatory regions (50, 51) could also result in missing heritability.

Given the etiological complexity of malocclusion, future studies could anticipate missing heritability problems. These could be minimized utilizing multivariate phenotypes which include both quantitative and dichotomous phenotypes simultaneously to capture more variation information and thus increase analytical power (52). Also, malocclusion studies could benefit from a 'comparison of extremes' approach. This powerful method requires fewer individuals and can reduce genetic heterogeneity as at the extremes of the trait's distribution, the expectation is to find fewer loci with greater effect sizes (49). To date, the comparison of extremes approach has been applied to individuals with long or short mandibular ramus, resulting in significant correlations for the growth hormone receptor (*GHR*) gene in Asian populations (53).

The benefits of identifying genotype–phenotype correlations rest on the potential to develop approaches to improve treatment outcomes. Carriers of risk alleles can be screened for prevention, and risk alleles can be targeted for pharmaceutical interventions that may increase the efficiency of orthopedic appliances in patients with maxillo-mandibular discrepancies. For example, 3D imaging to date has provided direct visualization of size and volume changes in the condyle produced by orthopedic effects (54). This knowledge coupled with the identification of molecular regulators of cell proliferation in the condylar cartilage following mandibular propulsion experiments (55) suggest exciting translational approaches to carry this research into the clinical setting. This includes the possibility of identifying detrimental genetic variation in patients with mandibular size deficiencies, particularly if these are found in molecular mediators of condylar cartilage proliferation. Moreover, studies may confirm that such variants lead to poor orthopedic force responses, and lastly, localized pharmaceutical interventions can be developed to target or compensate for the functional deficiencies caused by such variants.

Conclusions and future directions

Understanding the genetics underlying the dentofacial variation in patients with malocclusion is fundamental to develop preventive strategies and innovative treatment modalities that will benefit individual patients. The technology to acquire comprehensive phenotypic and genetic data to accomplish these discoveries is within our reach, and therefore, it is important for the orthodontic academic centers to establish large consortiums of images and data to speed up such discoveries.

Advances in this field will require researchers to move beyond the discovery of genetic variants conferring susceptibility to malocclusion into translational research to identify those with clinical utility. Moreover, methods to provide access to this information need to be developed almost simultaneously so that actionable genetic information gets rapidly implemented in the clinical setting (56), one of the constant challenges in genomic medicine to date (57).

Clinical relevance

As progress in comprehensive phenotyping and genomic data generation continue, clinically relevant phenotype–genotype correlations will be discovered, providing orthodontists with opportunities to use this information for clinical action. This article summarizes the current knowledge on these topics with the goal of making orthodontists aware of their future potential for making significant impacts in clinical practice.

Acknowledgements: We are thankful for the funding received to complete projects in dentofacial phenotyping and malocclusion. AAOF OFDFA 2008–2011, AAO BRA 2012–2013, the Carver Charitable Trust Grant (#12-4058), the National Center for Advancing Translational Sciences, and the National Institutes of Health (NIH), through Grants CDCR01D000295, 2 UL1 TR000442-06, and T32-DEO14678-09. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

References

- Proff P. Malocclusion, mastication and the gastrointestinal system: a review. *J Orofac Orthop* 2010;71:96–107.
- Claudino D, Traebert J. Malocclusion, dental aesthetic self-perception and quality of life in a 18 to 21 year-old population: a cross section study. *BMC Oral Health* 2013;13:3.
- Baccetti T, Franchi L, McNamara JA Jr, Tollaro I. Early dentofacial features of Class II malocclusion: a longitudinal study from the deciduous through the mixed dentition. *Am J Orthod Dentofacial Orthop* 1997;111:502–9.
- Stahl F, Baccetti T, Franchi L, McNamara JA Jr. Longitudinal growth changes in untreated subjects with Class II Division 1 malocclusion. *Am J Orthod Dentofacial Orthop* 2008;134:125–37.
- Moreno Uribe LM, Howe SC, Kummet C, Vela KC, Dawson DV, Southard TE. Phenotypic diversity in white adults with moderate to severe Class II malocclusion. *Am J Orthod Dentofacial Orthop* 2014;145:305–16.
- Jacobson A, Evans WG, Preston CB, Sadowsky PL. Mandibular prognathism. *Am J Orthod* 1974;66:140–71.
- Miyajima K, McNamara JA Jr, Sana M, Murata S. An estimation of craniofacial growth in the untreated Class III female with anterior cross-bite. *Am J Orthod Dentofacial Orthop* 1997;112:425–34.
- Moreno Uribe LM, Vela KC, Kummet C, Dawson DV, Southard TE. Phenotypic diversity in white adults with moderate to severe Class III malocclusion. *Am J Orthod Dentofacial Orthop* 2013;144:32–42.
- Bui C, King T, Proffit W, Frazier-Bowers S. Phenotypic characterization of Class III patients. *Angle Orthod* 2006;76:564–9.
- Naini FB, Moss JP. Three-dimensional assessment of the relative contribution of genetics and environment to various facial parameters with the twin method. *Am J Orthod Dentofacial Orthop* 2004;126:655–65.
- Townsend G, Hughes T, Luciano M, Bockmann M, Brook A. Genetic and environmental influences on human dental variation: a critical evaluation of studies involving twins. *Arch Oral Biol* 2009c;54(Suppl 1):S45–51.
- Wolff G, Wienker TF, Sander H. On the genetics of mandibular prognathism: analysis of large European noble families. *J Med Genet* 1993;30:112–6.
- El-Gheriani AA, Maher BS, El-Gheriani AS, Sciote JJ, Abu-Shahba FA, Al-Azemi R et al. Segregation analysis of mandibular prognathism in Libya. *J Dent Res* 2003;82:523–7.
- Cruz RM, Krieger H, Ferreira R, Mah J, Hartsfield J Jr, Oliveira S. Major gene and multifactorial inheritance of mandibular prognathism. *Am J Med Genet A* 2008;146A:71–7.
- Otero L, Quintero L, Champsaur D. Inheritance of craniofacial features in Colombian families with class III malocclusion. *Appl Clin Genet* 2010;3:1.
- Nikopensius T, Saag M, Jagomagi T, Annilo T, Kals M, Kivistik PA et al. A missense mutation in DUSP6 is associated with Class III malocclusion. *J Dent Res* 2013;92:893–8.
- Harris JE. Genetic factors in the growth of the head. Inheritance of the craniofacial complex and malocclusion. *Dent Clin North Am* 1975;19:151–60.
- Peck S, Peck L, Kataja M. Class II Division 2 malocclusion: a heritable pattern of small teeth in well-developed jaws. *Angle Orthod* 1998;68:9–20.
- Morrison MW. Relative risk of Class II Division 2 malocclusion in first-degree relatives of probands with Class II Division 2 malocclusion. 2008.
- Attanasio C, Nord AS, Zhu Y, Blow MJ, Li Z, Liberton DK et al. Fine tuning of craniofacial morphology by distant-acting enhancers. *Science* 2013;342:1241006.
- Manolio TA. Study designs to enhance identification of genetic factors in healthy aging. *Nutr Rev* 2007;65(12 Pt 2):S228–33.
- Hu YJ, Berndt SI, Gustafsson S, Ganna A, Genetic Investigation of ANthropometric Traits (GIANT) Consortium, Hirschhorn J et al. Meta-analysis of gene-level associations for rare variants based on single-variant statistics. *Am J Hum Genet* 2013;93:236–48.
- Cruz RM, Hartsfield JK Jr, Falcão-Alencar G, Koller DL, Pereira RW, Mah J et al. Exclusion of Class III malocclusion candidate loci in Brazilian families. *J Dent Res* 2011;90:1202–5.
- Rodrigues JB, Araujo S, Guedes-Pinto H, San Roman F, Viegas C, Bastos E. Analysis of new Matrilin-1 gene variants in a case-control study related to dental malocclusions in *Equus asinus*. *Gene* 2013;522:70–4.
- Li C, Scott DA, Hatch E, Tian X, Mansour SL. Dusp6 (Mkp3) is a negative feedback regulator of FGF-stimulated ERK signaling during mouse development. *Development* 2007;134:167–76.
- Gutierrez SJ, Gomez M, Rey JA, Ochoa M, Gutierrez SM, Prieto JC. Polymorphisms of the noggin gene and mandibular micrognathia: a first approximation. *Acta Odontol Latinoam* 2010;23:13–9.
- Stottmann RW, Anderson RM, Klingensmith J. The BMP antagonists Chordin and Noggin have essential but redundant roles in mouse mandibular outgrowth. *Dev Biol* 2001;240:457–73.
- Pillas D, Hoggart CJ, Evans DM, O'Reilly PF, Sipila K, Lahdesmaki R et al. Genome-wide association study reveals multiple loci associated with primary tooth development during infancy. *PLoS Genet* 2010;6:e1000856.
- Ting TY, Wong RW, Rabie AB. Analysis of genetic polymorphisms in skeletal Class I crowding. *Am J Orthod Dentofacial Orthop* 2011;140:e9–15.
- Yamaguchi T, Park SB, Narita A, Maki K, Inoue I. Genome-wide linkage analysis of mandibular prognathism in Korean and Japanese patients. *J Dent Res* 2005;84:255–9.
- Xue F, Wong R, Rabie AB. Identification of SNP markers on 1p36 and association analysis of EPB41 with mandibular prognathism in a Chinese population. *Arch Oral Biol* 2010;55:867–72.
- Jang JY, Park EK, Ryoo HM, Shin HI, Kim TH, Jang JS et al. Polymorphisms in the Matrilin-1 gene and risk of mandibular prognathism in Koreans. *J Dent Res* 2010;89(11):1203–7.
- Ikuno K, Kajii TS, Oka A, Inoko H, Ishikawa H, Iida J. Microsatellite genome-wide association study for

- mandibular prognathism. *Am J Orthod Dentofacial Orthop* 2014;145:757–62.
34. Frazier-Bowers S, Rincon-Rodriguez R, Zhou J, Alexander K, Lange E. Evidence of linkage in a Hispanic cohort with a Class III dentofacial phenotype. *J Dent Res* 2009;88:56–60.
 35. Li Q, Zhang F, Li X, Chen F. Genome scan for locus involved in mandibular prognathism in pedigrees from China. *PLoS ONE* 2010;5:e12678.
 36. Xue F, Rabie AB, Luo G. Analysis of the association of COL2A1 and IGF-1 with mandibular prognathism in a Chinese population. *Orthod Craniofac Res* 2014;17:144–9.
 37. Tassopoulou-Fishell M, Deeley K, Harvey EM, Sciote J, Vieira AR. Genetic variation in myosin 1H contributes to mandibular prognathism. *Am J Orthod Dentofacial Orthop* 2012;141:51–9.
 38. Li Q, Li X, Zhang F, Chen F. The identification of a novel locus for mandibular prognathism in the Han Chinese population. *J Dent Res* 2011;90:53–7.
 39. Robinson PN. Deep phenotyping for precision medicine. *Hum Mutat* 2012;33:777–80.
 40. Yong R, Ranjitkar S, Townsend G, Smith R, Evans A, Hughes T et al. Dental phenomics: advancing genotype to phenotype correlations in craniofacial research. *Aust Dent J* 2014;59(1 Suppl):34–47.
 41. Hammond P, Suttie M. Large-scale objective phenotyping of 3D facial morphology. *Hum Mutat* 2012;33:817–25.
 42. Miller SF, Weinberg SM, Nidey NL, Defay DK, Marazita ML, Wehby GL et al. Exploratory genotype–phenotype correlations of facial form and asymmetry in unaffected relatives of children with non-syndromic cleft lip and/or palate. *J Anat* 2014;224:688–709.
 43. Boehringer S, van der Lijn F, Liu F, Gunther M, Sinigerova S, Nowak S et al. Genetic determination of human facial morphology: links between cleft-lips and normal variation. *Eur J Hum Genet* 2011;19:1192–7.
 44. Peng S, Tan J, Hu S, Zhou H, Guo J, Jin L et al. Detecting genetic association of common human facial morphological variation using high density 3D image registration. *PLoS Comput Biol* 2013;9:e1003375.
 45. Paternoster L, Zhurov AI, Toma AM, Kemp JP, St Pourcain B, Timpson NJ et al. Genome-wide association study of three-dimensional facial morphology identifies a variant in PAX3 associated with nasion position. *Am J Hum Genet* 2012;90:478–85.
 46. Liu F, van der Lijn F, Schurmann C, Zhu G, Chakravarty MM, Hysi PG et al. A genome-wide association study identifies five loci influencing facial morphology in Europeans. *PLoS Genet* 2012;8:e1002932.
 47. Fatemifar G, Hoggart C, Paternoster L, Kemp JP, Prokopenko I, Horikoshi M et al. Genome-wide association study of primary tooth eruption identifies pleiotropic loci associated with height and craniofacial distances. *Hum Mol Genet* 2013;22(18):3807–17.
 48. Lango Allen H, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F et al. Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature* 2010;467:832–8.
 49. Berndt SI, Gustafsson S, Magi R, Ganna A, Wheeler E, Feitosa MF et al. Genome-wide meta-analysis identifies 11 new loci for anthropometric traits and provides insights into genetic architecture. *Nat Genet* 2013;45:501–12.
 50. Zuk O, Schaffner SF, Samocha K, Do R, Hechter E, Kathiresan S et al. Searching for missing heritability: designing rare variant association studies. *Proc Natl Acad Sci U S A* 2014;111:E455–64.
 51. Corradin O, Saiakhova A, Akhtar-Zaidi B, Myeroff L, Willis J, Cowper-Salari R et al. Combinatorial effects of multiple enhancer variants in linkage disequilibrium dictate levels of gene expression to confer susceptibility to common traits. *Genome Res* 2014;24:1–13.
 52. Melton PE, Pankratz N. Joint analyses of disease and correlated quantitative phenotypes using next-generation sequencing data. *Genet Epidemiol* 2011;35(Suppl 1):S67–73.
 53. Tomoyasu Y, Yamaguchi T, Tajima A, Nakajima T, Inoue I, Maki K. Further evidence for an association between mandibular height and the growth hormone receptor gene in a Japanese population. *Am J Orthod Dentofacial Orthop* 2009;136:536–41.
 54. Yildirim E, Karacay S, Erkan M. Condylar response to functional therapy with Twin-Block as shown by cone-beam computed tomography. *Angle Orthod* 2014;84:1018–25.
 55. Patil A, Sable R, Kothari R. Genetic expression of MMP-Matrix-metallproteinases (MMP-1 and MMP-13) as a function of anterior mandibular repositioning appliance on the growth of mandibular condylar cartilage with and without administration of Insulin like growth factor (IGF-1) and Transforming growth factor-B (TGF-beta). *Angle Orthod* 2012;82:1053–9.
 56. Ramos EM, Din-Lovinescu C, Berg JS, Brooks LD, Duncanson A, Dunn M et al. Characterizing genetic variants for clinical action. *Am J Med Genet C Semin Med Genet* 2014;166:93–104.
 57. Manolio TA, Green ED. Leading the way to genomic medicine. *Am J Med Genet C Semin Med Genet* 2014;166:1–7.

Copyright of Orthodontics & Craniofacial Research is the property of Wiley-Blackwell and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.