

Guideline on Adolescent Oral Health Care

Originating Committee
Clinical Affairs Committee

Review Council
Council on Clinical Affairs and Committee on the Adolescent

Adopted
1986

Revised
1999, 2003, 2005

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that the adolescent patient has unique needs. This guideline addresses these unique needs and proposes general recommendations for their management.

Methods

This guideline is based on a review of the current dental and medical literature related to adolescent oral health. A MEDLINE search was conducted using the term "adolescent" combined with "dental", "gingivitis", "oral piercing", "sealants", "oral health", "caries", "tobacco use", "dental trauma", "orofacial trauma", "periodontal", "dental esthetics", "smokeless tobacco", "nutrition", and "diet".

Background

There is no standard definition of "adolescent".¹ Adolescents are defined very broadly as youths between the ages of 10 to 18. Using this definition, there were approximately 39.9 million adolescents in the United States in 2003, according to the US Census Bureau.² The adolescent patient is recognized as having distinctive needs^{3,4} due to: (1) a potentially high caries rate; (2) increased risk for traumatic injury and periodontal disease; (3) a tendency for poor nutritional habits; (4) an increased esthetic desire and awareness; (5) complexity of combined orthodontic and restorative care (eg, congenitally missing teeth); (6) dental phobia; (7) initiation of tobacco use; (8) pregnancy; (9) eating disorders; and (10) unique social and psychological needs.⁵⁻⁸

Treatment of the adolescent patient can be multifaceted and complex. An accurate, comprehensive, and up-to-date medical history is necessary for correct diagnosis and effective treatment planning. Familiarity with the patient's medical history is essential to decreasing the risk of aggravating a medical condition while rendering dental care. If the parent is unable to provide adequate details regarding a patient's medical history, consultation with the medical health care provider may be indicated. The practitioner also may need to obtain additional information confidentially from an adolescent patient.

Recommendations

This guideline addresses some of the special needs within the adolescent population and proposes general recommendations for their management.

Caries

Adolescence marks a period of significant caries activity for many individuals. Current research suggests that the overall caries rate is declining, yet remains highest during adolescence.⁹ These carious lesions often are confined to developmental pits and fissures.¹⁰

Immature permanent tooth enamel,¹¹ a total increase in susceptible tooth surfaces, and environmental factors such as diet, independence to seek care or avoid it, a low priority for oral hygiene, and additional social factors also may contribute to the upward slope of caries in adolescence.^{1,12-14} It is important for the dental provider to emphasize the positive effects that fluoridation, routine professional care, patient education, and personal hygiene can have in counteracting the changing pattern of caries in the adolescent population.^{5,6,15}

Management of Caries

Primary prevention

Fluoride: Fluoridation has proven to be the most economical and effective caries prevention measure. The adolescent can benefit from fluoride throughout the teenage years and into early adulthood. Although the systemic benefit of fluoride incorporation into developing enamel is not considered necessary past 16 years of age, the topical benefits of remineralization and antimicrobial activity still can be obtained through water fluoridation, professionally applied and prescribed compounds, and fluoridated dentifrices.¹⁶

Recommendations: The adolescent should receive maximum fluoride benefit:

1. Systemic fluoride intake via optimal fluoridation of drinking water or professionally prescribed supplements is recommended to 16 years of age or the eruption of the second permanent molars, whichever comes first.
2. Fluoridated dentifrice is recommended to provide continuing topical benefits through adolescence.

3. Professionally applied fluoride treatments should be based on the individual patient's caries-risk assessment, as determined by the patient's dental provider.
4. Topical fluoride supplementation via home-applied compounds should be a professional recommendation when indicated by an individual's caries pattern or caries-risk status.
5. The criteria for determination of need and the methods of delivery should be those currently recommended by the American Dental Association and the AAPD.¹⁷

Oral hygiene: Adolescence can be a time of heightened caries activity and periodontal disease due to an increased intake of cariogenic substances and inattention to oral hygiene procedures.^{1,18} Tooth-brushing with a fluoridated dentifrice and flossing can provide an anticaries benefit through plaque removal from tooth surfaces and the topical effect of the fluoride.

Recommendations:

1. Adolescents should be educated and motivated to maintain personal oral hygiene through daily plaque removal, including flossing, with the frequency and pattern based on the individual's disease pattern and oral hygiene needs.
2. Professional removal of plaque and calculus is recommended highly for the adolescent, with the frequency of such intervention based on the individual's assessed risk for caries/periodontal disease, as determined by the patient's dental provider.¹⁹

Diet management: The role of carbohydrates in caries initiation is unequivocal. Adolescents are exposed to and consume high quantities of refined carbohydrates and acid-containing beverages.^{12,13,20} The adolescent can benefit from diet analysis and modification.

Recommendations: Diet analysis, along with professionally determined recommendations for maximal general and dental health, should be part of an adolescent's dental health management. A diet analysis and management should consider:

1. dental disease patterns;
2. overall nutrient and energy needs;
3. psychosocial aspects of adolescent nutrition;
4. dietary carbohydrate intake and frequency;
5. intake and frequency of acid-containing beverages;
6. wellness considerations.

Sealants: Sealant placement is an effective caries-preventive technique that should be considered on an individual basis. Sealants have been recommended on any adolescent tooth, primary or permanent, that is judged to be at risk for pit and fissure caries.^{6,13,21-24} Caries risk may increase due to changes in patient habits, oral microflora, or physical condition, and unsealed teeth subsequently might benefit from sealant applications.²⁴

Recommendations: Adolescents at risk for caries should have sealants placed. An individual's caries risk may change

over time; periodic reassessment for sealant need is indicated throughout adolescence.²⁴

Secondary prevention

Professional preventive care: Professional preventive dental care, on a routine basis, may prevent oral disease or disclose existing disease in its early stages. The adolescent patient whose oral health has not been monitored routinely by a dentist may have advanced caries, periodontal disease, or other oral involvement urgently in need of professional evaluation and extensive treatment.

Recommendations:

1. Timing of periodic oral examinations should take into consideration the individual's needs and risk indicators to determine the most cost-effective, disease-preventive benefit to the adolescent.
2. Initial and periodic radiographic evaluation should be a part of a clinical evaluation. The type, number, and frequency of radiographs should be determined only after an oral examination and history taking. Previously exposed radiographs should be available, whenever possible, for comparison. Currently accepted guidelines for radiographic exposures (ie, appropriate films based upon medical history, caries risk, history of periodontal disease, and growth and development assessments) should be followed.²⁵

Restorative dentistry: In cases where remineralization of noncavitated, demineralized tooth surfaces is not successful, as demonstrated by progression of carious lesions, dental restorations are necessary. Preservation of tooth structure, esthetics, and each individual patient's needs must be considered when selecting a restorative material.²⁶ Molars with extensive caries or malformed, hypoplastic enamel—for which traditional amalgam or composite resin restorations are not feasible—may require full coverage restorations.^{24,27}

Recommendations: Each adolescent patient and restoration must be evaluated on an individual basis. Preservation of noncarious tooth structure is desirable. Referral to an appropriately trained and/or experienced dentist should be considered when treatment needs are beyond the treating dentist's ability or interest.

Periodontal diseases

Adolescence can be a critical period in the human being's periodontal status. Epidemiologic and immunologic data suggests that irreversible tissue damage from periodontal disease begins in late adolescence and early adulthood.⁸ Pubertal changes characteristically affect the periodontium of the young adolescent, with an increase in inflammation which is, in most cases, manageable through oral hygiene and regular professional care.²⁸

Acute conditions: The adolescent may be subjected to acute conditions such as acute necrotizing ulcerative gingivitis, periodontitis, and traumatic injuries, which can require immediate and occasional long-term management. In most

cases, early diagnosis, treatment, and appropriate management can prevent irreversible damage.²⁹⁻³¹

Recommendations: Acute intraoral infection involving the periodontium and oral mucosa requires immediate treatment. Therapeutic management should be based on currently accepted techniques of periodontal therapy. Traumatic injuries to the teeth and periodontium always require dental evaluation and treatment. Referral to an appropriately trained and/or experienced dentist should be considered when the treatment needs are beyond the treating dentist's ability or interest.

Chronic conditions: Chronic conditions affecting the adolescent include, but are not limited to, gingivitis, puberty gingivitis, hyperplastic gingivitis related to orthodontic therapy, gingival recession that may or may not be related to orthodontic therapy, drug-related gingivitis, pregnancy gingivitis, localized juvenile periodontitis, and periodontitis.^{21,28,29,32} Personal oral hygiene and regular professional intervention can minimize occurrence of these conditions and prevent irreversible damage.

Recommendations: The adolescent will benefit from an individualized preventive dental health program, which includes the following items aimed specifically at periodontal health:

1. Patient education emphasizing the etiology, characteristics, and prevention of periodontal diseases, as well as self-hygiene skills.
2. A personal, age-appropriate oral hygiene program including plaque removal, oral health self-assessment, and diet. Sulcular brushing and flossing should be included in plaque removal, and frequent follow-up to determine adequacy of plaque removal and improvement of gingival health should be considered.
3. Regular professional intervention, the frequency of which should be based on individual needs and should include evaluation of personal oral hygiene success, periodontal status, and potential complicating factors, such as medical conditions, malocclusion, or handicapping conditions. Periodontal probing, periodontal charting, and radiographic periodontal diagnosis should be a consideration when caring for the adolescent. The extent and nature of the periodontal evaluation should be determined professionally on an individual basis. Those patients with progressive periodontal disease should be referred to an appropriately trained and/or experienced dentist for evaluation and treatment.
4. Appropriate evaluation for procedures to facilitate orthodontic treatment including, but not limited to, tooth exposure, frenectomy, fiberotomy, gingival augmentation, and implant placement.³³

Occlusal considerations

Malocclusion can be a significant treatment need in the adolescent population as both environmental and genetic factors come into play. Although the genetic basis of much

malocclusion makes it unpreventable, numerous methods exist to treat the occlusal disharmonies, temporomandibular joint dysfunction, periodontal disease, and disfigurement which may be associated with malocclusion. Within the area of occlusal problems are several tooth/jaw-related discrepancies, which can affect the adolescent. Third molar malposition and temporomandibular disorders require special attention to avoid long-term problems. Congenitally missing teeth present complex problems for the adolescent and often require combined orthodontic and restorative care for satisfactory resolution.

Malocclusion: Any tooth/jaw positional problems that present significant esthetic, functional, physiologic, or emotional dysfunction are potential difficulties for the adolescent. These can include single or multiple tooth malpositions, tooth/jaw size discrepancies, and craniofacial disfigurements.

Recommendations: Any malposition of teeth, malrelationship of teeth to jaws, tooth/jaw size discrepancy, bimaxillary malrelationship, or craniofacial malformations or disfigurement that presents functional, esthetic, physiologic, or emotional problems to the adolescent should be evaluated by the appropriately trained dentist or professional team. Treatment of malocclusion by an appropriately trained and/or experienced dentist should be based on professional diagnosis, available treatment options, patient motivation and readiness, and other factors to maximize progress.

Third molars: Third molars can present acute and chronic problems for the adolescent. Impaction or malposition leading to such problems as pericoronitis, caries, cysts, or periodontal problems merits evaluation for removal.^{32,34} The role of the third molar as a functional tooth should also be considered. Although prophylactic removal of all impacted or unerupted disease-free third molars is not indicated, consideration should be given to removal by the third decade when there is a high probability of disease or pathology and/or the risks associated with early removal are less than the risks of later removal.

Recommendations: Evaluation of third molars, including radiographic diagnostic aids, should be an integral part of the dental examination of the adolescent.²⁵ For diagnostic and extraction criteria, refer to pages 158-164 for the AAPD's Clinical Guideline on Pediatric Oral Surgery.³⁵ Treatment of third molars that are potential or active problems should be performed by an appropriately trained and/or experienced dentist.

Temporomandibular joint problems: Disorders of the temporomandibular joint can occur at any age, but adolescence may provide the stimulus to trigger problems.³⁶⁻³⁹

Recommendations: Evaluation of the temporomandibular joint and related structures should be a part of the examination of the adolescent. Abnormalities should be managed by an appropriately trained and/or experienced dentist following accepted clinical procedures.^{40,41}

Congenitally missing teeth: The impact of a congenitally missing permanent tooth on the developing dentition can be significant.³ When treating adolescent patients with congenitally missing teeth, many factors must be taken into consideration including, but not limited to, esthetics, patient age, and growth potential, as well as periodontal and oral surgical needs.⁴²⁻⁴⁴

Recommendations: Evaluation of congenitally missing permanent teeth should include both immediate and long-term management. Management should be by an appropriately trained and/or experienced dentist, and a team approach may be indicated.⁴⁵

Ectopic eruption: Abnormal eruption patterns of the adolescent's permanent teeth can contribute to root resorption, bone loss, gingival defects, space loss, and esthetic concerns. Early diagnosis and treatment of ectopically erupting teeth can result in a healthier and more esthetic dentition. Prevention and treatment may include extraction of deciduous teeth, surgical intervention, and/or endodontic, orthodontic, periodontal, and/or restorative care.⁴⁶⁻⁵⁰

Recommendations: The dentist should be proactive in diagnosing and treating ectopic eruption in the young adolescent. Early diagnosis, including appropriate radiographic examination²⁵ of ectopic eruption, is important. An appropriately trained and/or experienced dentist should manage treatment, and a team approach may be necessary.⁴⁵

Traumatic injuries

The most common injuries to permanent teeth occur secondary to falls, followed by traffic accidents, violence, and sports.⁵¹⁻⁵⁴ All sporting activities have an associated risk of orofacial injuries due to falls, collisions, and contact with hard surfaces.⁵⁵ The administrators of youth, high school, and college organized sports have demonstrated that dental and facial injuries can be reduced significantly by introducing mandatory protective equipment such as face guards and mouthguards. Additionally, youths participating in leisure activities such as skateboarding, rollerskating, and bicycling also benefit from appropriate protective equipment.^{56,57}

Recommendations: Dentists should introduce a comprehensive trauma prevention program to help reduce the incidence of traumatic injury to the adolescent dentition. This prevention plan should consider assessment of the patient's sport or activity, including level and frequency of activity.⁵⁸ Once this information is acquired, recommendation and fabrication of an age-appropriate, sport-specific, and properly-fitted mouthguard/faceguard can be initiated.⁵⁸ Players must be warned about altering the protective equipment that will disrupt the fit of the appliance. In addition, players and parents must be informed that injury may occur, even with properly fitted protective equipment.⁵⁸

Additional considerations in oral/dental management of the adolescent

The adolescent can present particular psychosocial characteristics that impact the health status of the oral cavity, care

seeking, and compliance. The self-concept development process, emergence of independence, and the influence of peers are just a few of the psychodynamic factors impacting dental health during this period.^{1,5,7,16}

Discolored or stained teeth: Desire to improve esthetics of the dentition by tooth whitening and removal of stained areas or defects can be a concern of the adolescent. Indications for the appropriate use of tooth-whitening methods and products are dependent upon correct diagnosis.⁵⁹

The dentist must determine the appropriate mode of treatment. Use of bleaching agents, microabrasion, placement of an esthetic restoration, or a combination of treatments all can be considered.⁶⁰

Recommendations: For the adolescent patient, judicious use of bleaching can be considered part of a comprehensive, sequenced treatment plan that takes into consideration the patient's dental developmental stage, oral hygiene, and caries status. A dentist should monitor the bleaching process, ensuring the least invasive, most effective treatment method. Dental professionals also should consider possible side effects when contemplating dental bleaching for adolescent patients.^{61,62}

Tobacco use: Significant oral, dental, and systemic health consequences and death are associated with all forms of tobacco use. Smoking and other tobacco use almost always are initiated and established in adolescence.⁶³⁻⁶⁸

Recommendations: Education of the adolescent patient on the oral and systemic consequences of tobacco use should be part of each patient's oral health education. For those adolescent patients who use tobacco products, the practitioner should provide or refer the patient to appropriate educational and counseling services.⁶⁹⁻⁷¹ When associated pathology is present, treatment should be managed by an appropriately trained and/or experienced health care provider.

Positive youth development: Treatment and management of adolescent oral health that takes into account the adolescent's psychological and social needs can be approached through the framework of positive youth development (PYD).⁷² The approach goes beyond traditional prevention, intervention, and treatment of risky behaviors and problems and suggests that a strong interpersonal relationship between the adolescent patient and the pediatric dentist can be influential in improving adolescent oral health and transitioning patients to adult care. In the office, dental professionals have a unique opportunity to serve as positive role models.

Recommendations: PYD should be recognized as containing a number of key elements relevant to care patients of this age:

1. providing youth with safe and supportive environments;
2. fostering relationships between young people and caring adults who can mentor and guide them;

3. promoting healthy lifestyles and teaching positive patterns of social interaction;
4. providing a safety net in times of need.⁷²

Integrating PYD into clinical practice can be attained through continuing education on adolescent development issues, as well as partnerships with community-based organizations and schools. The dentist can be a part of the myriad of adolescent support and services.

Psychosocial and other considerations: Behavioral considerations when treating an adolescent may include anxiety, phobia, or intellectual dysfunction.¹ These special needs should be managed by appropriately trained dentists. Referral to nondental professionals or a team approach may be indicated.

Additional examples of oral problems associated with adolescent behaviors include, but are not limited to:

1. oral manifestations of venereal diseases;
2. effects of oral contraceptives or antibiotics on periodontal structures;
3. perimyolysis in bulimia⁷³;
4. traumatic injury to teeth and oral structures in athletic or other activities (short- and long-term management)^{57,74-76};
5. intraoral and perioral piercing with possible local and systemic effects.^{77,78}

The impact of psychosocial factors relating to oral health must include consideration of the following:

1. changes in dietary habits (eg, fads, freedom to snack, increased energy needs, access to carbohydrates);
2. use and abuse of drugs;
3. motivation for maintenance of good oral hygiene;
4. potential for traumatic injury;
5. adolescent as responsible for care;
6. lack of knowledge about periodontal disease.

Physiologic changes also can account for significant oral problems in the adolescent. These include:

1. loss of remaining primary teeth;
2. eruption of remaining permanent teeth;
3. gingival maturity;
4. facial growth;
5. hormonal changes.

Recommendations:

1. Oral health care of the adolescent should be provided by a dentist who has appropriate training in managing the patient's specific needs. The primary care dentist should consider referral to a specialist for treatment of particular problems outside his or her expertise. This may include both dental and nondental problems.
2. Attention should be given to the particular psychosocial aspects of adolescent dental care. Other issues such as consent, confidentiality, and compliance should be addressed in the care of these patients.^{79,80}
3. A complete oral health care program for the adolescent requires an educational component that addresses the particular concerns and needs of the adolescent patient and focuses on:

- a. specific behaviorally and physiologically induced oral manifestations in this age group;
- b. shared responsibility for care and health by the adolescent and provider;
- c. consequences of adolescent behavior on oral health.

Transitioning to adult care: As adolescent patients approach the age of majority, it is important to educate the patient and parent on the value of transitioning to a dentist who is knowledgeable in adult oral health care. The adult's oral health needs may go beyond the scope of the pediatric dentist's training. The transitioning adolescent should continue professional oral health care in an environment sensitive to his/her individual needs. Many adolescent patients independently will choose the time to seek care from a general dentist and may elect to seek treatment from a parent's primary care provider. In some instances, however, the treating pediatric dentist will be required to suggest transfer to adult care.

Pediatric dentists are concerned about decreased access to oral health care for persons with special health care needs (SHCN)⁸¹ as they transition beyond the age of majority. Pediatric hospitals, by imposing age restrictions, can create a barrier to care for these patients. Transitioning to a dentist who is knowledgeable and comfortable with adult oral health care needs often is difficult due to a lack of trained providers willing to accept the responsibility of caring for SHCN patients.

Recommendations: At a time agreed upon by the patient, parent, and pediatric dentist, the patient should be transitioned to a dentist knowledgeable and comfortable with managing that patient's specific oral care needs. For the SHCN patient, in cases where it is not possible or desired to transition to another practitioner, the dental home can remain with the pediatric dentist and appropriate referrals for specialized dental care should be recommended when needed.⁸²

References

1. American Psychological Association. *Developing Adolescents: A Reference for Professionals*. Washington, DC: American Psychological Association; 2002.
2. US Census Bureau. (2001) Census 2000 PHC-T-9. Population by age, sex, race for the United States. Census 2003 Summary File I. Available at: <http://factfinder.census.gov>. Accessed February 7, 2005.
3. Pinkham JR, Casamassimo PS, Fields HW, McTigue DJ, Nowak AJ. Adolescent. In: *Pediatric Dentistry: Infancy Through Adolescence*. 4th ed. Philadelphia, Pa: WB Saunders Co; 2005:649-718.
4. National Institutes of Health. Consensus development conference statement: Diagnosis and management of dental caries throughout life, March 26-28, 2001. J Am Dent Assoc 2001;132:153-1161.
5. Macgregor ID, Regis D, Balding J. Self-concept and dental health behaviors in adolescents. J Clin Periodontol 1997;24:335-339.

6. Yu SM, Bellamy HA, Schwalberg RH, Drum MA. Factors associated with use of preventive dental and health services among US adolescents. *J Adolesc Health* 2001; 29:395-405.
7. American Academy of Pediatric Dentistry. Policy on prevention of sports-related orofacial injuries. *Pediatr Dent* 2004;26(suppl):44.
8. US Dept of Health and Human Services. *Oral Health In America: A Report of the Surgeon General—Executive Summary*. Rockville, Md: US Dept of Health and Human Services, National Institute of Dental and Craniofacial Research, National Institutes of Health; 2000.
9. Kaste LM, Selwitz RH, Oldakowski JA, Brunelle JA, Winn DM, Brown LJ. Coronal caries in the primary and permanent dentition of children and adolescents 1-17 years of age: United States, 1988-1991. *J Dent Res* 1996;75(special issue):631-641.
10. Burt BA. Prevention policies in light of the changed distribution of dental caries. *Acta Odontol Scand* 1998;56:179-186.
11. Kirkham J, Robinson C, Strong M, Shore RC. Effects of frequency of acid exposure on demineralization/remineralization behavior of human enamel in vitro. *Caries Res* 1994;28:9-13.
12. Howze KA. *Health for Teens in Care: A Judge's Guide 2002*. Washington, DC: American Bar Association; 2002.
13. Majewski RF. Dental caries in adolescents associated with caffeinated carbonated beverages. *Pediatr Dent* 2001;23:198-203.
14. Marshall TA, Levy SM, Broffitt B, et al. Dental caries and beverage consumption in young children. *Pediatrics* 2003;112:e184-e191.
15. Irwin CE, Millstein SG. Biophysical correlates of risk-taking behaviors during adolescence. *J Adolesc Health Care* 1986;7:825-965.
16. CDC. Recommendations for using fluoride to prevent and control dental caries in the United States. *MMWR* 2001;50:1-42.
17. American Academy of Pediatric Dentistry. Clinical guideline on fluoride therapy. *Pediatr Dent* 2004;26(suppl): 87-88.
18. Macgregor ID, Balding J, Regis D. Tooth-brushing schedule, motivation, and lifestyle behaviors in 7,770 young adolescents. *Community Dent Health* 1996;13: 232-237.
19. American Academy of Pediatric Dentistry. Clinical guideline on periodicity of examination, preventive dental services, anticipatory guidance, and oral treatment for children. *Pediatr Dent* 2004;26(suppl):81-83.
20. Freeman R, Sheiham A. Understanding decision-making process for sugar consumption in adolescents. *Community Dent Oral Epidemiol* 1997;25:228-232.
21. Feigal RJ. The use of pit and fissure sealants. *Pediatr Dent* 2002;24:415-422.
22. Ahovuo-Saloranta A, Hiiri A, Nordblad A, Worthington H, Makela M. Pit and fissure sealants for preventing dental decay in the permanent teeth of children and adolescents [review]. *Cochrane Database Syst Rev* 2004;3:CD001830.
23. Macek MD, Beltran-Aguilar ED, Lockwood SA, Malvitz DM. Updated comparison of the caries susceptibility of various morphological types of permanent teeth. *J Public Health Dent* 2003;63:174-182.
24. American Academy of Pediatric Dentistry. Clinical guideline on pediatric restorative dentistry. *Pediatr Dent* 2004;26(suppl):106-114.
25. American Academy of Pediatric Dentistry. Clinical guideline on prescribing dental radiographs for infants, children, adolescents, and persons with special health care needs. *Pediatr Dent* 2005;27(suppl):185-186.
26. Donly K. Pediatric Restorative Dentistry Consensus Conference. *Pediatr Dent* 2002;24:374.
27. Croll TP, Castaldi CR. The preformed stainless steel crown for restoration of permanent posterior teeth in special cases. *J Am Dent Assoc* 1978;97:644-649.
28. American Academy of Periodontology. Oral health info for the public: Adolescents and oral health care. Available at: <http://www.perio.org/consumerchildren.htm>. Accessed February 7, 2005.
29. Modeer T, Wondimu B. Periodontal diseases in children and adolescents. *Dent Clin North Am* 2000;44: 633-658.
30. Grossi SG, Zambon JJ, Ho AW, et al. Assessment of risk for periodontal disease. I. Risk indicators for attachment loss. *J Periodontol* 1994;65:260-267.
31. Grossi SG, Genco RJ, Machtei EE, et al. Assessment of risk for periodontal disease. II. Risk indicators for alveolar bone loss. *J Periodontol* 1995;66:23-29.
32. Litonjua LS. Pericoronitis, deep fascial space infections, and the impacted third molar. *J Philipp Dent Assoc* 1996;47:43-47.
33. American Academy of Periodontology. Periodontal therapy. *J Periodontol* 2001;72:1624-1628.
34. Waller JH, Malden N. Rapid cystic involvement of a lower third molar. *Dent Update* 1999;26:166-167.
35. American Academy of Pediatric Dentistry. Clinical guideline on pediatric oral surgery. *Pediatr Dent* 2005;27(suppl):158-164.
36. American Academy of Pediatric Dentistry. Treatment of temporomandibular disorders in children: Summary statements and recommendations. *J Am Dent Assoc* 1990;120:265-269.
37. Riolo ML, ten Have TR, Brandt D. Clinical validity of the relationship between TMJ signs and symptoms in children and youth. *J Dent Child* 1988;55:110-113.
38. Alamoudi N, Farsi N, Salako N, Feteih R. Temporomandibular disorders among school children. *J Clin Pediatr Dent* 1998;22:323-329.
39. Nydell A, Helkimo M, Koch G. Craniomandibular disorders in children: A critical review of the literature. *Swed Dent J* 1994;18:191-205.

40. National Institutes of Health. *Management of Temporomandibular Disorders*. Bethesda, Md; NIH Techno Assess Statement; 1996:1-31. Available at: http://consensus.nih.gov/ta/018/018_statement.htm. Accessed February 7, 2005.
41. Skeppar J, Nilner M. Treatment of craniomandibular disorders in children and young adults. *J Orofac Pain* 1993;7:362-369.
42. Garg AK. Treatment of congenitally missing maxillary incisors: Orthodontics, bone grafts, and osseointegrated implants. *Dent Implantol Update* 2002;13:9-14.
43. Wexler G. Missing upper lateral incisors: Orthodontic considerations in young patients. *Ann R Australas Coll Dent Surg* 2002;15:136-140.
44. Richardson G, Russell KA. Congenitally missing maxillary incisors and orthodontic treatment considerations for the single tooth implant. *J Can Dent Assoc* 2001;67:25-28.
45. American Academy of Pediatric Dentistry. Clinical guideline on management of the developing dentition and occlusion in pediatric dentistry. *Pediatr Dent* 2005;27(suppl):143-155.
46. Chaushu S, Sharabi S, Becker A. Dental morphologic characteristics of normal versus delayed developing dentitions with palatally displaced canines. *Am J Orthod Dentofacial Orthop* 2002;121:399-346.
47. Kojima R, Taguchi Y, Kabayashi H, Noda T. External root resorption of the maxillary permanent incisors caused by ectopically erupting canines. *J Clin Pediatr Dent* 2002;26:193-197.
48. Ericson S, Kurol PJ. Resorption of incisors after ectopic eruption of maxillary canines. *Angle Orthod* 2000;70:415-423.
49. Shapira Y, Borell G, Kuftinec MM. Bringing impacted mandibular second premolars into occlusion. *J Am Dent Assoc* 1996;127:1075-1078.
50. Kurol J. Early treatment of tooth eruption disturbances. *Am J Orthod Dentofacial Orthop* 2002;121:588-591.
51. Rocha MJdC, Cardoso M. Traumatized permanent teeth in Brazilian children assisted at the Federal University of Santa Catarina, Brazil. *Dent Traumatol* 2001;17:245-249.
52. deFranca Caldas A Jr, Burgos MEA. A retrospective study of traumatic dental injuries in a Brazilian dental trauma clinic. *Dent Traumatol* 2001;17:250-253.
53. Skaare AB, Jacobsen I. Dental injuries in Norwegians aged 7-18 years. *Dent Traumatol* 2003;19:67-71.
54. Tapias Ma, Jimenez-Garcia R, Lamas F, Bil AA. Prevalence of traumatic crown fractures to permanent incisors in a childhood population: Mostoles, Spain. *Dent Traumatol* 2003;19:119-122.
55. Gassner R, Bosch R, Tuli T, Emschhoff R. Prevalence of dental trauma in 6,000 patients with facial injuries: Implications for prevention. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1999;87:27-33.
56. Tesini DA, Soporowski NJ. Epidemiology of orofacial sports-related injuries. *Dent Clin North Am* 2000;44:1-18.
57. Ranalli DN. Prevention of sport-related dental traumatic injuries. *Dent Clin North Am* 2000;44:19-33.
58. Ranalli DN. A sports dentistry trauma control plan for children and adolescents. *J Southeast Soc Pediatr Dent* 2002;8:8-9.
59. Sarrett DC. Tooth whitening today. *J Am Dent Assoc* 2002;133:1535-1538.
60. Donly KJ. The adolescent patient: Special whitening challenges. *Compend Contin Educ Dent* 2003;24:390-396.
61. Li Y. Tooth bleaching using peroxide containing agents: Current status of safety issues. *Compendium* 1998;19:783-796.
62. American Academy of Pediatric Dentistry. Policy on dental bleaching for child and adolescent patients. *Pediatr Dent*. 2004;26(suppl):45-47.
63. US Dept of Health and Human Services. *Preventing Tobacco Use Among Young People: Report of the Surgeon General*. Atlanta, Ga: US Dept of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 1994.
64. CDC. Tobacco use among high school students—United States, 1997. *MMWR* 1998;47:229-233.
65. Tomar SL, Winn DM, Swango PA, Giovino GA, Kleinman DV. Oral mucosal smokeless tobacco lesions among adolescents in the United States. *J Dent Res* 1997;76:1277-1286.
66. Audrain-McGovern J, Rodriguez D, Tercyak KP, Cuevas J, Rodgers K, Patterson F. Identifying and characterizing adolescent smoking trajectories. *Cancer Epidemiol Biomarkers Prev* 2004;13:2023-2034.
67. Zullig KJ, Valois RF, Huebner ES, Drane JW. Evaluating the performance of the Centers for Disease Control and Prevention core Health-Related quality of life scale with adolescents. *Public Health Rep* 2004;119:577-584.
68. Johnson CC, Myers L, Webber LS, Boris NW. Profiles of the adolescent smoker: Models of tobacco use among 9th grade high school students; Acadiana Coalition of Teens against Tobacco (ACTT). *Prev Med* 2004;39: 551-558.
69. American Dental Association. Summary of policy and recommendations regarding tobacco: 1964-present. ADA Resolution 1H-1992. In: *ADA Transactions 1992*. Chicago, Ill: ADA; 1993:598.
70. American Cancer Society, National Cancer Institute, National Institutes of Health. *How to Help Your Patients Stop Using Tobacco: A National Cancer Institute Manual for the Oral Health Team*. Bethesda, Md: National Institutes of Health, US Dept of Health and Human Services, Public Health Service; 1998. NIH publication No. 98-3191.
71. American Academy of Pediatric Dentistry. Policy on Tobacco Use. *Pediatr Dent* 2004;27(suppl):42-43.

72. US Department of Health and Human Services Administration for Children and Families. Toward a Blueprint for Youth: Making Positive Youth Development a National Priority. Rockville, Md: US Department of Health and Human Services; 2002. Available at: <http://www.acf.dhhs.gov/programs/fysb/youthinfo/blueprint2.htm>. Accessed February 7, 2005.
73. Christenson GJ. Oral care for patients with bulimia. *J Am Dent Assoc* 2002;133:1689-1691.
74. Cortes MIS, Marcenes W, Sheiham A. Impact of traumatic injuries to the permanent teeth on the oral health-related quality of life in 12- to 14-year-old children. *Community Dent Oral Epidemiol* 2002;30:193-198.
75. Gassner R, Tuli T, Hachl O, Moreira R, Ulmer H. Craniomaxillofacial trauma in children: A review of 3,385 cases with 6,060 injuries in 10 years. *J Oral Maxillofac Surg* 2004;62: 399-407.
76. Barnett F. Prevention of sports-related dental trauma: The role of mouthguards. *Pract Proced Aesthet Dent* 2003;15:391-394.
77. American Dental Association. Statement on intraoral/perioral piercing. Available at: <http://www.ada.org/prof/recoveries/positions/statements/piercings.asp.html>. Accessed February 7, 2005.
78. Boardman R Smith RA. Dental implications of oral piercing. *J Calif Dent Assoc* 1997;25:200-207.
79. American Academy of Pediatric Dentistry. Clinical guideline on record-keeping. *Pediatr Dent* 2004;26(suppl): 134-139.
80. American Academy of Pediatric Dentistry. Clinical guideline on informed consent. *Pediatr Dent* 2005; 27(suppl):182-183.
81. American Academy of Pediatric Dentistry. Definition of persons with special health care needs. *Pediatr Dent* 2004;26(suppl):15.
82. American Academy of Pediatric Dentistry. Clinical guideline on management of persons with special health care needs. *Pediatr Dent* 2004;26(suppl):77-80.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.