Policy on Minimizing Occupational Health Hazards Associated With Nitrous Oxide

Originating Committee Clinical Affairs Committee

Review Council Council on Clinical Affairs

> Adopted 1987

Revised 1993, 1996, 2000, 2003

Purpose

The American Academy of Pediatric Dentistry (AAPD) recommends that exposure to ambient nitrous oxide be minimized to reduce occupational health hazards associated with nitrous oxide.

Methods

This document is based on current dental, medical, and public health literature regarding the potential risks of ambient nitrous oxide exposure. Guidelines and recommendations from the National Institute for Occupational Safety and Health (NIOSH) were reviewed also.¹⁻²

Background

Epidemiologic studies provide strong evidence that there are increased general health problems and reproductive difficulties among dental personnel chronically exposed to significant levels of ambient nitrous oxide.³⁻⁷ Nitrous oxide acts by oxidizing vitamin B_{12} from the active, reduced cobalamin to the inactive form. In turn, this inactivates the enzyme methionine synthetase, which requires both the active cobalamin and folate as cofactors. The inactivation of methionine synthetase decreases DNA production, thereby interfering with cell proliferation.⁸

While nitrous oxide has been linked epidemiologically to reproductive, hematologic, immunologic, neurologic, hepatic, and renal disorders, symptoms are time and dose dependent. Neurologic symptoms are reported most frequently in cases of chronic (recreational) abuse. Absolute occupational effects are still uncertain.⁹ Epidemiologic conclusions have been challenged.¹⁰ Adverse reproductive outcomes are linked to B₁₂ deficient individuals and those exposed to "high nitrous oxide levels".⁹ A maximum permissible level of ambient nitrous oxide in the dental environment has not been determined.⁹⁻¹⁴

Collection of ambient nitrous oxide involves 2 separate mechanisms. The first, the scavenging system, is part of the nitrous oxide delivery system. It begins at the nitrous oxide tanks and terminates at the expiratory valve in the mask. Canadian studies in hospital settings have shown that frequent and regular inspection and maintenance of the nitrous oxide delivery system, together with the use of a scavenging system, can reduce ambient nitrous oxide significantly.¹⁵ In the dental environment, patient behaviors such as talking, crying, and moving have been shown to result in significant increases in baseline ambient nitrous oxide levels despite the use of the mask-type scavenging systems.¹⁷ By using a well-fitted mask and an appropriate amount of suction via the scavenging system, the increased pressure on the patient's face by the mask will reduce leakage.¹⁸ NIOSH has recommended an oral evacuation rate of 45 L/min for maximizing scavenger effectiveness. However, scavenging at this rate has been shown to reduce the level of pyschosedation achieved with nitrous oxide inhalation.¹⁹

The second mechanism, the "exhaust system", collects escaped nitrous oxide and includes 2 entities. First, an appropriate nonrecirculating ventilation system is recommended by NIOSH to provide continuous rapid air exchange. It is important to vent waste gases outside of the building and away from fresh air intakes.² Second, a high-volume aspirator, placed near or within 20 cm of the patient's mouth, has been shown to reduce significantly ambient nitrous oxide levels in the dental environment.²⁰⁻²² Diligent use of these 2 mechanisms in the pediatric dental environment has allowed for the reduction of ambient nitrous oxide to the levels recommended by NIOSH.²³

Policy statement

The AAPD recommends that dentists and dental auxiliaries minimize their exposure to nitrous oxide by maintaining the lowest practical levels in the dental environment. Adherence to the recommendations below can help minimize occupational exposure to nitrous oxide.

- 1. Scavenging systems must be used when nitrous oxide is employed.
- 2. Exhaust systems that adequately vent scavenged air and gases to the outside of the building and away from fresh air intake vents should be employed.
- 3. Careful, regular surveillance and maintenance of the nitrous oxide/oxygen delivery equipment must be practiced.
- 4. Mask size should be such as to ensure proper fit for each patient.
- 5. Nitrous oxide discharge from the oral cavity of the patient should be minimized during dental procedures.

References

- 1. National Institute of Occupational Safety and Health. Control of nitrous oxide in dental operatories. Appl Occup Environ Hyg 1999;14:218-220.
- 2. National Institute of Occupational Safety and Health. Controlling exposures of nitrous oxide during anesthetic administration. Cincinnati, Ohio: National Institute of Occupational Safety and Health; 1994. DHHS/NIOSH Publication No. 94-100.
- Rowland AS, Baird DD, Weinberg CR, Shore DL, Shy CM, Wilcox AJ. Reduced fertility among women employed as dental assistants exposed to high levels of nitrous oxide. N Engl J Med 1992;327:993-997.
- 4. Cohen EN, Brown BW Jr, Bruce DL, et al. A survey of anesthetic health hazards among dentists. J Am Dent Assoc 1975;90:1291-1296.
- 5. Cohen EN, Gift HC, Brown BW Jr, et al. Occupational disease in dentistry and chronic exposure to trace anesthetic gases. J Am Dent Assoc 1980;101:21-31.
- Brodsky JB, Cohen EN, Brown BW Jr, Wu ML, Whitcher CE. Exposure to nitrous oxide and neurologic disease among dental professionals. Anesth Analg 1981;60:297-301.
- Rowland AS, Baird DD, Shore DL, Weinberg CR, Savitz DA, Wilcox AJ. Nitrous oxide and spontaneous abortion in female dental assistants. Am J Epidemiol 1995;141:531-537.
- 8. Yagiela JA. Health hazards and nitrous oxide: A time for reappraisal. Anesth Prog 1991;38:1-11.
- 9. Louis-Ferdinand RT. Myelotoxic, neurotoxic, and reproductive adverse effects of nitrous oxide. Adverse Drug React Toxicol Rev 1994;13:193-206.
- Clark MS, Renehan BW, Jeffers BW. Clinical use and potential biohazards of nitrous oxide/oxygen. Gen Dent 1997;45:486-491.
- 11. Howard WR. Nitrous oxide in the dental environment: Assessing the risk and reducing the exposure. J Am Dent Assoc 1997;128:356-360.
- 12. American Denal Association Council on Scientific Affairs, American Dental Association Council on Dental Practice. Nitrous oxide in the dental office. J Am Dent Assoc 1997;128:364-365.

- 13. Donaldson D, Meechan JG. The hazards of chronic exposure to nitrous oxide: An update. Br Dent J 1995; 178:95-100.
- 14. Johnston J. Nitrous oxide: Your health not theirs. Br J Theatre Nurs 1993;3:29-30.
- 15. Sass-Kortasak AM, Wheeler IP, Purdham JT. Exposure of operating room personnel to anaesthetic agents: An examination of the effectiveness of scavenging systems and the importance of maintenance programs. Can Anaesth Soc J 1981;28:22-28.
- 16. Flowerdew RM, Brummitt WM. Reduction of nitrous oxide contamination in a pediatric hospital. Can Anaesth Soc J 1979;26:370-374.
- 17. Henry RJ, Primosch RE, Courts FJ. The effects of various dental procedures and patient behaviors upon nitrous oxide scavenger effectiveness. Pediatr Dent 1992;14:19-25.
- Crouch KG, Johnston OE. Nitrous oxide control in the dental operatory: Auxiliary exhaust and mask leakage, design, and scavenging flow rate as factors. American Industrial Hygiene Association Journal, AIHAJ, 1996;57:272-278.
- 19. Primosch R, McLellan M, Jerrell G, Venezie R. Effect of scavenging on the psychomotor and cognitive function of subjects sedated with nitrous oxide and oxygen inhalation. Pediatr Dent 1992;14:19-25.
- 20. Borganelli GN, Primosch RE, Henry RJ. Operatory ventilation and scavenger evacuation rate influence on ambient nitrous oxide levels. J Dent Res 1993;72:1275-1278.
- 21. Carlsoon P, Ljungqvist B, Hallen B. The effect of local scavenging on occupational exposure to nitrous oxide. Acta Anaesthesiol Scand 1983;27:470-475.
- 22. Henry RJ, Borganelli GN. High-volume aspiration as a supplemental scavenging method for reducing ambient nitrous oxide levels in the operatory: A laboratory study. Int J Paediatr Dent 1995;5:157-161.
- 23. Crouch KG, McGlothin JD, Johnston OE. A longterm study of the development of N₂O controls at a pediatric dental facility. AIHAJ 2000;61:753-756.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.