

Guideline on Fluoride Therapy

Originating Committee
Liaison with Other Groups Committee

Review Council
Council on Clinical Affairs

Adopted
1967

Reaffirmed
1972, 1977

Revised
1978, 1995, 2000, 2003

Purpose

The American Academy of Pediatric Dentistry (AAPD) intends this guideline to help practitioners and parents/caregivers make decisions concerning appropriate use of fluoride as part of the comprehensive oral health care for infants, children, adolescents, and persons with special health care needs.

Methods

A thorough review of the scientific literature pertaining to the use of systemic and topical fluoride was completed to revise and update the guideline on fluoride therapy.

Background

Use of fluorides for the prevention and control of caries is documented to be both safe and highly effective.¹⁻³ Optimizing fluoride levels in water supplies is an ideal public health measure because it is effective and inexpensive and does not require conscious daily cooperation from individuals.³⁻⁴ Daily fluoride exposure through water supplies or supplementation and monitored use of fluoride toothpaste after 6 months of age can be effective primary preventive procedures. Before supplements are prescribed, it is essential to review all dietary sources of fluoride (eg, all drinking water sources such as home, day care, and school, consumed beverages, prepared food, toothpaste) to determine the patient's true exposure to fluoride.^{5,6} The use of fluoridated toothpaste in children who cannot expectorate predictably carries an increased risk of dental fluorosis.⁵⁻⁷ Clinical stud-

ies have shown the effectiveness of a professionally applied topical fluoride treatment.⁸ Eight percent stannous fluoride solution, 1.23% acidulated phosphate fluoride solution or gel, and 5% sodium fluoride varnish are agents used for professionally applied fluoride treatments.^{9,10} Children at higher caries risk¹¹ may require additional fluoride therapies.

Recommendations

Systemically administered fluoride supplements

Fluoride supplements should be considered for all children drinking fluoride-deficient (<0.6 ppm) water. After the fluoride level of the water supply or supplies has been determined, either through contacting public health officials or water analysis, and after evaluating other dietary sources of fluoride and assessing the infant's, child's, or adolescent's caries risk, the daily fluoride supplement dosage schedule can be determined using the Dietary Fluoride Supplementation Schedule (Table 1).

Professionally-applied topical fluoride treatment

Professional topical fluoride treatments should be based on caries-risk assessment.³ A pumice prophylaxis is not an essential prerequisite to this treatment.¹² Appropriate precautionary measures should be taken to prevent swallowing of any professionally-applied topical fluoride.

Self- or parentally-applied fluoride

The use of fluoride-containing toothpaste should be recommended as a primary preventive procedure. Because ingestion of fluoridated toothpaste carries an increased risk of fluorosis, this risk must be weighed against the benefit of caries prevention in determining the use of a fluoridated toothpaste by a child.³ Parents/caregivers should be counseled on the frequency of tooth-brushing and use no more than a "pea-size" amount of toothpaste.¹³

Children at high risk for caries (eg, children with orthodontic/prosthetic appliances, with reduced salivary function, who are unable to clean teeth properly, at dietary risk, having mothers or siblings with caries, or with high oral

Table 1. Dietary Fluoride Supplementation Schedule

Age	<0.3 ppm F	0.3-0.6 ppm F	>0.6 ppm F
Birth-6 mo	0	0	0
6 mo-3 y	0.25 mg	0	0
3-6 y	0.50 mg	0.25 mg	0
6 y up to at least 16 y	1.00 mg	0.50 mg	0

levels of cariogenic bacteria) or children with active caries should be considered for additional fluoride therapy. Home fluoride programs using fluoride mouth rinses or brush-on fluoride gels should be recommended for use by school-aged child at high risk for caries. If a patient at high risk for caries cannot or will not comply with home fluoride therapy, frequent professional fluoride treatments may be substituted.

References

1. Whitford GM. The physiological and toxicological characteristics of fluoride. *J Dent Res* 1990;69(special issue):539-549.
2. Workshop Reports I, II, III from "A symposium on changing patterns of fluoride intake" held at UNC-CH in April 1991. *J Dent Res* 1992;71:1214-1227.
3. CDC. Recommendations for using fluoride to prevent and control dental caries in the United States. *MMWR* 2001;50(RR-14):1-42.
4. CDC. Achievements in Public Health, 1990-1999: Fluoridation of drinking water to prevent dental caries. *JAMA* 2000;283:1283-1286.
5. Levy S, Cohout S, Kiritsy M, Heillman J, Wefel J. Infants fluoride ingestion from water, supplements, and dentifrice. *J Am Dent Assoc* 1995;126:1625-1632.
6. Bowen WH. Fluorosis, is it really a problem? *J Am Dent Assoc* 2002;133:1405-1407.
7. Lalumandier J, Rozier G. The prevalence and risk factors of fluorosis among patients in a pediatric dental practice. *Pediatr Dent* 1995;17:19-25.
8. Ripa LW. In evaluation of the use of professional (operator-applied) topical fluoride. *J Dent Res* 1990;60:86-96.
9. Adair S. Current fluoride therapy in dentistry for children. *Current Opinions in Dentistry* 1991;1:583-591.
10. Bawden JW. Fluoride varnish: A useful new tool for public health dentistry. *J Pub Health Dent* 1998;58:266-269.
11. American Academy of Pediatric Dentistry. Policy on use of a caries-risk assessment tool (CAT) for infants, children, and adolescents. *Pediatr Dent* 2003;25(suppl):18-20.
12. Johnston DW, Lewis DW. Three-year randomized trial of professionally applied topical fluoride gel comparing annual and biannual applications with/without prior prophylaxis. *Caries Res* 1995;29:331-336.
14. Pang D, Vann W. The use of fluoride-containing toothpaste in young children: The scientific evidence for recommending a small quantity. *Pediatr Dent* 1992;14:384-387.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.