Scientific Article



Determining the Prevalence and Risk Factors for Early Childhood Caries in a Community Dental Health Clinic

Robert | Schroth, DMD, MSc¹ • Vivek Cheba, BSc²

Abstract: *Purpose:* The purposes of this study were to: (1) determine the prevalence of early childhood caries (ECC) among young children accessing dental services at a community dental clinic; (2) identify factors associated with the presence of ECC; and (3) determine the percentage of children who received treatment for ECC in this setting and the number who required referral to specialists. *Methods:* The study population comprised children younger than 72 months attending the clinic between 1991 and 2004. A chart review was conducted. *Results:* Eight hundred thirty-four charts met inclusion criteria; 71% had ECC, while the mean deft was 3.7 ± 3.9 (SD). The average age at the first visit was 50.0 ± 12.7 (SD) months. Those with ECC were significantly older at the first visit (P<.001), and the prevalence increased with family size (P=.011) and number of siblings (P=.019). ECC children were significantly more likely to come from households with lower monthly incomes (P=.033). The prevalence of ECC did not vary according to specific areas in Winnipeg where children resided (P=.20). Conclusions: Key risk factors for ECC included: (1) the child's sex; (2) low monthly income; (3) whether the child resided with both parents; and (4) a history of failed dental visits. These data may assist in identifying children at greatest risk for ECC and may help public health agencies develop appropriate prevention strategies, including promoting early dental visits for infants. (Pediatr Dent 2007;29:387-96) Received August 21, 2006 / Revision Accepted January 2, 2007.

KEYWORDS: EARLY CHILDHOOD CARIES, CHILD, PRESCHOOL, COMMUNITY HEALTH CENTERS, PUBLIC HEALTH DENTISTRY, DENTAL CARIES

Young children from low-income households are at increased risk for early childhood caries (ECC). Infants and preschoolers are particularly vulnerable to caries because they rely on parents and caregivers for regular oral hygiene and dietary intakes. Additionally, many dental professionals refuse to see this population in their practices. This risk is further enhanced for those with limited discretionary resources, including families accessing care in publicly funded dental programs and community health centers in Canada.

ECC is a form of decay affecting very young children under the age of 6 years.^{1,2} This broad case definition runs the gamut of decay manifestations from minor to rampant and encompasses all antecedent terms, including "baby bottle tooth decay" and "nursing caries." Severe manifestations can often result in: (1) pain; (2) infections; (3) malnutrition; and (4) a significantly poorer oral health-related quality of life (QOL).³⁻⁵ Oral health QOL, however, often improves once dental treatment is provided.^{6.7} While ECC can have serious and immediate consequences for the child, one often forgotten reality is that decay exhibited during preschool life is also a predictor for future decay in both primary and permanent dentitions.⁸⁻¹⁰

The prevalence of ECC among Canadian children varies considerably, which may be attributed to several factors¹¹⁻¹³¹

- The prevalence may be dependent on the population being studied, as investigations focusing on disadvantaged and low socioeconomic status (SES) populations may naturally report higher prevalences (28%-98%) and rates.^{2,11,12}
- 2. Reports involving younger preschoolers often report a lower prevalence, since the chance of having decay in the primary dentition increases with age.¹⁴
- 3. Case definitions may differ.^{11,15,16} Previous attempts to clinically define forms of primary tooth decay have included specific decayed surfaces of teeth and specific presentation patterns of caries.^{11,15} As these definitions frequently vary, proper comparison of identified risk factors between studies have not always been possible. Only recently has a standardized case definition been published.¹²

¹Dr. Schroth is assistant professor, Faculty of Dentistry, University of Manitoba, Winnipeg, Manitoba, Canada, and researcher, Manitoba Institute of Child Health, Winnipeg, Manitoba, Canada; ²Mr. Cheba is student, Faculty of Dentistry, University of Manitoba. Correspond with Dr. Schroth at umschrot@cc.umanitoba.ca

The etiology underlying ECC was originally ascribed to inappropriate bottle-feeding behaviors (eg, bottle at bedtime or prolonged bottle-feeding) and prolonged breast-feeding.¹⁷⁻²¹ More recently, studies have found other important risk factors while others have found the practice of breastfeeding itself to be protective.²² Numerous contributing risk factors include:

- 1. host factors;
- 2. cariogenic micro-organisms;
- 3. diet;
- 4. individual, familial, and cultural traits such as:
 - a. age;
 - b. ethnicity;
 - c. low SES;
 - d. parental education; and
 - e. mothers with poor oral health and oral health behaviors. $^{\mathtt{2a-28}}$
- 5. limited access to care;
- 6. lack of water fluoridation;
- 7. psychosocial issues;
- 8. microbiological factors (Streptococcus mutans, lactobacilli);
- 9. oral hygiene behaviors;
- 10. the presence of debris, and
- 11. the use of sugar containing products in bottles. 11.29-32

In addition, evidence also indicates that delayed first dental visits may result in greater dental treatment needs.^{33,34}

Many of these contributory factors seem to be more pronounced in areas of low SES, influencing the high degree of caries experienced by inner city residents and those from minority groups. Access to dental care for children from these populations is inadequate. Consequently, one of the few options is to attend publicly funded dental programs, by which time they are often diagnosed with ECC.

The Mount Carmel Clinic (MCC) is a community health center located in the core area of Winnipeg, Manitoba, Canada, that serves an ethnically heterogeneous population of low-income families that predominantly reside in the urban Winnipeg region of Manitoba. It is a nonprofit community health center with the purpose of creating and promoting healthy inner city communities. Families must meet established financial eligibility criteria to qualify for the dental program and pay a minor fee for each dental visit.³⁵ The current assumptions are that the majority of preschool children attending MCC have ECC and that the average age at the first visit is considerably higher than current recommendations.

This study's purposes were to: (1) determine the prevalence of ECC among young children accessing dental services at the clinic; (2) identify factors associated with the presence of ECC; and (3) determine the percentage of children who received treatment for ECC in this setting and the number who required referral to pediatric specialists.

Methods

The study population involved infant and preschool-aged patients who attended the MCC Dental Department between 1991 and 2004. This clinic is located in the Point Douglas community of Winnipeg (Figure 1), the fluoridated capital city of Manitoba with a population of 650,000. Eligibility to attend is based upon a sliding scale of household income. A retrospective chart review was conducted to determine the



Figure 1. Distribution, prevalence of early childhood caries (ECC), and deft rates of children attending the Mount Carmel Clinic based on their neighborhood of residence in Winnipeg (* denotes location of MCC).

prevalence of ECC among young children presenting for dental care. All existing clinical records for preschool-aged children served as the available study sample from 1991-2004. As ECC is age specific, we limited the chart review to those younger than <72 months, based on the definition endorsed by the American Academy of Pediatric Dentistry.^{1,2}

In this institutionally-approved study, a data collection form was developed by the authors to glean information from charts for numerous variables, including: (1) household and family characteristics; (2) dental visitation status; (3) childhood issues; and (4) demographics. The specific community of residence in Winnipeg was determined by reported postal codes. The presence of ECC and primary caries rates (deft), where possible, were determined for each child. The deft score only involved teeth that were extracted, thereby excluding those that had naturally exfoliated. The form was pilot tested prior to the actual data collection. For the majority of charts, information was available and recorded. Data were collected by a single individual (VC).

After the initial data collection period, a random sample of 10% of all charts was re-reviewed to check for reliability of variables collected. Mean values of the 2 data samples were compared using paired t tests to confirm the original data's validity. Data from collection forms were analyzed using the Number Cruncher Statistical System (NCSS, v. 2001; Kaysville, Utah). Statistical analyses included: descriptive tests; and bivariate tests. Descriptive statistics included: (1) mean; (2) standard deviation (±SD); (3) frequencies; and (4) valid percent. Bivariate tests included: (1) analysis of variance (ANOVA); (2) t tests; and (3) chi-square tests. Backwards logistic regression was also employed for ECC. The threshold of significance was $P \le .05$.

Results

Overview of charts. A total of 834 charts met the inclusion criteria. The greatest number was from active patient charts from 2004 (N=85), while the lowest number was from children of the target ages last seen in 1999 (N=42). Ten percent of the charts belonged to children participating in a University of Manitoba dental outreach program that occurred during 1995 and 1996, while 8% (N=64) were derived from charts of children attending the Anne Ross Day Nursery program at MCC, a new outreach initiative at the clinic that began in 2004.

Descriptive statistics. *Geographical*. Seven hundred sixtytwo children resided in Winnipeg (92%), while 70 lived outside the city. Of those from Winnipeg, the majority resided in northern sections of the city, including the following communities (Figure 1): (1) Downtown (23%); (2) Point Douglas (20%); (3) Inkster (9%); (4) River East (12%); and (5) Seven Oaks (10%). This distribution was based upon 759 available observations. Children from outside Winnipeg came from 38 different towns within a 1½-hour driving distance.

Child and family characteristics. Fifty-three percent of the subjects were male (Table 1). Most children (70%) belonged to families consisting of 4 or more persons, and 80% of children had at least 1 or more siblings. The average total family size was 4.2 ± 1.4 persons and the average number of siblings was 1.5 ± 1.2 . On average, total family monthly income was \$1,481.88±673.14 (all dollar amounts are Canadian [Cdn]). Few children had documented medical conditions (13%, N=111). Among those with medical issues, allergies (42%) and asthma (41%) were the most prevalent.

Caries experience and dental visitation status. Overall, 71% of children were found to have at least 1 primary tooth affected by decay (ECC prevalence), while the estimated mean deft for the sample was 3.7 ± 3.9 (range=0-20). Mean scores for d, e, and f components and overall deft appear in Table 2.

The average age at the first dental visit was 50.0 ± 12.7 months (4 years of age), and the average age at the time of first diagnosis of caries was 52.4 ± 12.3 months, with both ranging between 11 and 71 months. Only 5 children came in by age 1 for their first dental visit (1%), while 15 were under 2 years of age (2%; Table 1). In total, 19% of children (N=154) attending MCC were referred for further care. Of those referred, 94% were referred to a pediatric dentist. The majority of preschool children (84%) referred outside the clinic were referred by dental therapists employed at MCC. Twenty-six percent were documented to have failed at least 1 dental appointment.

Bivariate associations. *Geographical.* The prevalence of ECC among Winnipeg children did not significantly differ from rural Manitoba children (70% vs 78%; chi-square=1.5; df=1; P=.21), but the mean deft score did differ between rural children and those from Winnipeg (4.7±4.1 vs 3.6±3.9 [range=0-20]; P=.047), suggesting that children outside the capital city have increased caries rates.

The proportion of children affected with decay differed according to their distinct Winnipeg communities. Overall, children attending MCC who lived in the Inkster area had the highest prevalence of ECC (79%) while those living in St. Boniface had the lowest (17%; chi-square=21.5; df=11; P=.o3). ANOVA, however, revealed that there was no significant association between the mean deft score and the community of residence (P=.o6; Figure 1). A post hoc chi-square analysis excluding the St. Boniface neighborhood revealed no statistical difference in ECC prevalence (P=.14), indicating that the original significant difference observed between the distinct Winnipeg communities was entirely due to this one neighborhood.

Table 1. CHILD AND	D FAMILY C	HARACTERIS	TICS AND DENTAL VISITATION CHARACTERI	STICS			
Variable	N	Valid %	Variable	N	Valid %		
Sex			Maternal employment status	Maternal employment status			
Male	441	53	Employed	286	35		
Female	391	47	Unemployed	479	58		
Age (ys)			Unsure	55	7		
1	7	1	Paternal employment status				
2	28	3	Employed	434	52		
3	85	10	Unemployed	136	17		
4	157	19	Unusre	260	31		
5	555	67	Documented medical conditions	Documented medical conditions			
Family Size			Yes	Yes 111			
2	74	10	No	721	87		
3	157	20	Early childhood caries				
4	251	32	Yes	588	71		
5	165	21	No	244	29		
6	84	11	Age at first dental visit (mos)	Age at first dental visit (mos)			
7	31	4	≤ 23	15	2		
≥ 8	18	2	24-25	91	11		
No. of siblings			36-47	260	31		
0	157	20	48-59	235	28		
1	281	36	60-71	231	28		
2	188	24	Form of Payment				
3	99	13	Self	705	84		
4	37	5	Insurance	29	4		
≥ 5	18	2	Employment and income assistance (social assistance)	90	11		
Monthly household income (Cdn)			Non-Insured Health Benefits (treaty status First Nations)	7	1		
\$ 0-1,000	199	28	Failed dental appointment				
\$ 1,001-2,000	360	51	Yes	220	26		
\$ 2,001-4,000	152	21	No	No 612			
No. of Parents			Treatment completed at Mount Ca	Treatment completed at Mount Carmel Clinic			
1	228	27	Yes	542	65		
2	603	73	No	280	34		
Space Maintainer			Unsure	Unsure 10 1			
Yes	29	4	Child referred to another provider				
No	803	96	Yes	154	19		
Age at first diagnosis	of caries (m	nos)	No	678	81		
≤ 23	6	1					
24-25	46	8					
36-47	156	27					
48-59	182	31					
60-7	193	33					

Child and family characteristics. There were no significant relationships between both the prevalence of ECC or rates of primary caries and the sex of the child (73% males vs 69% females, P=.2; 3.9 ± 4.0 [range=0-20] males vs 3.5 ± 3.9 [range=0-17] females, P=.13). The prevalence of ECC, however, was significantly associated with: (1) family size (chi-square=16.6; df=6; P=.01); (2) the number of siblings in the family unit (chi-square=13.5; df=5; P=.02); and (3) the child's age (chi-square=41.4; df=4; P<.001).

Generally, as the number of family members, number of siblings, and age increased so did the prevalence of ECC. On the other hand, ANOVA results indicated that only the number of siblings and age, but not total family size, were significantly associated with increased deft scores (P=.03, P<.001, and P=.14 respectively). ANOVA also revealed that the mean family size (4.3 ± 1.4 vs 4.2 ± 1.4) and the mean number of siblings (1.5 ± 1.2 siblings vs 1.5 ± 1.2) was not significantly associated with the presence of ECC (P=.40 and P=.80, respectively).

Table 2.MEAN NUMBER (±SD) OF DECAYED (d), EXTRACTED (e), AND FILLED (f) PRIMARY TEETH AND DEFT						
	N	Mean±SD	Range			
d score	753	1.6±2.9	0-16			
Age 1		1.4±1.9	0-4			
Age 2		0.05±0.2	0-1			
Age 3		2.1±3.7	0-14			
Age 4		1.9±3.2	0-16			
Age 5		1.6±2.7	0-15			
e score	753	0.4±1.0	0-8			
Age 1		0	0			
Age 2	0		0			
Age 3	0.03±0.2 0		0-2			
Age 4	0.2±0.8		0-6			
Age 5		0.4±1.0	0-8			
f score	753	1.7±2.6	0-16			
Age 1		0 0				
Age 2		0.05±0.2 0-1				
Age 3		0.4±1.5 0-11				
Age 4		0.8±1.7 0-9				
Age 5		2.2±2.8	0-16			
deft score	753	3.7±3.9	0-20			
Age 1		1.4±1.9	0-4			
Age 2		0.1±0.3	0-1			
Age 3		2.5±3.8	0-14			
Age 4		2.9±4.0	0-20			
Age 5		4.2±3.8	0-16			

Children from single-parent homes did not experience significantly more decay (chi-square=2.5; df=1; P=.11), but did appear to have lower caries rates than those from 2parent homes (t test, P=.057). The family's monthly income bracket (\$500 Cdn increments) was not found to be associated with the presence of decay (chi-square=7.2; df=5; P=.20) or rates of decay (P=.56). A post hoc decision was made to divide monthly income by the number of family members in the household to create a new variable of monthly income per family member, since all families were considered to be low-income and fell into similar income brackets. The average was \$359.82 Cdn/member/month±152.90. Children with decay were more likely to come from families with lower monthly income per family member compared with those who were caries free (\$351.95/month/member±147.09 vs \$378.98/month/member±164.99, P=.03).

Chi-square analysis revealed that the presence of decay was not associated with the employment status of either the mother (chi-square=0.41, df=1, P=.50; 73% employed vs 71% unemployed) or the father (chi-square=0.002, df=1, P=.96; 73% employed vs 73% unemployed). ANOVA also revealed there was no significant relationship between the employment status of each parent and mean deft (P=.054 and P=.11, respectively, for mothers and fathers). Two-tailed t tests were used to compare deft rates for the employed and unemployed mothers and fathers, as the unsure category was likely influencing the analyses. Results revealed that there were no differences between maternal and paternal employment status and deft scores (P=.08 and P=.8, respectively).

While 104 children had documented underlying medical conditions, there were no significant differences in prevalence (67% vs 71%) or rates of decay (3.4 ± 3.7 [range=0-15] vs 3.8 ± 4.0 [range=0-20]) between healthy children and those with medical issues.

Dental visitation status. The prevalence of ECC and the mean deft scores remained relatively unchanged from 1991 and 2004 (chi-square=15.2, df=13, P=0.40; ANOVA P=.17). The prevalence of ECC ranged from its lowest of 57% in 2003 to its highest of 80% in 1994. Interestingly, children who had failed scheduled dental appointments were significantly more likely to have ECC (80% vs 68%; odds ratio=1.9) and had an increased mean deft score (Table 3).

There were also significant associations between ECC and the extent of decay based on the child's age at the time of their first visit (Table 4). ANOVA also revealed that ECC children were significantly older at the first visit (51.4 ± 12.3 months vs 46.6 ± 13.2 months; P < .001). No significant difference existed between the mean deft scores and the age when children were first diagnosed with caries (P=.10).

Finally, statistical analyses revealed that, over the years, there was a significant association between the mean age at

No. with ECC (%)	No. caries free (%)	Mean deft±SD
33 (72) †	13 (28)	3.6±3.9 (range=0-15) †
113 (72)	45 (28)	3.5±3.6 (range=0-15)
168 (73)	61 (27)	3.9±4.0 (range=0-17)
98 (75)	33 (25)	4.2±4.0 (range=0-18)
68 (69)	31 (31)	3.5±3.7 (range=0-15)
30 (57)	23 (43)	3.5±4.7 (range=0-20)
No. with ECC (%)	No. caries free (%)	Mean deft±SD
175 (80) ‡	45 (20)	4.4±3.7 (range=0-18) §
413 (68)	199 (32)	3.5±4.0 (range=0-20)
No. with ECC (%)	No. caries free (%)	Mean deft±SD
7 (47) ‡	8 (53)	1.4±1.9 (range=0-5)‡
51 (56)	40 (44)	2.2±3.5 (range=0-17)
174 (67)	86 (33)	3.5±4.2 (range=0-20)
177 (75)	58 (25)	4.4±4.0 (range=0-16)
179 (78)	52 (22)	4.0±3.6 (range=0-14)
	No. with ECC (%) 33 (72) † 113 (72) 168 (73) 98 (75) 68 (69) 30 (57) No. with ECC (%) 175 (80) ‡ 413 (68) No. with ECC (%) 7 (47) ‡ 51 (56) 174 (67) 177 (75) 179 (78)	No. with ECC (%) No. caries free (%) 33 (72) † 13 (28) 113 (72) 45 (28) 168 (73) 61 (27) 98 (75) 33 (25) 68 (69) 31 (31) 30 (57) 23 (43) No. with ECC (%) No. caries free (%) 175 (80) ‡ 45 (20) 413 (68) 199 (32) No. with ECC (%) No. caries free (%) 7 (47) ‡ 8 (53) 51 (56) 40 (44) 174 (67) 86 (33) 177 (75) 58 (25) 179 (78) 52 (22)

Lable 3. PREVALENCE OF EARLY CHILDHOOD CARIESEARLY CHILDHOOD CARIES (ECC) AND MEAN DEFT BY TOTAL MONTHLY INCOME (CDN), FAILED DENTAL APPOINTMENTS. AND AGE UPON FIRST DENTAL VISIT*

* Chi-square for ECC; ANOVA for deft. $\ddagger P > .05$. $\ddagger P < .001$. $\S P < .01$.

the time of the first clinic visit and the year archived (P<.001). A Tukey's post hoc test showed that there was a significant difference in age at the first visit during 2003 and 2004 compared to every year except 1999 and 2001. During 2003 and 2004, the children visiting the clinic for a first visit were significantly younger (45.1 ± 14.4 and 42.1 ± 14.0 months) than the other years (range= 49.2 ± 11.3 [1995]- 55.0 ± 14.1 [2002]).

Logistic regression. Backwards logistic regression was performed, with the final model appearing in Table 4. Variables removed from the model during the backward variable iteration included: (1) single child (P=.89); (2) father's employment (P=.62); (3) medical condition (P=.34); and (4) family size (P=.34).

Discussion

Relatively few chart reviews have been conducted to determine the prevalence of preschool dental decay in clinical environments.³⁶⁻³⁸To date, there have been no published chart reviews in Canada examining the prevalence and risk factors for ECC among children attending dental public health programs.

Overall, 71% of the eligible children in this retrospective chart review had ECC. Of those with ECC. 5.5±3.6 teeth were affected by decay. The prevalence of ECC in this population is cause for concern, but not unexpected, since all were from low-income households. In fact, children from lower-income families were more likely to suffer from ECC. Decay is prominent in disadvantaged populations, including those with limited access to care. Two previous retrospective studies in the United States involving clinical records reported similar prevalence rates (67% and 76%), while a third reported a much lower prevalence (34%).36-38 The deft in this study's sample also approximates recent published rates of primary tooth decay among disadvantaged preschool children in the Point Douglas community of Winnipeg, where MCC is located."

Undoubtedly, age is a significant predictor of ECC. Not surprisingly, both the prevalence of ECC and the rate of decay increased as the age at the first visit increased. Children

who visit the dentist at later ages have more teeth for longer lengths of time, increasing their odds of developing decay. Therefore, primary health centers such as MCC must become dental homes for high-risk infants and preschoolers.

The concept of the "dental home" is a relatively new one and likely not well disseminated among general practitioners.³⁹ Private dental offices and public health clinics can serve as "dental homes" for young children by providing care and prevention services that meet each child's unique needs,^{39.4°} including: (1) caries risk assessments; (2) prevention, including fluoride varnish and other chemotherapeutics; (3) anticipatory guidance; and (4) parental education.^{39.4°}

Organized dentistry recommends a first dental visit by 12 months of age.^{41,42} The average age at the first visit for MCC children, however, was significantly higher (50.0 ± 12.7 months) than the current recommendation. This study confirms that dental visits before 2 years of age are important to keep young children free of cavities. Screening children within months of eruption of their first tooth is integral to ensuring that they receive early primary prevention, thus setting the foundation for good childhood oral health.⁴³ Further, early preventive care can provide parents with much needed dental anticipatory guidance so that they are empowered to practice dental friendly behaviors.^{39,41}

Considering MCC's mandate to serve low SES populations, those drawing on the services are at increased risk for dental disease, including ECC. Public health clinics such as MCC must begin to promote and institute first visits by 1 year of age to curb the problem of ECC. The profession and public health agencies, however, must first ensure that the primary focus applies to those at greatest risk with the

Variable	Regression coefficient ± (SD)	P-value	Odds ratio	95% confidence interval	
Sex (male)	0.324±0.163	.048	1.38	1.003-1.904	
Community of residence (urban)	-0.606±0.323	.060	0.55	0.290-1.026	
Age at the first dental	-1643+0630	009	0.19	0.056-0.664	
visit (<24 mos)	1.04520.050	.005	0110		
Low monthly income (≤\$2,000/mo)	0.581±0.236	.014	1.79	1.125-2.841	
Low monthly income per family member (<\$325/member/mo)	0.319±0.189	.092	1.28	0.949-1.993	
Maternal employment status (employed)	0.351±0.182	.054	1.42	0.995-2.030	
Failed dental appointments	0.597±0.196	.002	1.82	1.237-2.667	
Single parent	-0.487±0.192	.011	0.61	0.421-0.896	

least access before such policies apply to the entire populace.44 Without such a strategic approach, it is quite possible that those with limited access may be unable to obtain visits. This is because infants from middle and upper income groups may overwhelm the profession's ability to undertake such a preventive service, given provider shortages and the limited number of those practitioners willing to see infants in their practices. Awareness of this recommendation has grown at MCC, which may account for the significant decline in the mean age of children attending for the first time during 2003 and 2004. This change, however, is more likely due to one dentist conducting a study on ECC during infancy. The high age at the first visit may be due to parents bringing in their children only when dental problems arise. Many other barriers to early and timely care exist, including: (1) parental education; (2) dentists refusing to see preschool-aged children; (3) unreliable transportation; (4) discrimination and delays in seeing a dentist; and (5) a lack of cultural sensitivity on the part of service providers.^{29,45-48}

By not promoting early childhood oral health care, the dental profession continues to propagate the myth that baby teeth are of little value to overall childhood well-being and dental development.⁴⁹

The prevalence of decay did not significantly differ between children residing in Winnipeg and those residing outside the capital city. This finding indicates that the need for dental care among disadvantaged preschoolers from rural Manitoba is just as large and should not be neglected. Several studies have reported that children residing in rural locales are more likely to suffer from decay in the primary dentition.50-53 Access to affordable, community-based dental care is severely limited for children in rural Manitoba. In addition, they may not be benefiting from water fluoridation and are dependent on reliable transportation to visit the dentist. Children who attended MCC and lived in the Inkster, Point Douglas, downtown, and River East communities had a higher prevalence of ECC than those from other Winnipeg communities, although the overall prevalence by community was not significant. This comes as little surprise, since these communities surround MCC and are considered to have more high-risk families. The 2004 Community Health Assessment (CHA) report noted that these communities share patterns of poor health, including: (1) chronic health conditions; (2) infant and maternal health; (3) communicable diseases; (4) injury; (5) mental health; and (6) death.53 Not surprisingly, the oral health of children from these areas is also less favorable. Overall, very few children were from the St. Boniface community of Winnipeg, a francophone section of the city. The lack of bilingual oral health services at MCC may account for the very low usage from this neighborhood and the low prevalence of ECC exhibited.

Children from single-parent homes did not experience a greater prevalence or rates of decay than those living with 2 parents. In fact, the former were more likely to be caries free. This result was somewhat surprising, since previous studies have reported higher rates of ECC in single-parent homes.^{22,54} The mean family size was significantly smaller in single-parent households (3.1±1.1 vs 4.7±1.2; P<.001). While this association does not allow us to develop a cause-andeffect relationship, it possibly suggests that children from smaller families might have less ECC because they may have more time to devote to early childhood oral health care and oral hygiene.^{12,55}

SES has been reported as a significant risk factor for ECC in a substantial number of studies.^{24,56} The CHA reported that the Point Douglas and downtown areas have the highest percentage of families living below the low-income cutoff. These communities represented nearly 60% of this study's population, withall subjects having high rates of ECC. The relationship between low-income and caries was confirmed in this study.

Logistic regression revealed that boys were more likely to have increased odds of ECC (OR=1.3). Most ECC studies have not found such a relationship to be true, but this does agree with findings from another investigation of high-risk aboriginal preschool children.⁵⁷

Irregular dental attendance was a predictor of caries among young children at MCC. Those who failed scheduled dental appointments were more likely to have ECC and had higher mean deft scores than those who did not miss appointments (OR=1.9). The issue at hand for this population may not entirely be due to parents neglecting childhood oral health, but rather other factors, some of which were previously discussed. Access to care for this population is a huge barrier, since many families attending the clinic lack reliable transportation. The following factors may also lead to broken appointments: (1) a lack of integration of oral health with primary health care; (2) previous bad dental experiences; (3) racial and ethnic barriers; and (4) limited hours of operation.^{47,48}

It is also probable that some parents fail to take their child for scheduled visits because they cannot afford to miss work, as this would result in lost income.

General limitations of retrospective chart reviews include: (1) the fact that important data may not be available (ie, missing data and limited variables); (2) difficulty deciphering providers progress notes; (3) charting errors; (4) difficulty in controlling bias and confounders; and (5) difficulty in establishing a cause-and-effect relationship.

Limitations of this study included:

- 1. missing information;
- 2. patient records lacking information on:
 - a. family income;
 - b. total family size;
 - c. parental employment status; and
 - d. medical conditions;
- the uncertainty of certain variables, such as: a. decayed (d);
 - b. extracted (e):

c. filled (f); and

d. deft; and

4. differing chart documentation habits of the providers.

The format of charts themselves also changed during the period under review. The majority of the service providers, however, remained constant over this period.

Although the authors assumed that the child's age at the time of their first visit to MCC was the true age when they first visited the dentist, this study was unable to determine whether children had received care prior to attending MCC. Given the challenges faced by this population in accessing care, the assumption is likely. This limitation, however, potentially changes the average age at the first visit. Recently, MCC has recognized the need to be more responsive to the growing health needs of populations residing in the Point Douglas and Downtown communities bordering the facility, especially those of Aboriginal ancestry. Strategies to achieve such improvements include: (1) focusing on community development; (2) early childhood development and care; (3) primary health for the economically disadvantaged; and (4) integrated programming. Early dental visits should be included in these strategies as a means to improve preschool oral health. Many people residing in the clinic's catchment area may already have existing dental benefits from the Non-Insured Health Benefits Program (NIHB) of the First Nations and Inuit Health Program or social assistance. Considering that MCC dental department has targeted the low-income sector, it has historically focused on the working poor, encouraging those with benefits to access care from other sources.

Unfortunately, the evidence is clear that dental benefits for vulnerable groups do not translate into better utilization or improved dental outcomes.^{58,59} In fact, only 22% of Canadian First Nations preschool children with NIHB for dental care had a dental visit during 2002/03.⁶⁰ Thus, it is imperative that the MCC and other community agencies continue to identify the needs in the community and work towards reducing health disparities and improving access to the underserved.

Conclusions:

Based on this study's results, the following conclusions can be made:

- The majority of preschool children attending the Mount Carmel Clinic had ECC.
- 2. Factors significantly associated with ECC on logistic regression analysis included: (a) sex of the child; (b) first dental visit at no earlier than 24 months of age; (c) low monthly household income; (d) residing with both parents; and (e) a history of failed dental visits.
- 3. Efforts should be made to follow current recommendations for a first dental visit by 12 months of age.

Acknowledgements

The authors wish to thank the Mount Carmel Clinic and Ms. Lori Black, dental program manager, for providing access to clinic records. Mr. Cheba received a BSc(Dent) summer studentship from the Manitoba Institute of Child Health a division of the Children's Hospital Foundation of Manitoba Inc, Winnipeg, Manitoba, Canada. Dr. Schroth is a Canadian Institutes of Health Research Strategic Training Fellow in the Canadian Child Health Clinician Scientist Program (www. cchcsp.ca, Shared Partners: SickKids Foundation, Child and Family Research Institute/BC Children's Hospital Foundation, and the Manitoba Institute of Child Health—a division of the Children's Hospital Foundation of Manitoba, Inc).

References

- American Academy of Pediatric Dentistry. Definition of Early Childhood CariesEarly childhood caries (ECC). Pediatr Dent. 2006;28 (suppl):13.
- 2. Drury TF, Horowitz AM, Ismail AI, et al. Diagnosing and reporting early childhood caries for research purposes. A report of a workshop sponsored by the National Institute of Dental and Craniofacial Research, the Health Resources and Services Administration, and the Health Care Financing Administration. J Public Health Dent 1999;59:192-7.
- 3. Low W, Tan S, Schwartz S. The effect of severe caries on the quality of life in young children. Pediatr Dent 1999;21:325-6.
- Filstrup SL, Briskie D, da Fonseca M, et al. Early childhood caries and quality of life: Child and parent perspectives. Pediatr Dent 2003;25:431-40.
- 5. Clarke M, Locker D, Berall G, et al. Malnourishment in a population of young children with severe early childhood caries. Pediatr Dent 2006;28:254-9.
- Acs G, Pretzer S, Foley M, Ng MW. Perceived outcomes and parental satisfaction following dental rehabilitation under general anesthesia. Pediatr Dent 2001;23:419-23.
- White H, Lee JY, Vann WF, Jr. Parental evaluation of quality of life measures following pediatric dental treatment using general anesthesia. Anesth Prog 2003;50:105-10.
- Kaste LM, Marianos D, Chang R, Phipps KR. The assessment of nursing caries and its relationship to high caries in the permanent dentition. J Public Health Dent 1992;52:64-8.
- Johnsen DC, Gerstenmaier JH, DiSantis TA, Berkowitz RJ. Susceptibility of nursing-caries children to future approximal molar decay. Pediatr Dent 1986;8:168-70.
- al Shalan TA, Erickson PR, Hardie NA. Primary incisor decay before age 4 as a risk factor for future dental caries. Pediatr Dent 1997;19:37-41.
- Schroth RJ, Moore P, Brothwell DJ. Prevalence of early childhood caries in 4 Manitoba communities. J Can Dent Assoc 2005;71:567-67f.

- Schroth RJ, Moffatt ME. Determinants of early childhood caries (ECC) in a rural Manitoba community: A pilot study. Pediatr Dent 2005;27:114–20.
- 13. Schroth RJ, Smith PJ, Whalen JC, Lekic C, Moffatt ME. Prevalence of caries among preschool-aged children in a northern Manitoba community. J Can Dent Assoc 2005;71:27-27f.
- Weinstein P, Smith WF, Fraser-Lee N, Shimono T, Tsubouchi J. Epidemiologic study of 19-month-old Edmonton, Alberta children: Caries rates and risk factors. J Dent Child 1996;63:426-33.
- Ismail AI, Sohn W. A systematic review of clinical diagnostic criteria of early childhood caries. J Public Health Dent 1999;59:171-91.
- Albert RJ, Cantin RY, Cross HG, Castaldi CR. Nursing caries in the Inuit children of the Keewatin. J Can Dent Assoc 1988;54:751-8.
- Dilley GJ, Dilley DH, Machen JB. Prolonged nursing habit: A profile of patients and their families. J Dent Child 1980;47:102-8.
- 18. Ripa LW. Nursing habits and dental decay in infants: "Nursing bottle caries." J Dent Child 1978;45:274-5.
- 19. Richardson BD, Cleaton-Jones PE. Nursing bottle caries. Pediatrics 1977;60:748-9.
- Bottle baby syndrome. Contact Intraocul Lens Med J 1966;11:12.
- Smith PJ, Moffatt ME. Baby-bottle tooth decay: Are we on the right track? Int J Circumpolar Health 1998;57 (suppl 1):155-62.
- 22. Hallett KB, O'Rourke PK. Early childhood caries and infant feeding practice. Community Dent Health 2002;19:237-42.
- 23. Reisine S, Douglass JM. Psychosocial and behavioral issues in early childhood caries. Community Dent Oral Epidemiol 1998;26:32-44.
- 24. Hallett KB, O'Rourke PK. Social and behavioural determinants of early childhood caries. Aust Dent J 2003;48:27-33.
- 25. Dini EL, Holt RD, Bedi R. Caries and its association with infant feeding and oral health-related behaviours in 3- to 4-year-old Brazilian children. Community Dent Oral Epidemiol 2000;28:241-8.
- Grytten J, Rossow I, Holst D, Steele L. Longitudinal study of dental health behaviors and other caries predictors in early childhood. Community Dent Oral Epidemiol 1988;16:356-9.
- 27. Schou L, Uitenbroek D. Social and behavioural indicators of caries experience in 5-year-old children. Community Dent Oral Epidemiol 1995;23:276-81.
- Sasahara H, Kawamura M, Kawabata K, Iwamoto Y. Relationship between mothers' gingival condition and caries experience of their 3-year-old children. Int J Paediatr Dent 1998;8:261-7.
- 29. Mouradian WE, Wehr E, Crall JJ. Disparities in children's oral health and access to dental care. JAMA 2000; 284:2625-31.

- 30. Booth JM, Mitropoulos CM, Worthington HV. A comparison between the dental health of 3-year-old children living in fluoridated Huddersfield and nonfluoridated Dewsbury in 1989. Community Dent Health 1992;9:151-7.
- Grindefjord M, Dahllof G, Modeer T. Caries development in children from 2.5 to 3.5 years of age: A longitudinal study. Caries Res 1995;29:449-54.
- Wendt LK, Birkhed D. Dietary habits related to caries development and immigrant status in infants and toddlers living in Sweden. Acta Odontol Scand 1995;53:339-44.
- Melhado FL, Cunha RF, Nery RS. Influence of dental care for infants on caries prevalence: A comparative study. J Dent Child 2003;70:120-3.
- Savage MF, Lee JY, Kotch JB, Vann WF, Jr. Early preventive dental visits: Effects on subsequent utilization and costs. Pediatrics 2004;114:e418-23.
- Mount Carmel Clinic. Mount Carmel Clinic Dental Program. Available at: "www.mountcarmel.ca/programs.dental.htm". Accessed March 11, 2005.
- Nainar SM, Crall JJ. Caries experience in inner-city preschoolers at the time of their initial dental visit. J Dent Child 1997;64:421-4.
- Albert DA, Park K, Findley S, Mitchell DA, McManus JM. Dental caries among disadvantaged 3- to 4-year-old children in northern Manhattan. Pediatr Dent 2002;24:229-33.
- Quartey JB, Williamson DD. Prevalence of early childhood caries at Harris County clinics. J Dent Child 1999; 66:85, 127-31.
- 39. Hale KJ. Oral health risk assessment timing and establishment of the dental home. Pediatrics 2003;111:1113-6.
- 40. Crall JJ. Development and integration of oral health services for preschool-age children. Pediatr Dent 2005;27:323-30.
- 41. American Acadmey of Pediatric Dentistry. Policy on Early Childhood Cariesearly childhood caries (ECC): Classifications, consequences, and preventive strategies. Pediatr Dent 2006;28(suppl):34-6.
- 42. Canadian Dental Association. Your child's first visit. Available at: "http://www.cda-adc.ca/english/your_oral_health/ caring_teeth/info_for_parents_teachers/first_visit.asp". Accessed August 2, 2005.
- 43. Lee JY, Bouwens TJ, Savage MF, Vann WF, Jr. Examining the cost-effectiveness of early dental visits. Pediatr Dent 2006;28:102-5.
- Jones K, Tomar SL. Estimated impact of competing policy recommendations for age at the first dental visit. Pediatrics 2005;115:906-14.
- 45. Casamassimo PS. Dental disease prevalence, prevention, and health promotion: The implications on pediatric oral health of a more diverse population. Pediatr Dent 2003;25:16-8.

- Lam M, Riedy CA, Milgrom P. Improving access for Medicaid-insured children: Focus on front-office personnel. J Am Dent Assoc 1999;130:365-73.
- 47. Riedy CA, Weinstein P, Milgrom P, Bruss M. An ethnographic study for understanding children's oral health in a multicultural community. Int Dent J 2001;51:305-12.
- 48. Mofidi M, Rozier RG, King RS. Problems with access to dental care for Medicaid-insured children: What caregivers think. Am J Public Health 2002;92:53-8.
- 49. Weinstein P. Provider versus patient-centered approaches to health promotion with parents of young children: What works/does not work and why. Pediatr Dent 2006; 28:172-6.
- 50. Barnes GP, Parker WA, Lyon TC, Jr., Drum MA, Coleman GC. Ethnicity, location, age, and fluoridation factors in baby bottle tooth decay and caries prevalence of Head Start children. Public Health Rep 1992;107:167-73.
- Marino RJ, Onetto JE. Caries experience in urban and rural Chilean 3-year-olds. Community Dent Oral Epidemiol 1995;23:60-1.
- 52. Mattila ML, Rautava P, Sillanpaa M, Paunio P. Caries in five-year-old children and associations with family-related factors. J Dent Res 2000;79:875-81.
- 53. Winnipeg Regional Health Authority. Community Health Assessment Report 2004. Winnipeg, Manitoba: Winnipeg Regional Health Authority; 2004.
- 54. Crall JJ, Edelstein B, Tinanoff N. Relationship of microbiological, social, and environmental variables to caries status in young children. Pediatr Dent 1990;12:233-6.
- 55. Muller M. Nursing-bottle syndrome: Risk factors. J Dent Child 1996;63:42-50.
- Moynihan PJ, Holt RD. The national diet and nutrition survey of 1.5- to 4.5-year-old children: Summary of the findings of the dental survey. Br Dent J 1996;181:328-32.
- Seow WK, Amaratunge A, Bennett R, Bronsch D, Lai PY. Dental health of aboriginal preschool children in Brisbane, Australia. Community Dent Oral Epidemiol 1996; 24:187-90.
- 58. Non-Insured Health Benefits Program Annual Report 2003/2004. Ottawa, Ontario: First Nations and Inuit Health Branch Non-Insured Health Benefits Directorate Program Analysis Division, Health Canada; 2005.
- Ismail AI, Sohn W. The impact of universal access to dental care on disparities in caries experience in children. J Am Dent Assoc 2001;132:295-303.
- 60. Non-Insured Health Benefits Annual Report 2002/2003. Ottawa, Ontario: First Nations and Inuit Health Branch, Non-Insured Health Benefits Directorate, Program Analysis Division, Health Canada; 2003.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.