Case Report

Laser-assisted Treatment of a Dentigerous Cyst: Case Report

Juan R. Boj, DDS, PhD¹ • Clervie Poirier, DDS, PhD² • Miguel Hernandez, DDS, PhD³ • Enric Espasa, DDS, PhD⁴

Abstract: Dentigerous cysts are benign maxillary odontogenic cysts associated with the crown of an unerupted tooth. They can expand the cortical bone to such an extent that they cause displacement of teeth and root resorption in the adjacent teeth. The purpose of this report was to describe the clinical case of a dentigerous cyst in a mandibular permanent molar of a 6-year-old child. The cyst caused the displacement of the bud of the permanent second molar and root resorption of the primary second molar. The cyst was treated by: (1) mucous fenestration using an erbium laser: (2) drainage of the fluid content; and (3) curettage of the bone cavity. The injury was successfully resolved in just 5 months with: (1) bone regeneration; (2) the repositioning of the displaced bud; and (3) correct eruption of the affected tooth. (Pediatr Dent 2007;29:521-4) Received December 6, 2006 / Revision Accepted February 20, 2007.

KEYWORDS: DENTIGEROUS CYST, PEDIATRIC DENTISTRY, LASER DENTISTRY

Dentigerous or follicular cysts are maxillary benign odontogenic cysts generally associated with the crown of an impacted, embedded, or unerupted permanent tooth.^{1,2} The cyst enclosing the unerupted tooth's crown is attached to the tooth's cervical region.³ The eruption cyst is the analogous case when it appears in the soft tissue, shortly before tooth eruption.⁴ Despite there being a number of theories about their origin,⁵ both seem to result from the separation of the epithelium from the enamel of the tooth's crown due to an accumulation of blood or other fluid in a dilated follicular space.⁶

In the dental literature, these cysts are considered to be the second most common odontogenic cysts of the jaws, after radicular cysts.⁷ They are more common in male patients and usually appear during the second or third decade of life.^{1.8} In mixed dentition, they represent less than 9% of dentigerous cysts.⁹ In 75% of cases, they are located in the mandible.10 In order, the teeth most often affected are: (1) mandibular third molars; (2) maxillary third molars; and (3) maxillary permanent canines.^{8,11} They have also been reported in the mandibular second premolars.³

Correspond with Dr. Boj at 16388jrb@comb.es

These lesions are often detected during a routine radiographic examination. This is because dentigerous cysts are typically painless, unless there is acute inflammatory exacerbation^{1,7,8} (ie, in mixed dentition when a necrotic primary tooth infects the follicle of the permanent successor).³⁹ Main designated such cysts as inflammatory follicular cysts.¹² Pain is a sign of secondary infection on these sterile cysts.¹³ If the cyst reaches a large size¹⁴ (>2 cm in diameter), we may observe facial swelling due to its potential for expanding the cortical bone.^{17,9}

Dentigerous cysts are usually solitary, and the radiographic pattern is characterized by a symmetric, well-defined, unilocular radiolucent lesion surrounding the unerupted tooth's crown, which can displace adjacent teeth and cause root resorption.^{1,11}

Clinical examination and radiography provide a preliminary diagnosis, but a definitive diagnosis requires histopathological analysis.⁷ Differential diagnoses should include: (1) hyperplasia of the tooth follicle¹; (2) ameloblastic fibroma^{1,15}; (3) unicystic ameloblastomas¹⁶; (4) glandular odontogenic cyst¹⁷; (5) odontogenic keratocyst¹⁸; and (6) adenomatoid odontogenic tumor.¹⁹

Several treatments have been proposed, including:

- 1. total enucleation7;
- 2. marsupialization²⁰;
- decompression opening the cyst and ensuring continual drainage^{1,7,9};
- combination of marsupialization or decompression and enucleation^{1,7,21}; and

¹Dr. Boj is chaîrman. ²Dr. Poirier is associate professor, 3Dr. Hernandez is professor, and ⁴Dr. Espasa is professor, all at Department of Pediatric Dentistry, Dental School, University of Barcelona, Spain.

 decompression combined with extraction of the causative infected tooth in inflammatory cysts.^{3:9}

This article describes the case of a dentigerous cyst treated with an Er, Cr:YSGG laser (erbium, chromium, yttrium, scandium, gallium, and garnet; Waterlase-Biolase Technology, San Clemente, Calif). This type of laser is a hydrokinetic system that liberates photons in an air-water spray and works at a wavelength of 2,780 nm. The effect of this laser with its wavelength is favored by the water present in soft tissues.²⁴ The laser's energy is carried by a system of fiber optics to a terminal point made of a sapphire crystal.²³

Case report

A 6-year-old boy was brought to the Department of Pediatric Dentistry at the Dental School of the University of Barcelona, Barcelona, Spain, because his family had noticed a swelling on his face at the mandible level on the right-hand side. The patient was asymptomatic and had no history of pain. The medical history was noncontributory.

In the extraoral examination, slight facial asymmetry was observed, with swelling at the right mandibular angle. There was, however, no noticeable presence of submandibular or cervical adenopathy. Intraoral examination revealed a bluish area on the gingiva (Figure 1) above the unerupted permanent mandibular right first molar. There was little buccolingual expansion of the bone.

The panoramic radiograph revealed a large unilocular radiolucency (osteolytic lesion) measuring 1.6 cm in diameter and associated with the crown of an unerupted permanent mandibular right first molar with open apices. There was resorption of the distal root of the primary mandibular second molar and distal displacement of the permanent second molar bud (Figure 2).

The lesion's clinical appearance was compatible with a dentigerous cyst.We decided to incise the upper limit of the



Figure 1. Clinical lesion demonstrating bluish appearance and handpiece with sapphire tip of the erbium laser.



Figure 3. Initial ablation with the laser. The amber-yellowish color of the cystic liquid can be appreciated.

cyst to expose the cavity. A total of 36 mg of 2% lidocaine with epinephrine 1:100,000 in a mandibular block and buccal infiltration was used for local anesthesia. The settings for the laser were those recommended by the manufacturer for soft tissue surgery (1.50 watts, 10% water, and 11% air). The sap-





Figure 2. Panoramic radiograph demonstrating a dentigerous cyst related to an unerupted permanent mandibular first molar, with displacement of the permanent second molar bud and resorption of the distal root of the primary second molar.



Figure 4. Intraoperative appearance of the molar in the cystic cavity. Rough laser incisions with tissue tags can be seen. The margins of the incision were subsequently smoothed with the laser. A cleaner incision margin could have been achieved by the use of a scalpel blade.

The patient was observed after surgical treatment, and no symptoms or signs of recurrence were detected. Five months after treatment, a panoramic radiograph revealed that the cystic lesion had disappeared and there was new bone growth (Figure 5). The unerupted molar had reached a proper position in the arch without orthodontic traction. Bone regeneration was also observed around the resorbed root of the mandibular primary second molar, which presented no signs of the pulp being affected. Furthermore, the permanent second molar's bud had corrected its position within the bone. The only pathologic finding was that the molar associated with the cyst was hypocalcified, while none of the other permanent molars was affected (Figure 6).

Discussion

The case described presents all the typical characteristics of a

dentigerous cyst—they:

1. are more frequent in $men^{7,11}$;

2. appear at the neck of an unerupted tooth's crown;

3. displace teeth; and

4. resorb roots as they grow.

The discovery of a hypomineralized molar in the cyst could be just a coincidence, due to the high incidence of this anomaly in the last decade.

Early diagnosis of this type of cyst in children is important, as growth can be rapid and can cause bone fractures and deformation.¹ Eruption cysts usually do not require treatment, and the affected tooth erupts normally. In the

Figure 5. Panoramic radiograph 5 months after surgery showing the resolution of the lesion.



Figure 6. Erupted hypomineralized permanent mandibular first molar 5 months after surgery.

dense connective fibrous wall covered with a fine layer of nonkeratinized squamous epithelium and containing variable amounts of myxoid tissue and odontogenic epithelial remnants. case of dentigerous cysts, however, treatment is always necessary.⁴ Children have a much greater capacity than adults to regenerate bone and teeth with open apices have a greater eruptive potential.^{9:20:24} These factors should make one consider large dentigerous cysts in children as entities distinct from those in adults, with much better prognosis for the teeth involved.

We employed conservative treatment. Exposing the cyst, aspirating the fluid content, and curetting the cavity proved to be sufficient. The erbium laser provided a good vision of the operational field, as we could take advantage of its coagulating effect, even though this effect is limited compared to $\rm CO_2$ lasers.²⁵ It also provides the possibility of remodelling the bone if necessary.^{26,27} The antibacterial²⁸ and anti-in-flammatory properties attributed to the laser may improve postoperative prognosis.²⁹ The only medication the patient required was 200 mg of ibuprofen on the day of the surgery. No sutures had been used, and the soft tissues scarred quickly and without any problems.

Kozelj et al.⁹ Ertas et al.²⁰ and Martinez et al²⁴ have also presented cases of conservative treatment in the mixed dentition, which permitted spontaneous eruption of the affected permanent tooth. In these cases, complete ossification of the bony defect occurred in 1 or 2 years. In the present case, ossification occurred in a shorter time since the origin was not due to an infection of a primary tooth as in the cases presented by these other authors.

The laser technology currently available can considerably help in the treatment of some cases that require surgery in pediatric dentistry.^{3°} Erbium lasers are the most versatile for pediatric dental patients, as they can be used for both soft and hard tissues. They are an additional tool for improving and modernizing the authors' treatments.

References

- Pinkham JR, Casamassimo PS, Fields HW, Mc Tigue DJ, Nowak AJ. *Pediatric Dentistry: Infancy Through Adolescence*. 4th ed. St. Louis, Mo: Elsevier Saunders; 2005;47.
- 2. Miller CS, Bean LR. Pericoronal radiolucencies with and without radiopacities. Dent Clin North Am 1994:38:51-61.
- Rubin DM, Vedrenne D, Portnof JE. Orthodontically guided eruption of mandibular second premolar following enucleation of an inflammatory cyst: Case report. J Clin Pediatr Dent 2002;27:19–24.
- Boj JR, Poirier C, Espasa E, Hernandez M, Jacobson B. Eruption cyst treated with a laser powered hydrokinetic system. J Clin Pediatr Dent 2006;30:199-202.
- Bodner L, Goldstein J, Sarnat H. Eruption cysts: A clinical report of 24 new cases. J Clin Pediatr Dent 2004;28:183-6.
- 6. Boj JR, García-Godoy F. Multiple eruption cysts: Report of a case. J Dent Child 2000;67:282-4.
- Delbem ACB, Cunha RF, Afonso RL, Bianco KG, Idem AP. Dentigerous cysts in the primary dentition: Report of 2 cases. Pediatr Dent 2006;28:269-72.
- Smith JL, Kellman RM. Dentigerous cysts presenting as head and neck infections. Otolaryngol Head Neck Surg 2005;133:715-7.
- Ko_elj V, Soto_ek B. Inflammatory dentigerous cysts of children treated by tooth extraction and decompression: Report of four cases. Br Dent J 1999;187:587-90.
- Weber AL. Imaging of the cysts and odontogenic tumors of the jaw: Definition and classification. Radiol Clin North Am 1993;31:101-20.
- Thoma KH. The circumferential dentigerous cyst. Oral Surg 1964;18:368-71.
- Main DMG. Epithelial jaw cysts: 10 years of the WHO classification. J Oral Pathol 1985;14:1-7.
- August M, Faquin WC, Troulis M, et al. Differentiation of odontogenic keratocysts from nonkeratinizing cysts by use of fine-needle aspiration biopsy and cytokeratin-10 staining. J Oral Maxillofac Surg 2000;58:935-40.

- Bodner L, Woldenberg Y, Bar-Ziv J. Radiographic features of large cysts lesions of the jaws in children. Pediatr Radiol 2003;33:3-6.
- Kim SG, Jang HS. Ameloblastic fibroma: Report of a case. J Oral Maxillofac Surg 2002;60:216-8.
- Patel H, Rees RT. Unicystic ameloblastoma presenting in Gardner's syndrome: A case report. Br Dent J 2005; 198:747-8.
- Kasabo_lu O, Ba_al Z, Usubütün A. Glandular odontogenic cyst presenting as a dentigerous cyst: A case report. J Oral Maxillofac Surg 2006;64:731-3.
- Stoll C, Stollenwerk C, Riediger D, Mittermayer C, Alfer J. Cytokeratin expression patterns for distinction of odontogenic keratocysts from dentigerous and radicular cysts. J Oral Pathol Med 2005;34:558-64.
- Bravo M, White D, Miles L, Cotton R. Adenomatoid odontogenic tumor mimicking a dentigerous cyst. Int J Pediatr Otorhinolaryngol 2005;69:1685-8.
- 20. Ertas Ü, Selim Yavuz M. Interesting eruption of 4 teeth associated with a large dentigerous cyst in the mandible by only marsupialization. J Oral Maxillofac Surg 2003; 61:728-30.
- Motamedi MHK, Talesh KT. Management of extensive dentigerous cysts. Br Dent J 2005;198:203-6.
- 22. Eversole LR, Rizoiu IM. Preliminary investigations on the utility of an erbium, chromium YSGG laser. J Calif Dent Assoc 1995;23:41-7.
- 23. Jacobson B, Berger J, Kravitz R, Ko J. Laser pediatric Class II composites utilizing no anesthesia. J Clin Pediatr Dent 2004;28:99-101.
- 24. Martínez D, Varela M. Conservative treatment of dentigerous cysts in children: A report of 4 cases. J Oral Maxillofac Surg 2001;59:331-4.
- Pavelec V, Polenik P. Use of Er, Cr:YSGG versus standard lasers in laser-assisted uvulopalatoplasty for treatment of snoring. Laringoscope 2006;116:1512-6.
- Kimura Y, Yu DG, Fujita A, Yamashita A, Murakami Y, Matsumoto K. Effects of erbium, chromium: YSGG laser irradiation on canine mandibular bone. J Periodontol 2001;72:1178-82.
- 27. Wang X, Zhang C, Matsumoto K. In vivo study of the healing processes that occur in the jaws of rabbits following perforation by an Er, Cr: YSGG laser. Lasers Med Sci 2005;20:21-7.
- Turkun M, Turkun LS, Celik EU, Ates M. Bactericidal effect of Er, Cr: YSGG laser on streptococcus mutans. Dent Mater J 2006;25:81-6.
- 29. Walinski CJ. Irritation fibroma removal: A comparison of two laser wavelengths. Gen Dent 2004;52:236-8.
- Boj JR. The future of laser pediatric dentistry. J Oral Laser Applic 2005;5:173-7

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.