

# Guideline on for Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures

## Developed and Endorsed by

American Academy of Pediatrics and the American Academy of Pediatric Dentistry

## Adopted

2006

### Abstract

The safe sedation of children for procedures requires a systematic approach that includes the following: no administration of sedating medication without the safety net of medical supervision, careful presedation evaluation for underlying medical or surgical conditions that would place the child at increased risk from sedating medications, appropriate fasting for elective procedures and a balance between depth of sedation and risk for those who are unable to fast because of the urgent nature of the procedure, a focused airway examination for large tonsils or anatomic airway abnormalities that might increase the potential for airway obstruction, a clear understanding of the pharmacokinetic and pharmacodynamic effects of the medications used for sedation as well as an appreciation for drug interactions, appropriate training and skills in airway management to allow rescue of the patient, age- and size-appropriate equipment for airway management and venous access, appropriate medications and reversal agents, sufficient numbers of people to both carry out the procedure and monitor the patient, appropriate physiologic monitoring during and after the procedure, a properly equipped and staffed recovery area, recovery to presedation level of consciousness before discharge from medical supervision, and appropriate discharge instructions.

### Introduction

Invasive diagnostic and minor surgical procedures on pediatric patients outside the traditional operating room setting have increased in the last decade. As a consequence of this change and the increased awareness of the importance of providing analgesia and anxiolysis, the need for sedation for procedures in physician offices, dental offices, subspecialty procedure suites, imaging facilities, emergency departments, and ambulatory surgery centers also has markedly increased.<sup>1-37</sup> In recognition of this need for both elective and emergency use of sedation in nontraditional settings, the American Academy of Pediatrics (AAP) and American Academy of Pediatric Dentistry (AAPD) have published a series of guidelines for the monitoring and management of pediatric patients during and after sedation for a procedure.<sup>38-42</sup> The purpose of this updated statement is to unify

the guidelines for sedation used by medical and dental practitioners, add clarifications regarding monitoring modalities, provide new information from medical and dental literature, and suggest methods for further improvement in safety and outcomes. With the revision of this document, the Joint Commission on Accreditation of Healthcare Organizations, the American Society of Anesthesiologists (ASA), the AAP, and the AAPD will use similar language to define sedation categories and the expected physiologic responses.<sup>41-44</sup>

This revised statement reflects the current understanding of appropriate monitoring needs both during and after sedation for a procedure.<sup>4-5,12,19,21,22,26,45-53</sup> The monitoring and care outlined in this guideline may be exceeded at any time, based on the judgment of the responsible practitioner. Although intended to encourage high-quality patient care, adherence to this guideline cannot guarantee a specific patient outcome. However, structured sedation protocols designed to incorporate the principles in this document have been widely implemented and shown to reduce morbidity.<sup>29,32-34,37,54,55</sup> This guideline is proffered with the awareness that, regardless of the intended level of sedation or route of administration, the sedation of a pediatric patient represents a continuum and may result in respiratory depression and the loss of the patient's protective reflexes.<sup>43,57-60</sup>

Sedation of pediatric patients has serious associated risks, such as hypoventilation, apnea, airway obstruction, laryngospasm, and cardiopulmonary impairment.<sup>2,6,22,45,46,54,60-69</sup> These adverse responses during and after sedation for a diagnostic or therapeutic procedure may be minimized, but not completely eliminated, by a careful preprocedure review of the patient's underlying medical conditions and consideration of how the sedation process might affect or be affected by these conditions.<sup>54</sup> Appropriate drug selection for the intended procedure as well as the presence of an individual with the skills needed to rescue a patient from an adverse response are essential. Appropriate physiologic monitoring and continuous observation by personnel not directly involved with the procedure allow for accurate and rapid diagnosis of complications and initiation of appropriate rescue interventions.<sup>46,51,54</sup>

The sedation of children is different from the sedation of adults. Sedation in children often is administered to control behavior to allow the safe completion of a procedure. A child's ability to control his or her own behavior to cooperate for a procedure depends both on his or her chronologic and developmental age. Often, children younger than 6 years and those with developmental delay require deep levels of sedation to gain control of their behavior.<sup>57</sup> Therefore, the need for deep sedation should be anticipated. Children in this age group are particularly vulnerable to the sedating medication's effects on respiratory drive, patency of the airway, and protective reflexes.<sup>46</sup> Studies have shown that it is common for children to pass from the intended level of sedation to a deeper, unintended level of sedation.<sup>56,59-70</sup> For older and cooperative children, other modalities, such as parental presence, hypnosis, distraction, topical local anesthetics, and guided imagery, may reduce the need for or the needed depth of pharmacologic sedation.<sup>31,71-81</sup>

The concept of rescue is essential to safe sedation. Practitioners of sedation must have the skills to rescue the patient from a deeper level than that intended for the procedure. For example, if the intended level of sedation is "minimal," practitioners must be able to rescue from "moderate sedation"; if the intended level of sedation is "moderate," practitioners must have the skills to rescue from "deep sedation"; if the intended level of sedation is "deep," practitioners must have the skills to rescue from a state of "general anesthesia." The ability to rescue means that practitioners must be able to recognize the various levels of sedation and have the skills necessary to provide appropriate cardiopulmonary support if needed. Sedation and anesthesia in a nonhospital environment (private physician or dental office or freestanding imaging facility) may be associated with an increased incidence of "failure to rescue" the patient should an adverse event occur, because the only backup in this venue may be to activate emergency medical services (EMS).<sup>46,82</sup> Rescue therapies require specific training and skills.<sup>46,54,83,84</sup> Maintenance of the skills needed to perform successful bag-valve-mask ventilation is essential to successfully rescue a child who has become apneic or developed airway obstruction. Familiarity with emergency airway management procedure algorithms is essential.<sup>83-87</sup> Practitioners should have an in-depth knowledge of the agents they intend to use and their potential complications. A number of reviews and handbooks for sedating pediatric patients are available.<sup>32,48,55,88-93</sup> This guideline is intended for all venues in which sedation for a procedure might be performed (hospital, surgical center, freestanding imaging facility, dental facility, or private office).

There are other guidelines for specific situations and personnel that are beyond the scope of this document. Specifically, guidelines for the delivery of general anesthesia and monitored anesthesia care (sedation or analgesia), outside or within the operating room by anesthesiologists or other practi-

tioners functioning within a department of anesthesiology, are addressed by policies developed by the ASA and by individual departments of anesthesiology.<sup>94</sup> Also, guidelines for the sedation of patients undergoing mechanical ventilation in a critical care environment or for providing analgesia for patients postoperatively, patients with chronic painful conditions, and hospice care are beyond the scope of this document.

### Definitions of Terms for This Report

- "Pediatric patients": all patients through 21 years of age, as defined by the AAP.
- "Must" or "shall": an imperative need or duty that is essential, indispensable, or mandatory.
- "Should": the recommended need and/or duty.
- "May" or "could": freedom or liberty to follow a suggested or reasonable alternative.
- "Medical supervision" or "medical personnel": a current, licensed practitioner in medicine, surgery, or dentistry trained in the administration of medications used for procedural sedation and the management of complications associated with these medications.
- "Are encouraged": a suggested or reasonable action to be taken.
- "ASA Physical Status Classification": guidelines for classifying the baseline health status according to the ASA (see Appendix B).
- "Minimal sedation" (old terminology "anxiolysis"): a drug-induced state during which patients respond normally to verbal commands. Although cognitive function and coordination may be impaired, ventilatory and cardiovascular functions are unaffected.
- "Moderate sedation" (old terminology "conscious sedation" or "sedation/analgesia"): a drug-induced depression of consciousness during which patients respond purposefully to verbal commands (eg, "open your eyes" either alone or accompanied by light tactile stimulation; a light tap on the shoulder or face, not a sternal rub). For older patients, this level of sedation implies an interactive state; for younger patients, age-appropriate behaviors (eg, crying) occur and are expected. Reflex withdrawal, although a normal response to a painful stimulus, is not considered as the only age-appropriate purposeful response (ie, it must be accompanied by another response, such as pushing away the painful stimulus so as to confirm a higher cognitive function). With moderate sedation, no intervention is required to maintain a patent airway, and spontaneous ventilation is adequate. Cardiovascular function is usually maintained. However, in the case of procedures that may themselves cause airway obstruction (eg, dental or endoscopic), the practitioner must recognize an obstruction and assist the patient in opening the airway. If the patient is not making spontaneous efforts to open his/her airway so as to relieve the obstruction, then the patient should be considered to be deeply sedated.
- "Deep sedation" ("deep sedation/analgesia"): a drug-

induced depression of consciousness during which patients cannot be easily aroused but respond purposefully (see discussion of reflex withdrawal above) after repeated verbal or painful stimulation (eg, purposefully pushing away the noxious stimuli). The ability to independently maintain ventilatory function may be impaired. Patients may require assistance in maintaining a patent airway, and spontaneous ventilation may be inadequate. Cardiovascular function is usually maintained. A state of deep sedation may be accompanied by partial or complete loss of protective airway reflexes.

- “General anesthesia”: a drug-induced loss of consciousness during which patients are not arousable, even by painful stimulation. The ability to independently maintain ventilatory function is often impaired. Patients often require assistance in maintaining a patent airway, and positive-pressure ventilation may be required because of depressed spontaneous ventilation or drug-induced depression of neuromuscular function. Cardiovascular function may be impaired.

## Goals of Sedation

The goals of sedation in the pediatric patient for diagnostic and therapeutic procedures are: 1) to guard the patient's safety and welfare; 2) to minimize physical discomfort and pain; 3) to control anxiety, minimize psychological trauma, and maximize the potential for amnesia; 4) to control behavior and/or movement so as to allow the safe completion of the procedure; and 5) to return the patient to a state in which safe discharge from medical supervision, as determined by recognized criteria, is possible (Appendix A).

These goals can best be achieved by selecting the lowest dose of drug with the highest therapeutic index for the procedure. It is beyond the scope of this document to specify which drugs are appropriate for which procedures; however, the selection of the fewest number of drugs and matching drug selection to the type and goal of the procedure are essential for safe practice.<sup>53,88,91-93,95-97</sup> For example, analgesic medications such as opioids are indicated for painful procedures. For nonpainful procedures, such as computed tomography or magnetic resonance imaging (MRI), sedatives/hypnotics are preferred. When both sedation and analgesia are desirable (eg, fracture reduction), either single agents with analgesic/sedative properties or combination regimens commonly are used. Anxiolysis and amnesia are additional goals that should be considered in selection of agents for particular patients. However, the potential for an adverse outcome may be increased when 3 or more sedating medications are administered.<sup>44,98</sup> Knowledge of each drug's time of onset, peak response, and duration of action is essential. Although the concept of titration of drug to effect is critical, one must know whether the previous dose has taken full effect before administering additional drug. Such management will improve safety and outcomes. Drugs with long durations of action (eg, chloral hydrate, intramuscular pentobarbital,

phenothiazines) will require longer periods of observation even after the child achieves currently used recovery and discharge criteria.<sup>45,99,100</sup> This concept is particularly important for infants and toddlers transported in car safety seats who are at risk of resedation after discharge because of residual prolonged drug effects with the potential for airway obstruction.<sup>45,46</sup>

## General Guidelines

### Candidates

Patients who are in ASA classes I and II are frequently considered appropriate candidates for minimal, moderate, or deep sedation (Appendix B). Children in ASA classes III and IV, children with special needs, and those with anatomic airway abnormalities or extreme tonsillar hypertrophy present issues that require additional and individual consideration, particularly for moderate and deep sedation.<sup>51</sup> Practitioners are encouraged to consult with appropriate subspecialists and/or an anesthesiologist for patients at increased risk of experiencing adverse sedation events because of their underlying medical/surgical conditions.

### Responsible Person

The pediatric patient shall be accompanied to and from the treatment facility by a parent, legal guardian, or other responsible person. It is preferable to have 2 or more adults accompany children who are still in car safety seats if transportation to and from a treatment facility is provided by 1 of the adults.<sup>101</sup>

### Facilities

The practitioner who uses sedation must have immediately available facilities, personnel, and equipment to manage emergency and rescue situations. The most common serious complications of sedation involve compromise of the airway or depressed respirations resulting in airway obstruction, hypoventilation, hypoxemia, and apnea. Hypotension and cardiopulmonary arrest may occur, usually from inadequate recognition and treatment of respiratory compromise. Other rare complications may also include seizures and allergic reactions. Facilities providing pediatric sedation should monitor for, and be prepared to treat, such complications.

### Back-up Emergency Services

A protocol for access to back-up emergency services shall be clearly identified, with an outline of the procedures necessary for immediate use. For nonhospital facilities, a protocol for ready access to ambulance service and immediate activation of the EMS system for life-threatening complications must be established and maintained. It should be understood that the availability of EMS services does not replace the practitioner's responsibility to provide initial rescue in managing life-threatening complications.

### On-Site Monitoring and Rescue Equipment

An emergency cart or kit must be immediately accessible. This cart or kit must contain equipment to provide the necessary age- and size-appropriate drugs and equipment to resuscitate a nonbreathing and unconscious child. The contents of the kit must allow for the provision of continuous life support while the patient is being transported to a medical facility or to another area within a medical facility. All equipment and drugs must be checked and maintained on a scheduled basis (see Appendices C and D for suggested drugs and emergency life support equipment to consider before the need for rescue occurs). Monitoring devices, such as electrocardiography (ECG) machines, pulse oximeters (with size-appropriate oximeter probes), end-tidal carbon dioxide monitors, and defibrillators (with size-appropriate defibrillator paddles), must have a safety and function check on a regular basis as required by local or state regulation.

### Documentation Before Sedation

Documentation shall include, but not be limited to, the guidelines that follow:

1. Informed consent. The patient record shall document that appropriate informed consent was obtained according to local, state, and institutional requirements.<sup>102</sup>
2. Instructions and information provided to the responsible person. The practitioner shall provide verbal and/or written instructions to the responsible person. Information shall include objectives of the sedation and anticipated changes in behavior during and after sedation. Special instructions shall be given to the adult responsible for infants and toddlers who will be transported home in a car safety seat regarding the need to carefully observe the child's head position so as to avoid airway obstruction. Transportation by car safety seat poses a particular risk for infants who have received medications known to have a long half-life, such as chloral hydrate, intramuscular pentobarbital, or phenothiazine.<sup>45,46,100,103</sup> Consideration for a longer period of observation shall be given if the responsible person's ability to observe the child is limited (eg, only 1 adult who also has to drive). Another indication for prolonged observation would be a child with an anatomic airway problem or a severe underlying medical condition. A 24-hour telephone number for the practitioner or his or her associates shall be provided to all patients and their families. Instructions shall include limitations of activities and appropriate dietary precautions.

### Dietary Precautions

Agents used for sedation have the potential to impair protective airway reflexes, particularly during deep sedation. Although a rare occurrence, pulmonary aspiration may occur if the child regurgitates and cannot protect his or her airway. Therefore, it is prudent that before sedation, the practitioner evaluate preceding food and fluid intake. It is likely that the risk of aspiration during procedural sedation differs from that during general anesthesia

involving tracheal intubation or other airway manipulation.<sup>104,105</sup> However, because the absolute risk of aspiration during procedural sedation is not yet known, guidelines for fasting periods before elective sedation generally should follow those used for elective general anesthesia. For emergency procedures in children who have not fasted, the risks of sedation and the possibility of aspiration must be balanced against the benefits of performing the procedure promptly (see below). Further research is needed to better elucidate the relationships between various fasting intervals and sedation complications.

### Before Elective Sedation

Children receiving sedation for elective procedures should generally follow the same fasting guidelines as before general anesthesia (Table 1). It is permissible for routine necessary medications to be taken with a sip of water on the day of the procedure.

### For the Emergency Patient

The practitioner must always balance the possible risks of sedating nonfasted patients with the benefits and necessity for completing the procedure. In this circumstance, the use of sedation must be preceded by an evaluation of food and fluid intake. There are few published studies with adequate statistical power to provide guidance to the practitioner regarding safety or risk of pulmonary aspiration of gastric contents during procedural sedation.<sup>104-109</sup> When protective airway reflexes are lost, gastric contents may be regurgitated into the airway. Therefore, patients with a history of recent oral intake or with other known risk factors, such as trauma, decreased level of consciousness, extreme obesity, pregnancy, or bowel motility dysfunction, require careful evaluation before administration of sedatives. When proper fasting has not been ensured, the increased risks of sedation must be carefully weighed against its benefits, and the lightest effective sedation should be used. The use of agents with less risk of depressing protective airway reflexes may be preferred.<sup>110</sup> Some emergency patients requiring deep sedation may require protection of the airway before sedation.

### Use of Immobilization Devices

Immobilization devices, such as papoose boards, must be applied in such a way as to avoid airway obstruction or chest restriction. The child's head position and respiratory excursions should be checked frequently to ensure airway patency. If an immobilization device is used, a hand or foot should be kept exposed, and the child should never be left unattended. If sedating medications are administered in conjunction with an immobilization device, monitoring must be used at a level consistent with the level of sedation achieved.

### Documentation at the Time of Sedation

1. Health evaluation. Before sedation, a health evaluation shall be performed by an appropriately-licensed practitioner and reviewed by the sedation team at the time of treatment for

possible interval changes. The purpose of this evaluation is not only to document baseline status but also to determine whether patients present specific risk factors that may warrant additional consultation before sedation. This evaluation will also screen out patients whose sedation will require more advanced airway or cardiovascular management skills or alterations in the doses or types of medications used for procedural sedation.

A new concern for the practitioner is the widespread use of medications that may interfere with drug absorption or metabolism and, therefore, enhance or shorten the effect time of sedating medications. Herbal medicines (eg, St John's wort, echinacea) may alter drug pharmacokinetics through inhibition of the cytochrome P450 system, resulting in prolonged drug effect and altered (increased or decreased) blood drug concentrations.<sup>111-116</sup> Kava may increase the effects of sedatives by potentiating gamma-aminobutyric acid inhibitory neurotransmission, and valerian may itself produce sedation that apparently is mediated through modulation of gamma-aminobutyric acid neurotransmission and receptor function.<sup>117,118</sup> Drugs such as erythromycin, cimetidine, and others may also inhibit the cytochrome P450 system, resulting in prolonged sedation with midazolam as well as other medications competing for the same enzyme systems.<sup>119-122</sup> Medications used to treat human immunodeficiency virus infection, some anticonvulsants, and some psychotropic medications may also produce clinically important drug-drug interactions.<sup>123-125</sup> Therefore, a careful drug history is a vital part of the safe sedation of children. The clinician should consult various sources (a pharmacist, textbooks, online services, or handheld databases) for specific information on drug interactions.<sup>126</sup>

The health evaluation should include:

- Age and weight
- Health history, including: 1) allergies and previous allergic or adverse drug reactions; 2) medication/drug history, including dosage, time, route, and site of administration for prescription, over-the-counter, herbal, or illicit drugs; 3) relevant diseases, physical abnormalities, and neurologic impairment that might increase the potential for airway obstruction, such as a history of snoring or obstructive sleep apnea<sup>127,128</sup>; 4) pregnancy status; 5) a summary of previous relevant hospitalizations; 6) history of sedation or general anesthesia and any complications or unexpected responses; and 7) relevant family history, particularly related to anesthesia
- Review of systems with a special focus on abnormalities of cardiac, pulmonary, renal, or hepatic function that might alter the child's expected responses to sedating/analgesic medications
- Vital signs, including heart rate, blood pressure, respiratory rate, and temperature (for some children

who are very upset or noncooperative, this may not be possible and a note should be written to document this occurrence)

- Physical examination, including a focused evaluation of the airway (tonsillar hypertrophy, abnormal anatomy—eg, mandibular hypoplasia) to determine whether there is an increased risk of airway obstruction<sup>54,129,130</sup>
- Physical status evaluation (ASA classification [see Appendix B])
- Name, address, and telephone number of the child's medical home

For hospitalized patients, the current hospital record may suffice for adequate documentation of presedation health; however, a brief note shall be written documenting that the chart was reviewed, positive findings were noted, and a management plan was formulated. If the clinical or emergency condition of the patient precludes acquiring complete information before sedation, this health evaluation should be obtained as soon as feasible.

2. Prescriptions. When prescriptions are used for sedation, a copy of the prescription or a note describing the content of the prescription should be in the patient's chart along with a description of the instructions that were given to the responsible person. **Prescription medications intended to accomplish procedural sedation must not be administered without the benefit of direct supervision by trained medical personnel.** Administration of sedating medications at home poses an unacceptable risk, particularly for infants and preschool-aged children traveling in car safety seats.<sup>46</sup>

### Documentation During Treatment

The patient's chart shall contain a time-based record that includes the name, route, site, time, dosage, and patient effect of administered drugs. Before sedation, a "time out" should be performed to confirm the patient's name, procedure to be performed, and site of the procedure.<sup>43</sup> During administration, the inspired concentrations of oxygen and inhalation sedation agents and the duration of their administration shall be documented. Before drug administrations, special attention must be paid to calculation of dosage (ie, mg/kg). The patient's chart shall contain documentation at the time of treatment that the patient's level of consciousness and responsiveness, heart rate, blood pressure, respiratory rate, and oxygen saturation were monitored until the patient attained predetermined discharge criteria (see Appendix A). A variety of sedation scoring systems are available and may aid this process.<sup>70,100</sup> Adverse events and their treatment shall be documented.

### Documentation After Treatment

The time and condition of the child at discharge from the treatment area or facility shall be documented; this should include documentation that the child's level of consciousness and oxygen saturation in room air have returned to a state that is safe

for discharge by recognized criteria (see Appendix A). Patients receiving supplemental oxygen before the procedure should have a similar oxygen need after the procedure. Because some

sedation medications are known to have a long half-life and may delay a patient's complete return to baseline or pose the risk of re sedation,<sup>45,103,131,132</sup> some patients might benefit from a longer period of less-intense observation (eg, a step-down observation area) before discharge from medical supervision.<sup>133</sup> Several scales to evaluate recovery have been devised and validated.<sup>70,134,135</sup> A recently described and simple evaluation tool may be the ability of the infant or child to remain awake for at least 20 minutes when placed in a quiet environment.<sup>100</sup>

### Continuous Quality Improvement

The essence of medical error reduction is a careful examination of index events and root cause analysis of how the event could be avoided in the future.<sup>137-141</sup> Therefore, each facility should maintain records that track adverse events, such as desaturation, apnea, laryngospasm, the need for airway interventions including jaw thrust, positive pressure ventilation, prolonged sedation, unanticipated use of reversal agents, unintended or prolonged hospital admission, and unsatisfactory sedation/analgesia/anxiolysis. Such events can then be examined for assessment of risk reduction and improvement in patient satisfaction.

### Preparation and Setting up for Sedation Procedures

Part of the safety net of sedation is to use a systematic approach so as to not overlook having an important drug, piece of equipment, or monitor immediately available at the time of a developing emergency. To avoid this problem, it is helpful to use an acronym that allows the same setup and checklist for every procedure. A commonly used acronym useful in planning and preparation for a procedure is **SOAPME**:

- S** = Size-appropriate **suction** catheters and a functioning **suction** apparatus (eg, Yankauer-type suction)
- O** = An adequate **oxygen** supply and functioning flow meters/other devices to allow its delivery
- A** = **Airway**: size-appropriate airway equipment (nasopharyngeal and oropharyngeal airways, laryngoscope blades [checked and functioning], endotracheal tubes, stylets, face mask, bag-valve-mask or equivalent device [functioning])
- P** = **Pharmacy**: all the basic drugs needed to support life during an emergency, including antagonists as indicated
- M** = **Monitors**: functioning pulse oximeter with size-appropriate oximeter probes<sup>141,142</sup> and other monitors as appropriate for the procedure (eg, noninvasive blood pressure, end-tidal carbon dioxide, ECG, stethoscope)
- E** = Special **equipment or drugs** for a particular case (eg, defibrillator)

## Specific Guidelines for Intended Level of Sedation

### Minimal Sedation

Minimal sedation (old terminology "anxiolysis") is a drug-induced state during which patients respond normally to verbal commands. Although cognitive function and coordination may be impaired, ventilatory and cardiovascular functions are unaffected. Children who have received minimal sedation generally will not require more than observation and intermittent assessment of their level of sedation. Some children will become moderately sedated despite the intended level of minimal sedation; should this occur, then the guidelines for moderate sedation apply.<sup>57</sup>

### Moderate Sedation

"Moderate sedation" (old terminology "conscious sedation" or "sedation/analgesia") is a drug-induced depression of consciousness during which patients respond purposefully to verbal commands or following light tactile stimulation (see Definition of Terms for This Report). No interventions are required to maintain a patent airway, and spontaneous ventilation is adequate. Cardiovascular function usually is maintained. The caveat that loss of consciousness should be unlikely is a particularly important aspect of the definition of moderate sedation. The drugs and techniques used should carry a margin of safety wide enough to render unintended loss of consciousness highly unlikely. Because the patient who receives moderate sedation may progress into a state of deep sedation and obtundation, the practitioner should be prepared to increase the level of vigilance corresponding to what is necessary for deep sedation.<sup>57</sup>

### Personnel

#### The Practitioner

The practitioner responsible for the treatment of the patient and/or the administration of drugs for sedation must be competent to use such techniques, to provide the level of monitoring provided in this guideline, and to manage complications of these techniques (ie, to be able to rescue the patient). Because the level of intended sedation may be exceeded, the practitioner must be sufficiently skilled to provide rescue should the child progress to a level of deep sedation. The practitioner must be trained in, and capable of providing, at the minimum, bag-valve-mask ventilation so as to be able to oxygenate a child who develops airway obstruction or apnea. Training in, and maintenance of, advanced pediatric airway skills is required; regular skills reinforcement is strongly encouraged.

#### Support Personnel

The use of moderate sedation shall include provision of a person, in addition to the practitioner, whose responsibility is to monitor appropriate physiologic parameters and to assist in any supportive or resuscitation measures, if required. This individual may also be responsible for assisting with interruptible patient-related tasks of short duration.<sup>44</sup> This individual

must be trained in and capable of providing pediatric basic life support. The support person shall have specific assignments in the event of an emergency and current knowledge of the emergency cart inventory. The practitioner and all ancillary personnel should participate in periodic reviews and practice drills of the facility's emergency protocol to ensure proper function of the equipment and coordination of staff roles in such emergencies.

#### *Monitoring and Documentation*

##### Baseline

Before administration of sedative medications, a baseline determination of vital signs shall be documented. For some children who are very upset or noncooperative, this may not be possible and a note should be written to document this happenstance.

##### During the procedure

The practitioner shall document the name, route, site, time of administration, and dosage of all drugs administered. There shall be continuous monitoring of oxygen saturation and heart rate and intermittent recording of respiratory rate and blood pressure; these should be recorded in a time-based record. Restraining devices should be checked to prevent airway obstruction or chest restriction. If a restraint device is used, a hand or foot should be kept exposed. The child's head position should be checked frequently to ensure airway patency. A functioning suction apparatus must be present.

##### After the procedure

The child who has received moderate sedation must be observed in a suitably equipped recovery facility (ie, the facility must have functioning suction apparatus as well as the capacity to deliver more than 90% oxygen and positive-pressure ventilation—eg, bag and mask with oxygen capacity as described previously). The patient's vital signs should be recorded at specific intervals. If the patient is not fully alert, oxygen saturation and heart rate monitoring shall be used continuously until appropriate discharge criteria are met (see Appendix A). Because sedation medications with a long half-life may delay the patient's complete return to baseline or pose the risk of resedation, some patients might benefit from a longer period of less-intense observation (eg, a step-down observation area where multiple patients can be observed simultaneously) before discharge from medical supervision (see also Documentation Before Sedation for instructions to families).<sup>45,103,131,132</sup> A recently described and simple evaluation tool may be the ability of the infant or child to remain awake for at least 20 minutes when placed in a quiet environment.<sup>100</sup> Patients who have received reversal agents, such as flumazenil or naloxone, will also require a longer period of observation, because the duration of the drugs administered may exceed the duration of the antagonist, which can lead to resedation.

#### **Deep Sedation**

Deep sedation is a drug-induced depression of consciousness during which patients cannot be easily aroused but respond purposefully after repeated verbal or painful stimulation (see Definition of Terms for This Report). The state and risks of deep sedation may be indistinguishable from those of general anesthesia.

#### *Personnel*

There must be 1 person available whose only responsibility is to constantly observe the patient's vital signs, airway patency, and adequacy of ventilation and to either administer drugs or direct their administration. At least 1 individual must be present who is trained in, and capable of, providing advanced pediatric life support, and who is skilled in airway management and cardiopulmonary resuscitation; training in pediatric advanced life support is required.

#### *Equipment*

In addition to the equipment previously cited for moderate sedation, an electrocardiographic monitor and a defibrillator for use in pediatric patients should be readily available.

#### *Vascular Access*

Patients receiving deep sedation should have an intravenous line placed at the start of the procedure or have a person skilled in establishing vascular access in pediatric patients immediately available.

#### *Monitoring and Documentation*

A competent individual shall observe the patient continuously. The monitoring shall include all parameters described for moderate sedation. Vital signs, including oxygen saturation and heart rate, must be documented at least every 5 minutes in a time-based record. The use of a precordial stethoscope or capnograph for patients difficult to observe (eg, during MRI, in a darkened room) to aid in monitoring adequacy of ventilation is encouraged.<sup>43</sup> The practitioner shall document the name, route, site, time of administration, and dosage of all drugs administered. The inspired concentrations of inhalation sedation agents and oxygen and the duration of administration shall be documented.

#### *Postsedation Care*

The facility and procedures followed for postsedation care shall conform to those described under "Moderate Sedation."

#### **Special Considerations**

##### **Local Anesthetic Agents**

All local anesthetic agents are cardiac depressants and may cause central nervous system excitation or depression. Particular attention should be paid to dosage in small children.<sup>64,66</sup> To ensure that the patient will not receive an excessive dose, the

maximum allowable safe dosage (eg, mg/kg) should be calculated before administration. There may be enhanced sedative effects when the highest recommended doses of local anesthetic drugs are used in combination with other sedatives or narcotics (see Tables 2 and 3 for limits and conversion tables of commonly used local anesthetics).<sup>64,144-157</sup> In general, when administering local anesthetic drugs, the practitioner should aspirate frequently so as to minimize the likelihood that the needle is in a blood vessel; lower doses should be used when injecting into vascular tissues.<sup>158</sup>

### Pulse Oximetry

The new generation of pulse oximeters is less susceptible to motion artifacts and may be more useful than older oximeters that do not contain the updated software.<sup>159-163</sup> Oximeters that change tone with changes in hemoglobin saturation provide immediate aural warning to everyone within hearing distance. It is essential that any oximeter probe is positioned properly; clip-on devices are prone to easy displacement, which may produce artifactual data (ie, under- or overestimation of oxygen saturation).<sup>141,142</sup>

### Capnography

Expired carbon dioxide monitoring is valuable to diagnose the simple presence or absence of respirations, airway obstruction, or respiratory depression, particularly in patients sedated in less-accessible locations, such as magnetic resonance imaging or computerized axial tomography devices or darkened rooms.<sup>47,49,50,143,164-173</sup> The use of expired carbon dioxide monitoring devices is encouraged for sedated children, particularly in situations where other means of assessing the adequacy of ventilation are limited. Several manufacturers have produced nasal cannulae that allow simultaneous delivery of oxygen and measurement of expired carbon dioxide values.<sup>164,165</sup> Although these devices can have a high degree of false-positive alarms, they are also very accurate for the detection of complete airway obstruction or apnea.<sup>166,168,173</sup>

### Adjuncts to Airway Management and Resuscitation

The vast majority of sedation complications can be managed with simple maneuvers, such as supplemental oxygen, opening the airway, suctioning, and bag-mask-valve ventilation. Occasionally, endotracheal intubation is required for more prolonged ventilatory support. In addition to standard endotracheal intubation techniques, a number of new devices are available for the management of patients with abnormal airway anatomy or airway obstruction. Examples include the laryngeal mask airway (LMA), the cuffed oropharyngeal airway, and a variety of kits to perform an emergency cricothyrotomy.

The largest clinical experience in pediatrics is with the LMA, which is available in a variety of sizes and can even be used in neonates. Use of the LMA is now being introduced into advanced airway training courses, and familiarity with insertion techniques can be life saving.<sup>174,175</sup> The LMA also can serve as a

bridge to secure airway management in children with anatomic airway abnormalities.<sup>176,177</sup> Practitioners are encouraged to gain experience with these techniques as they become incorporated into pediatric advanced life support courses.

An additional emergency device with which to become familiar is the intraosseous needle. Intraosseous needles also are available in several sizes and can be life saving in the rare situation when rapid establishment of intravenous access is not possible. Familiarity with the use of these adjuncts for the management of emergencies can be obtained by keeping current with resuscitation courses, such as Pediatric Advanced Life Support and Advanced Pediatric Life Support or other approved programs.

### Patient Simulators

Advances in technology, particularly patient simulators that allow a variety of programmed adverse events (eg, apnea, bronchospasm, laryngospasm), response to medical interventions, and printouts of physiologic parameters, are now available. The use of such devices is encouraged to better train medical professionals to respond more appropriately and effectively to rare events.<sup>178-180</sup>

### Monitoring During MRI

The powerful magnetic field and the generation of radiofrequency emissions necessitate the use of special equipment to provide continuous patient monitoring throughout the MRI scanning procedure. Pulse oximeters capable of continuous function during scanning should be used in any sedated or restrained pediatric patient. Thermal injuries can result if appropriate precautions are not taken; avoid coiling the oximeter wire and place the probe as far from the magnetic coil as possible to diminish the possibility of injury. Electrocardiogram monitoring during magnetic resonance imaging has been associated with thermal injury; special MRI-compatible ECG pads are essential to allow safe monitoring.<sup>181-184</sup> Expired carbon dioxide monitoring is strongly encouraged in this setting.

### Nitrous Oxide

Inhalation sedation/analgesia equipment that delivers nitrous oxide must have the capacity of delivering 100% and never less than 25% oxygen concentration at a flow rate appropriate to the size of the patient. Equipment that delivers variable ratios of nitrous oxide to oxygen and that has a delivery system that covers the mouth and nose must be used in conjunction with a calibrated and functional oxygen analyzer. All nitrous oxide-to-oxygen inhalation devices should be calibrated in accordance with appropriate state and local requirements. Consideration should be given to the National Institute of Occupational Safety and Health Standards for the scavenging of waste gases.<sup>185</sup> Newly constructed or reconstructed treatment facilities, especially those with piped-in nitrous oxide and oxygen, must have appropriate state or local inspections to certify proper function of inhalation sedation/analgesia systems before any delivery of patient care.



**Table 1. APPROPRIATE INTAKE OF FOOD AND LIQUIDS BEFORE ELECTIVE SEDATION\***

Ingested Material	Minimum Fasting Period (h)
Clear liquids: water, fruit juices without pulp, carbonated beverages, clear tea, black coffee	2
Breast milk	4
Infant formula	6
Nonhuman milk: because nonhuman milk is similar to solids in gastric emptying time, the amount ingested must be considered when determining an appropriate fasting period	6
Light meal: a light meal typically consists of toast and clear liquids. Meals that include fried or fatty foods or meat may prolong gastric emptying time. Both the amount and type of foods ingested must be considered when determining an appropriate fasting period.	6

\* American Society of Anesthesiologists. Practice Guidelines for Preoperative Fasting and the Use of Pharmacologic Agents to Reduce the Risk of Pulmonary Aspiration: Application to Healthy Patients Undergoing Elective Procedures. A Report of the American Society of Anesthesiologists. Available at: [www.asahq.org/publicationsAndServices/npoguide.html](http://www.asahq.org/publicationsAndServices/npoguide.html)

**Table 2. COMMONLY USED LOCAL ANESTHETIC AGENTS: DOSES, DURATION, AND CALCULATIONS\***

Local Anesthetic	Maximum Dose with Epinephrine (kg)†		Duration of Action (min)‡
	Medical	Dental	
<i>Esters</i>			
Procaine	10.0	6	60-90
Chloroprocaine	20.0	12	30-60
Tetracaine	1.5	1	180-600
<i>Amides</i>			
Lidocaine	7.0	4.4	90-200
Mepivacaine	7.0	4.4	120-240
Bupivacaine	3.0	1.3	180-600
Levobupivacaine	3.0	2	180-600
Ropivacaine	3.0	2	180-600
Articaine		7	60-230

\* Maximum recommended doses and duration of action. Note that lower doses should be used in very vascular areas.

† These are maximum doses of local anesthetics combined with epinephrine; lower doses are recommended when used without epinephrine. Doses of amides should be decreased by 30% in infants younger than 6 months. When lidocaine is being administered intravascularly (eg, during intravenous regional anesthesia), the dose should be decreased to 3 to 5 mg/kg; long-acting local anesthetic agents should not be used for intravenous regional anesthesia.

‡ Duration of action is dependent on concentration, total dose, and site of administration; use of epinephrine; and the patient's age.

**Table 3. LOCAL ANESTHETIC PERCENT CONCENTRATION: CONVERSION TO mg/mL**

Concentration (%)	mg/mL
3.0	30.0
2.5	25.0
2.0	20.0
1.0	10.0
0.5	5.0
0.25	2.5
0.125	1.25

Nitrous oxide in oxygen with varying concentrations has been successfully used for many years to provide analgesia for a variety of painful procedures in children.<sup>15,186-210</sup> The use of nitrous oxide for minimal sedation is defined as the administration of nitrous oxide (50% or less) with the balance as oxygen, without any other sedative, narcotic, or other depressant drug before or concurrent with the nitrous oxide to an otherwise healthy patient in ASA class I or II. The patient is able to maintain verbal communication throughout the procedure.

It should be noted that although local anesthetics have sedative properties, for purposes of this guideline, they are not considered sedatives in this circumstance. If nitrous oxide in oxygen is combined with other sedating medications, such as chloral hydrate, midazolam, or an opioid, or if nitrous oxide is used in concentrations greater than 50%, the likelihood for moderate or deep sedation increases.<sup>211,212</sup> In this situation, the clinician must be prepared to institute the guidelines for moderate or deep sedation as indicated by the patient's response.<sup>213</sup>

## Appendix A. Recommended Discharge Criteria

1. Cardiovascular function and airway patency are satisfactory and stable.
2. The patient is easily arousable, and protective reflexes are intact.
3. The patient can talk (if age appropriate).
4. The patient can sit up unaided (if age appropriate).
5. For a very young or handicapped child incapable of the usually expected responses, the presedation level of responsiveness or a level as close as possible to the normal level for that child should be achieved.
6. The state of hydration is adequate.

## Appendix C. Drugs\* That May Be Needed to Rescue a Sedated Patient<sup>44</sup>

Albuterol for inhalation  
 Ammonia spirits  
 Atropine  
 Diphenhydramine  
 Diazepam  
 Epinephrine (1:1000, 1:10 000)  
 Flumazenil  
 Glucose (25% or 50%)  
 Lidocaine (cardiac lidocaine, local infiltration)  
 Lorazepam  
 Methylprednisolone  
 Naloxone  
 Oxygen  
 Fosphenytoin  
 Racemic epinephrine  
 Rocuronium  
 Sodium bicarbonate  
 Succinylcholine

\* The choice of emergency drugs may vary according to individual or procedural needs.

## Appendix B. ASA Physical Status Classification

Class I	A normally healthy patient.
Class II	A patient with mild systemic disease (eg, controlled reactive airway disease).
Class III	A patient with severe systemic disease (eg, a child who is actively wheezing).
Class IV	A patient with severe systemic disease that is a constant threat to life (eg, a child with status asthmaticus).
Class V	A moribund patient who is not expected to survive without the operation (eg, a patient with severe cardiomyopathy requiring heart transplantation).

## Appendix D. Emergency Equipment\* That May Be Needed to Rescue a Sedated Patient †

### Intravenous Equipment

Assorted IV catheters (eg, 24-, 22-, 20-, 18-, 16-gauge)  
 Tourniquets  
 Alcohol wipes  
 Adhesive tape  
 Assorted syringes (eg, 1-, 3-, 5-, 10-mL)  
 IV tubing  
   Pediatric drip (60 drops/mL)  
   Pediatric burette  
   Adult drip (10 drops/mL)  
   Extension tubing  
   3-way stopcocks  
 IV fluid  
   Lactated Ringer solution  
   Normal saline solution  
   D<sub>5</sub> 0.25 normal saline solution  
 Pediatric IV boards  
 Assorted IV needles (eg, 25-, 22-, 20-, and 18-gauge)  
 Intraosseous bone marrow needle  
 Sterile gauze pads

\* The choice of emergency equipment may vary according to individual or procedural needs.

† The practitioner is referred to the SOAPME acronym described in the text in preparation for sedating a child for a procedure.

### Airway Management Equipment

Face masks (Infant, child, small adult, medium adult, large adult)  
 Breathing bag and valve set  
 Oropharyngeal airways (Infant, child, small adult, medium adult, large adult)  
 Nasopharyngeal airways (Small, medium, large)  
 Laryngeal mask airways (1, 1.5, 2, 2.5, 3, 4, and 5)  
 Laryngoscope handles (with extra batteries)  
 Laryngoscope blades (with extra light bulbs)  
   Straight (Miller) No. 1, 2, and 3  
   Curved (Macintosh) No. 2 and 3  
 Endotracheal tubes (2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, and 6.0 uncuffed and 6.0, 7.0, and 8.0 cuffed)  
 Stylettes (appropriate sizes for endotracheal tubes)  
 Surgical lubricant  
 Suction catheters (appropriate sizes for endotracheal tubes)  
 Yankauer-type suction  
 Nasogastric tubes  
 Nebulizer with medication kits  
 Gloves (sterile and nonsterile, latex free)

## References

1. Milnes AR. Intravenous procedural sedation: An alternative to general anesthesia in the treatment of early childhood caries. *J Can Dent Assoc* 2003;69:298-302.
2. Law AK, Ng DK, Chan KK. Use of intramuscular ketamine for endoscopy sedation in children. *Pediatr Int* 2003;45:180-5.
3. Rothermel LK. Newer pharmacologic agents for procedural sedation of children in the emergency department—Etomidate and propofol. *Curr Opin Pediatr* 2003;15:200-3.
4. Flood RG, Krauss B. Procedural sedation and analgesia for children in the emergency department. *Emerg Med Clin North Am* 2003;21:121-39.
5. Jaggar SI, Haxby E. Sedation, anaesthesia and monitoring for bronchoscopy. *Pediatr Respir Rev* 2002;3:321-7.
6. de Blic J, Marchac V, Scheinmann P. Complications of flexible bronchoscopy in children: Prospective study of 1,328 procedures. *Eur Respir J* 2002;20:1271-6.
7. Mason KP, Michna E, DiNardo JA, et al. Evolution of a protocol for ketamine-induced sedation as an alternative to general anesthesia for interventional radiologic procedures in pediatric patients. *Radiology* 2002;225:457-65.
8. Houpt M. Project USAP 2000—Use of sedative agents by pediatric dentists: A 15-year follow-up survey. *Pediatr Dent* 2002;24:289-94.
9. Vinson DR, Bradbury DR. Etomidate for procedural sedation in emergency medicine. *Ann Emerg Med* 2002;39:592-8.
10. Everitt IJ, Barnett P. Comparison of two benzodiazepines used for sedation of children undergoing suturing of a laceration in an emergency department. *Pediatr Emerg Care* 2002;18:72-4.
11. Karian VE, Burrows PE, Zurakowski D, Connor L, Poznauskis L, Park MK. The development of a pediatric radiology sedation program. *Pediatr Radiol* 2002;32:348-53.
12. Kaplan RF, Yang CI. Sedation and analgesia in pediatric patients for procedures outside the operating room. *Anesthesiol Clin North America* 2002;20:181-94.
13. Wheeler DS, Jensen RA, Poss WB. A randomized, blinded comparison of chloral hydrate and midazolam sedation in children undergoing echocardiography. *Clin Pediatr* 2001;40:381-7.
14. Hain RD, Campbell C. Invasive procedures carried out in conscious children: Contrast between North American and European paediatric oncology centres. *Arch Dis Child* 2001;85:12-5.
15. Kennedy RM, Luhmann JD. Pharmacological management of pain and anxiety during emergency procedures in children. *Pediatr Drugs* 2001;3:337-54.
16. Kanagasundaram SA, Lane LJ, Cavalletto BP, Keneally JP, Cooper MG. Efficacy and safety of nitrous oxide in alleviating pain and anxiety during painful procedures. *Arch Dis Child* 2001;84:492-5.

17. Younge PA, Kendall JM. Sedation for children requiring wound repair: A randomised controlled double blind comparison of oral midazolam and oral ketamine. *Emerg Med J* 2001;18:30-3.
18. Ljungman G, Gordh T, Sorensen S, Kreuger A. Lumbar puncture in pediatric oncology: Conscious sedation vs. general anesthesia. *Med Pediatr Oncol* 2001;36:372-9.
19. Poe SS, Nolan MT, Dang D, et al. Ensuring safety of patients receiving sedation for procedures: Evaluation of clinical practice guidelines. *Jt Comm J Qual Improv* 2001;27:28-41.
20. D'Agostino J, Terndrup TE. Chloral hydrate versus midazolam for sedation of children for neuroimaging: A randomized clinical trial. *Pediatr Emerg Care* 2000;16:1-4.
21. Green SM, Kuppermann N, Rothrock SG, Hummel CB, Ho M. Predictors of adverse events with intramuscular ketamine sedation in children. *Ann Emerg Med* 2000;35:35-42.
22. Pena BM, Krauss B. Adverse events of procedural sedation and analgesia in a pediatric emergency department. *Ann Emerg Med* 1999;34:483-91.
23. Hopkins KL, Davis PC, Sanders CL, Churchill LH. Sedation for pediatric imaging studies. *Neuroimaging Clin North Am* 1999;9:1-10.
24. Bauman LA, Kish I, Baumann RC, Politis GD. Pediatric sedation with analgesia. *Am J Emerg Med* 1999;17:1-3.
25. Bhatt-Mehta V, Rosen DA. Sedation in children: Current concepts. *Pharmacotherapy* 1998;18:790-807.
26. Morton NS, Oomen CJ. Development of a selection and monitoring protocol for safe sedation of children. *Pediatr Anaesth* 1998;8:65-8.
27. Murphy MS. Sedation for invasive procedures in paediatrics. *Arch Dis Child* 1997;77:281-4.
28. Webb MD, Moore PA. Sedation for pediatric dental patients. *Dent Clin North Am* 2002;46:803-14.
29. Malviya S, Voepel-Lewis T, Tait AR, Merkel S. Sedation/analgesia for diagnostic and therapeutic procedures in children. *J Perianesth Nurs* 2000;15:415-22.
30. Zempsky WT, Schechter NL. Office-based pain management. The 15-minute consultation. *Pediatr Clin North Am* 2000;47:601-15.
31. Kennedy RM, Luhmann JD. The "ouchless emergency department." Getting closer: Advances in decreasing distress during painful procedures in the emergency department. *Pediatr Clin North Am* 1999;46:1215-47.
32. Rodriguez E, Jordan R. Contemporary trends in pediatric sedation and analgesia. *Emerg Med Clin North Am* 2002;20:199-222.
33. Ruess L, O'Connor SC, Mikita CP, Creamer KM. Sedation for pediatric diagnostic imaging: use of pediatric and nursing resources as an alternative to a radiology department sedation team. *Pediatr Radiol* 2002;32:505-10.
34. Weiss S. Sedation of pediatric patients for nuclear medicine procedures. *Semin Nucl Med* 1993;23:190-198.
35. Wilson S. Pharmacologic behavior management for pediatric dental treatment. *Pediatr Clin North Am* 2000;47:1159-75.
36. McCarty EC, Mencio GA, Green NE. Anesthesia and analgesia for the ambulatory management of fractures in children. *J Am Acad Orthop Surg* 1999;7:81-91.
37. Egelhoff JC, Ball WS Jr, Koch BL, Parks TD. Safety and efficacy of sedation in children using a structured sedation program. *AJR Am J Roentgenol* 1997;168:1259-62.
38. American Academy of Pediatrics Committee on Drugs and Section on Anesthesiology. Guidelines for the elective use of conscious sedation, deep sedation, and general anesthesia in pediatric patients. *Pediatrics* 1985;76:317-21.
39. American Academy of Pediatric Dentistry. Guidelines for the elective use of conscious sedation, deep sedation, and general anesthesia in pediatric patients. *ASDC J Dent Child* 1986;53:21-2.
40. American Academy of Pediatrics, Committee on Drugs. Guidelines for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures. *Pediatrics* 1992;89:1110-5.
41. American Academy of Pediatrics, Committee on Drugs. Guidelines for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures: Addendum. *Pediatrics* 2002;110:836-8.
42. American Academy of Pediatric Dentistry. Guidelines on the Elective Use of Minimal, Moderate, and Deep Sedation and General Anesthesia for Pediatric Dental Patients. Chicago, Ill. *Pediatr Dent* 2004;26 (suppl):95-105.
43. Joint Commission on Accreditation of Healthcare Organizations. Comprehensive Accreditation Manual for Hospitals: The Official Handbook by the JCAHO. Oakbrook Terrace, Ill: Joint Commission on Accreditation of Healthcare Organizations; 2005.
44. American Society of Anesthesiologists, Task Force on Sedation and Analgesia by Non-Anesthesiologists. Practice guidelines for sedation and analgesia by non-anesthesiologists. *Anesthesiology* 2002;96:1004-17.
45. Coté CJ, Karl HW, Notterman DA, Weinberg JA, McCloskey C. Adverse sedation events in pediatrics: Analysis of medications used for sedation. *Pediatrics* 2000;106:633-44.
46. Coté CJ, Notterman DA, Karl HW, Weinberg JA, McCloskey C. Adverse sedation events in pediatrics: A critical incident analysis of contributory factors. *Pediatrics* 2000;105:805-14.
47. Kim G, Green SM, Denmark TK, Krauss B. Ventilatory response during dissociative sedation in children—A pilot study. *Acad Emerg Med* 2003;10:140-5.

48. Coté CJ. Sedation for the pediatric patient. A review. *Pediatr Clin North Am* 1994;41:31-58.
49. Mason KP, Burrows PE, Dorsey MM, Zurakowski D, Krauss B. Accuracy of capnography with a 30 foot nasal cannula for monitoring respiratory rate and end-tidal CO<sub>2</sub> in children. *J Clin Monit Comput* 2000; 16:259-62.
50. McQuillen KK, Steele DW. Capnography during sedation/analgesia in the pediatric emergency department. *Pediatr Emerg Care* 2000;16:401-404.
51. Malviya S, Voepel-Lewis T, Tait AR. Adverse events and risk factors associated with the sedation of children by nonanesthesiologists. *Anesth Analg* 1997;85:1207-13.
52. Coté CJ, Rolf N, Liu LM, et al. A single-blind study of combined pulse oximetry and capnography in children. *Anesthesiology* 1991;74:980-7.
53. Scottish Intercollegiate Guidelines Network. Safe Sedation of Children Undergoing Diagnostic and Therapeutic Procedures. A National Clinical Guideline. Edinburgh, Scotland: Scottish Intercollegiate Guidelines Network; 2003. Available at: [www.sign.ac.uk/pdf/sign58.pdf](http://www.sign.ac.uk/pdf/sign58.pdf). Accessed February 7, 2006.
54. Hoffman GM, Nowakowski R, Troshynski TJ, Berens RJ, Weisman SJ. Risk reduction in pediatric procedural sedation by application of an American Academy of Pediatrics/American Society of Anesthesiologists process model. *Pediatrics* 2002;109:236-43.
55. Krauss B. Management of acute pain and anxiety in children undergoing procedures in the emergency department. *Pediatr Emerg Care* 2001;17:115-22.
56. Dial S, Silver P, Bock K, Sagy M. Pediatric sedation for procedures titrated to a desired degree of immobility results in unpredictable depth of sedation. *Pediatr Emerg Care* 2001;17:414-20.
57. Maxwell LG, Yaster M. The myth of conscious sedation. *Arch Pediatr Adolesc Med*. 1996;150:665-7.
58. Coté CJ. "Conscious sedation": Time for this oxymoron to go away! *J Pediatr* 2001;139:15-7.
59. Motas D, McDermott NB, VanSickle T, Friesen RH. Depth of consciousness and deep sedation attained in children as administered by nonanaesthesiologists in a children's hospital. *Pediatr Anaesth* 2004;14:256-60.
60. Nahata MC, Clotz MA, Krogg EA. Adverse effects of meperidine, promethazine, and chlorpromazine for sedation in pediatric patients. *Clin Pediatr* 1985; 24:558-60.
61. Brown ET, Corbett SW, Green SM. Iatrogenic cardio pulmonary arrest during pediatric sedation with meperidine, promethazine, and chlorpromazine. *Pediatr Emerg Care* 2001;17:351-3.
62. Benusis KP, Kapaun D, Furnam LJ. Respiratory depression in a child following meperidine, promethazine, and chlorpromazine premedication: Report of case. *J Dent Child* 1979;46:50-3.
63. Garriott JC, Di Maio VJ. Death in the dental chair: Three drug fatalities in dental patients. *J Toxicol Clin Toxicol* 1982;19:987-995.
64. Goodson JM, Moore PA. Life-threatening reactions after pedodontic sedation: An assessment of narcotic, local anesthetic, and antiemetic drug interaction. *J Am Dent Assoc* 1983;107:239-45.
65. Jastak JT, Pallasch T. Death after chloral hydrate sedation: Report of case. *J Am Dent Assoc* 1988;116:345-8.
66. Jastak JT, Peskin RM. Major morbidity or mortality from office anesthetic procedures: A closed-claim analysis of 13 cases. *Anesth Prog* 1991;38:39-44.
67. Kaufman E, Jastak JT. Sedation for outpatient dental procedures. *Compend Contin Educ Dent* 1995;16:462, 464, 466.
68. Wilson S. Pharmacological management of the pediatric dental patient. *Pediatr Dent* 2004;26:131-6.
69. Sams DR, Thornton JB, Wright JT. The assessment of two oral sedation drug regimens in pediatric dental patients. *J Dent Child* 1992;59:306-12.
70. Malviya S, Voepel-Lewis T, Tait AR, Merkel S, Tremper K, Naughton N. Depth of sedation in children undergoing computed tomography: Validity and reliability of the University of Michigan Sedation Scale (UMSS). *Br J Anaesth* 2002;88:241-5.
71. Newton JT, Shah S, Patel H, Sturmey P. Non-pharmacological approaches to behaviour management in children. *Dent Update* 2003;30:194-9.
72. Peretz B, Bimstein E. The use of imagery suggestions during administration of local anesthetic in pediatric dental patients. *ASDC J Dent Child* 2000;67:263-7.
73. Iserson KV. Hypnosis for pediatric fracture reduction. *J Emerg Med* 1999;17:53-6.
74. Rusy LM, Weisman SJ. Complementary therapies for acute pediatric pain management. *Pediatr Clin North Am* 2000;47:589-99.
75. Langley P. Guided imagery: A review of effectiveness in the care of children. *Pediatr Nurs* 1999;11:18-21.
76. Ott MJ. Imagine the possibilities! Guided imagery with toddlers and pre-schoolers. *Pediatr Nurs* 1996; 22:34-8.
77. Singer AJ, Stark MJ. LET versus EMLA for pretreating lacerations: A randomized trial. *Acad Emerg Med* 2001; 8:223-30.
78. Taddio A, Gurguis MG, Koren G. Lidocaine-prilocaine cream versus tetracaine gel for procedural pain in children. *Ann Pharmacother* 2002;36:687-92.
79. Eichenfield LF, Funk A, Fallon-Friedlander S, Cunningham BB. A clinical study to evaluate the efficacy of ELA-Max (4% liposomal lidocaine) as compared with eutectic mixture of local anesthetics cream for pain reduction of venipuncture in children. *Pediatrics* 2002;109:1093-9.
80. Shaw AJ, Welbury RR. The use of hypnosis in a sedation clinic for dental extractions in children: Report of 20 cases. *ASDC J Dent Child* 1996;63:418-20.

81. Aitken JC, Wilson S, Coury D, Moursi AM. The effect of music distraction on pain, anxiety and behavior in pediatric dental patients. *Pediatr Dent* 2002;24:114-8.
82. Domino KB. Office-based anesthesia: Lessons learned from the closed claims project. *ASA Newsletter* 2001; 65:9-11,15.
83. American Heart Association. Pediatric Advanced Life Support Provider Manual. Dallas, Tx: American Heart Association; 2002.
84. American Academy of Pediatrics, American College of Emergency Physicians. Advanced Pediatric Life Support. 4th ed. Fuchs S, Gausche-Hill M, Yamoto L, eds. Boston, Ma: Jones and Bartlett Publishers; 2004.
85. Wheeler M. Management strategies for the difficult pediatric airway. In: Riaz J, ed. *Anesthesiology Clinics of North America*. Philadelphia, Pa: WB Saunders Co; 1998:743-61.
86. Sullivan KJ, Kissoon N. Securing the child's airway in the emergency department. *Pediatr Emerg Care* 2002; 18:108-21.
87. Levy RJ, Helfaer MA. Pediatric airway issues. *Crit Care Clin* 2000;16:489-504.
88. Krauss B, Green SM. Procedural sedation and analgesia in children. *Lancet* 2006;367:766-80.
89. Krauss B, Green SM. Sedation and analgesia for procedures in children. *N Engl J Med*. 2000;342:938-45.
90. Ferrari LR. *Anesthesia and Pain Management for the Pediatrician*. Baltimore, Md: John Hopkins University Press; 1999.
91. Malviya S, Naughton NN, Tremper KT, eds. *Sedation Analgesia for Diagnostic and Therapeutic Procedures*. Totowa, NJ: Humana Press; 2001.
92. Yaster M, Krane EJ, Kaplan RF, Coté CJ, Lappe DG, eds. *Pediatric Pain Management and Sedation Handbook*. St. Louis, Mo: CV Mosby; 1997.
93. Cravero JP, Blike GT. Review of pediatric sedation. *Anesth Analg* 2004;99:1355-64.
94. *Standards for Basic Intra-operative Monitoring*. Chicago, Ill: American Society of Anesthesiologists; 1991.
95. Mace SE, Barata IA, Cravero JP, et al. Clinical policy: Evidence-based approach to pharmacologic agents used in pediatric sedation and analgesia in the emergency department. *Ann Emerg Med* 2004;44:342-77.
96. Deshpande JK, Tobias JD, eds. *The Pediatric Pain Handbook*. St. Louis, Mo: Mosby; 1996.
97. Alcaino EA. Conscious sedation in paediatric dentistry: Current philosophies and techniques. *Ann R Australas Coll Dent Surg* 2000;15:206-10.
98. Mitchell AA, Louik C, Lacouture P, Slone D, Goldman P, Shapiro S. Risks to children from computed tomographic scan premedication. *JAMA* 1982;247:2385-8.
99. Pershad J, Palmisano P, Nichols M. Chloral hydrate: the good and the bad. *Pediatr Emerg Care* 1999;15:432-5.
100. Malviya S, Voepel-Lewis T, Ludomirsky A, Marshall J, Tait AR. Can we improve the assessment of discharge readiness? A comparative study of observational and objective measures of depth of sedation in children. *Anesthesiology* 2004;100:218-24.
101. American Academy of Pediatrics, Committee on Injury and Poison Prevention. Transporting children with special health care needs. *Pediatrics* 1999;104:988-92.
102. American Academy of Pediatrics, Committee on Bioethics. Informed consent, parental permission, and assent in pediatric practice. *Pediatrics* 1995;95:314-7.
103. Malviya S, Voepel-Lewis T, Prochaska G, Tait AR. Prolonged recovery and delayed side effects of sedation for diagnostic imaging studies in children. *Pediatrics* 2000;105(3):e42.
104. Babl FE, Puspitadewi A, Barnett P, Oakley E, Spicer M. Preprocedural fasting state and adverse events in children receiving nitrous oxide for procedural sedation and analgesia. *Pediatr Emerg Care* 2005;21:736-43.
105. Roback MG, Bajaj L, Wathen JE, Bothner J. Preprocedural fasting and adverse events in procedural sedation and analgesia in a pediatric emergency department: Are they related? *Ann Emerg Med* 2004;44:454-9.
106. Agrawal D, Manzi SF, Gupta R, Krauss B. Preprocedural fasting state and adverse events in children undergoing procedural sedation and analgesia in a pediatric emergency department. *Ann Emerg Med* 2003; 42:636-46.
107. Green SM. Fasting is a consideration—not a necessity—for emergency department procedural sedation and analgesia. *Ann Emerg Med* 2003;42:647-50.
108. Green SM, Krauss B. Pulmonary aspiration risk during emergency department procedural sedation—An examination of the role of fasting and sedation depth. *Acad Emerg Med* 2002;9:35-42.
109. Treston G. Prolonged pre-procedure fasting time is unnecessary when using titrated intravenous ketamine for paediatric procedural sedation. *Emerg Med Australas* 2004;16:145-50.
110. Green SM, Krauss B. Ketamine is a safe, effective, and appropriate technique for emergency department paediatric procedural sedation. *Emerg Med J* 2004;21:271-2.
111. Gorski JC, Huang SM, Pinto A, et al. The effect of echinacea (*Echinacea purpurea* root) on cytochrome P450 activity in vivo. *Clin Pharmacol Ther* 2004;75:89-100.
112. Hall SD, Wang Z, Huang SM, et al. The interaction between St John's wort and an oral contraceptive. *Clin Pharmacol Ther* 2003;74:525-35.
113. Markowitz JS, Donovan JL, DeVane CL, et al. Effect of St John's wort on drug metabolism by induction of cytochrome P450 3A4 enzyme. *JAMA* 2003;290:1500-4.

114. Spinella M. Herbal medicines and epilepsy: The potential for benefit and adverse effects. *Epilepsy Behav* 2001; 2:524-32.
115. Wang Z, Gorski JC, Hamman MA, Huang SM, Lesko LJ, Hall SD. The effects of St John's wort (*Hypericum perforatum*) on human cytochrome P450 activity. *Clin Pharmacol Ther* 2001;70:317-26.
116. Xie HG, Kim RB. St John's wort-associated drug interactions: Short-term inhibition and long-term induction? *Clin Pharmacol Ther* 2005 Jul;78(1):19-24.
117. Ang-Lee MK, Moss J, Yuan CS. Herbal medicines and perioperative care. *JAMA* 2001;286:208-16.
118. Abebe W. Herbal medication: Potential for adverse interactions with analgesic drugs. *J Clin Pharm Ther* 2002; 27:391-401.
119. von Rosensteil NA, Adam D. Macrolide antibacterials. Drug interactions of clinical significance. *Drug Saf* 1995; 13:105-22.
120. Hiller A, Olkkola KT, Isohanni P, Saarnivaara L. Unconsciousness associated with midazolam and erythromycin. *Br J Anaesth* 1990;65:826-8.
121. Mattila MJ, Idanpaan-Heikkilä JJ, Tornwall M, Vanakoski J. Oral single doses of erythromycin and roxithromycin may increase the effects of midazolam on human performance. *Pharmacol Toxicol* 1993;73:180-5.
122. Olkkola KT, Aranko K, Luurila H, et al. A potentially hazardous interaction between erythromycin and midazolam. *Clin Pharmacol Ther* 1993;53:298-05.
123. Flockhart DA, Oesterheld JR. Cytochrome P450-mediated drug interactions. *Child Adolesc Psychiatr Clin N Am* 2000;9:43-76.
124. Yuan R, Flockhart DA, Balian JD. Pharmacokinetic and pharmacodynamic consequences of metabolism-based drug interactions with alprazolam, midazolam, and triazolam. *J Clin Pharmacol* 1999;39:1109-25.
125. Young B. Review: Mixing new cocktails: Drug interactions in antiretroviral regimens. *AIDS Patient Care STDS* 2005;19:286-97.
126. Wilkinson GR. Drug metabolism and variability among patients in drug response. *N Engl J Med* 2005; 352:2211-21.
127. American Academy of Pediatrics, Section on Pediatric Pulmonology, Subcommittee on Obstructive Sleep Apnea Syndrome. Clinical practice guideline: Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics* 2002;109:704-12.
128. Schechter MS. Technical report: Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics* 2002;109(4):e69.
129. Litman RS, Kottra JA, Berkowitz RJ, Ward DS. Upper airway obstruction during midazolam/nitrous oxide sedation in children with enlarged tonsils. *Pediatr Dent* 1998; 20:318-20.
130. Fishbaugh DF, Wilson S, Preisch JW, Weaver JM II. Relationship of tonsil size on an airway blockage maneuver in children during sedation. *Pediatr Dent* 1997;19:277-81.
131. Mayers DJ, Hindmarsh KW, Sankaran K, Gorecki DK, Kasian GF. Chloral hydrate disposition following single-dose administration to critically ill neonates and children. *Dev Pharm & Ther* 1991;16:71-77.
132. Terndrup TE, Dire DJ, Madden CM, Davis H, Cantor RM, Gavula DP. A prospective analysis of intramuscular meperidine, promethazine, and chlorpromazine in pediatric emergency department patients. *Ann Emerg Med* 1991;20:31-5.
133. Coté CJ. Discharge criteria for children sedated by non-anesthesiologists: Is "safe" really safe enough? *Anesthesiology* 2004;100:207-9.
134. Macnab AJ, Levine M, Glick N, Susak L, Baker-Brown G. A research tool for measurement of recovery from sedation: The Vancouver Sedative Recovery Scale. *J Pediatr Surg* 1991;26:1263-7.
135. Chernik DA, Gillings D, Laine H, et al. Validity and reliability of the Observer's Assessment of Alertness/Sedation Scale: Study with intravenous midazolam. *J Clin Psychopharmacol* 1990;10:244-51.
136. Bagian JP, Lee C, Gosbee J, et al. Developing and deploying a patient safety program in a large health care delivery system: You can't fix what you don't know about. *Jt Comm J Qual Improv* 2001;27:522-532.
137. May T, Aulisio MP. Medical malpractice, mistake prevention, and compensation. *Kennedy Inst Ethics J* 2001;11:135-146.
138. Kazandjian VA. When you hear hoofs, think horses, not zebras: An evidence-based model of health care accountability. *J Eval Clin Pract* 2002;8:205-13.
139. Connor M, Ponte PR, Conway J. Multidisciplinary approaches to reducing error and risk in a patient care setting. *Crit Care Nurs Clin North Am* 2002;14:359-67.
140. Gosbee J. Human factors engineering and patient safety. *Qual Saf Health Care* 2002;11:352-4.
141. Barker SJ, Hyatt J, Shah NK, Kao YJ. The effect of sensor malpositioning on pulse oximeter accuracy during hypoxemia. *Anesthesiology* 1993;79:248-54.
142. Kelleher JF, Ruff RH. The penumbra effect: Vasomotion-dependent pulse oximeter artifact due to probe malposition. *Anesthesiology* 1989;71:787-91.
143. Hart LS, Berns SD, Houck CS, Boenning DA. The value of end-tidal CO<sub>2</sub> monitoring when comparing three methods of conscious sedation for children undergoing painful procedures in the emergency department. *Pediatr Emerg Care* 1997;13:189-93.
144. Aubuchon RW. Sedation liabilities in pedodontics (abstr). *Pediatr Dent* 1982;4:171-80.

145. Fitzmaurice LS, Wasserman GS, Knapp JF, Roberts DK, Waeckerle JF, Fox M. TAC use and absorption of cocaine in a pediatric emergency department. *Ann Emerg Med* 1990;19:515-8.
146. Tipton GA, DeWitt CW, Eisenstein SJ. Topical TAC (tetracaine, adrenaline, cocaine) solution for local anesthesia in children: Prescribing inconsistency and acute toxicity. *South Med J* 1989;82:1344-6.
147. Gunter JB. Benefit and risks of local anesthetics in infants and children. *Paediatr Drugs* 2002;4:649-72.
148. Resar LM, Helfaer MA. Recurrent seizures in a neonate after lidocaine administration. *J Perinatol* 1998;18:193-5.
149. Yagiela JA, Neidle EA, Dowd FJ. Local anesthetics. In: *Pharmacology and Therapeutics for Dentistry*. Elsevier Health Sciences 2004;251-70.
150. Haas DA. An update on local anesthetics in dentistry. *J Can Dent Assoc* 2002;68:546-51.
151. Malamed SF. Local anesthetic considerations in dental specialties. In: *Handbook of Local Anesthesia*, 5th ed. St. Louis, Mo: Mosby; 2004;269, 274-5.
152. Malamed SF. The needle. *Handbook of Local Anesthesia*, 5th ed. St. Louis, Mo: Mosby; 2004;99-107.
153. Malamed SF. Clinical action of specific agents. *Handbook of Local Anesthesia*, 5th ed. St. Louis, Mo: Mosby; 2004;55-81.
154. Ram D, Amir E. Comparison of articaine 4% and lidocaine 2% in pediatric dental patients. *Intl J Pediatric Dent* 2006;16:252-6.
155. Jakobs W, Ladwig B, Cichon P, Ortel R, Kirch W. Serum levels of articaine 2% and 4% in children. *Anesth Prog* 1995;42:113-5.
156. Wright GZ, Weinberger SJ, Friedman CS, Plotzke OB. Use of articaine local anesthesia in children under 4 years of age—A retrospective report. *Anesth Report* 1989;36:268-71.
157. Malamed SF, Gagnon S, Leblanc D. A comparison between articaine HCl and lidocaine HCl in pediatric dental patients. *Pediatr Dent* 2000;22:307-11.
158. American Academy of Pediatric Dentistry. Guideline on appropriate use of local anesthesia for pediatric dental patients. *Pediatr Dent* 2005;27(Suppl):101-6.
159. Next-generation pulse oximetry. *Health Devices* 2003;32:49-103.
160. Barker SJ. "Motion-resistant" pulse oximetry: A comparison of new and old models. *Anesth Analg* 2002;95:967-72.
161. Malviya S, Reynolds PI, Voepel-Lewis T, et al. False alarms and sensitivity of conventional pulse oximetry versus the Masimo SET technology in the pediatric postanesthesia care unit. *Anesth Analg* 2000;90:1336-40.
162. Barker SJ, Shah NK. Effects of motion on the performance of pulse oximeters in volunteers. *Anesthesiology* 1996;85:774-81.
163. Barker SJ, Shah NK. The effects of motion on the performance of pulse oximeters in volunteers (revised publication). *Anesthesiology* 1997;86:101-8.
164. Colman Y, Krauss B. Microstream capnography technology: A new approach to an old problem. *J Clin Monit Comput* 1999;15:403-9.
165. Wright SW. Conscious sedation in the emergency department: The value of capnography and pulse oximetry. *Ann Emerg Med* 1992;21:551-5.
166. Croswell RJ, Dilley DC, Lucas WJ, Vann WF Jr. A comparison of conventional versus electronic monitoring of sedated pediatric dental patients. *Pediatr Dent* 1995;17:332-9.
167. Tobias JD. End-tidal carbon dioxide monitoring during sedation with a combination of midazolam and ketamine for children undergoing painful, invasive procedures. *Pediatr Emerg Care* 1999;15:173-5.
168. Primosch RE, Buzzi IM, Jerrell G. Monitoring pediatric dental patients with nasal mask capnography. *Pediatr Dent* 2000;22:120-4.
169. Roelofse J. Conscious sedation: Making our treatment options safe and sound. *SADJ* 2000;55:273-6.
170. Wilson S, Creedon RL, George M, Troutman K. A history of sedation guidelines: Where we are headed in the future. *Pediatr Dent* 1996;18:194-9.
171. Miner JR, Heegaard W, Plummer D. End-tidal carbon dioxide monitoring during procedural sedation. *Acad Emerg Med* 2002;9:275-280.
172. Vascello LA, Bowe EA. A case for capnographic monitoring as a standard of care. *J Oral Maxillofac Surg* 1999;57:1342-7.
173. Iwasaki J, Vann WF Jr, Dilley DC, Anderson JA. An investigation of capnography and pulse oximetry as monitors of pediatric patients sedated for dental treatment. *Pediatr Dent* 1989;11:111-7.
174. Berry AM, Brimacombe JR, Verghese C. The laryngeal mask airway in emergency medicine, neonatal resuscitation, and intensive care medicine. *Int Anesthesiol Clin* 1998;36:91-109.
175. Patterson MD. Resuscitation update for the pediatrician. *Pediatr Clin North Am* 1999;46:1285-303.
176. Selim M, Mowafi H, Al Chamdi A, Adu-Gyamfi Y. Intubation via LMA in pediatric patients with difficult airways. *Can J Anaesth* 1999;46:891-3.
177. Munro HM, Butler PJ, Washington EJ, Freeman-Sheldon (whistling face) syndrome. *Anaesthetic and airway management*. *Paediatr Anaesth* 1997;7:345-8.
178. Rowe R, Cohen RA. An evaluation of a virtual reality airway simulator. *Anesth Analg* 2002;95:62-6.
179. Medina LS, Racadio JM, Schwid HA. Computers in radiology. The sedation, analgesia, and contrast media computerized simulator: A new approach to train and evaluate radiologists' responses to critical incidents. *Pediatr Radiol* 2000;30:299-305.



180. Blike G, Cravero J, Nelson E. Same patients, same critical events-different systems of care, different outcomes: Description of a human factors approach aimed at improving the efficacy and safety of sedation/analgesia care. *Qual Manag Health Care* 2001;10:17-36.
181. Kanal E, Shellock FG, Talagala L. Safety considerations in MR imaging. *Radiology* 1990;176:593-606.
182. Shellock FG, Kanal E. Burns associated with the use of monitoring equipment during MR procedures. *J Magn Reson Imaging* 1996;6:271-2.
183. Shellock FG. Magnetic resonance safety update 2002: Implants and devices. *J Magn Reson Imaging* 2002;16:485-96.
184. Dempsey MF, Condon B, Hadley DM. MRI safety review. *Semin Ultrasound CT MR*. 2002;23:392-401
185. National Institute for Occupational Safety and Health (NIOSH). Criteria for a Recommended Standard: Occupational Exposure to Waste Anesthetic Gases and Vapors. Cincinnati, Oh. Publication 77-140. 1977. RefType: Statute.
186. O'Sullivan I, Bengner J. Nitrous oxide in emergency medicine. *Emerg Med J* 2003;20:214-7.
187. Kennedy RM, Luhmann JD, Luhmann SJ. Emergency department management of pain and anxiety related to orthopedic fracture care: A guide to analgesic techniques and procedural sedation in children. *Paediatr Drugs* 2004;6:11-31.
188. Frampton A, Browne GJ, Lam LT, Cooper MG, Lane LG. Nurse administered relative analgesia using high concentration nitrous oxide to facilitate minor procedures in children in an emergency department. *Emerg Med J* 2003;20:410-3.
189. Everitt I, Younge P, Barnett P. Paediatric sedation in emergency department: What is our practice? *Emerg Med (Fremantle)* 2002;14:62-6.
190. Krauss B. Continuous-flow nitrous oxide: Searching for the ideal procedural anxiolytic for toddlers. *Ann Emerg Med* 2001;37:61-2.
191. Otley CC, Nguyen TH. Conscious sedation of pediatric patients with combination oral benzodiazepines and inhaled nitrous oxide. *Dermatol Surg* 2000;26:1041-4.
192. Luhmann JD, Kennedy RM, Jaffe DM, McAllister JD. Continuous-flow delivery of nitrous oxide and oxygen: A safe and cost-effective technique for inhalation analgesia and sedation of pediatric patients. *Pediatr Emerg Care* 1999;15:388-92.
193. Burton JH, Auble TE, Fuchs SM. Effectiveness of 50% nitrous oxide/50% oxygen during laceration repair in children. *Acad Emerg Med* 1998;5:112-7.
194. Gregory PR, Sullivan JA. Nitrous oxide compared with intravenous regional anesthesia in pediatric forearm fracture manipulation. *J Pediatr Orthop* 1996;16:187-91.
195. Hennrikus WL, Shin AY, Klingelberger CE. Self-administered nitrous oxide and a hematoma block for analgesia in the outpatient reduction of fractures in children. *J Bone Joint Surg Am* 1995;77:335-9.
196. Hennrikus WL, Simpson RB, Klingelberger CE, Reis MT. Self-administered nitrous oxide analgesia for pediatric fracture reductions. *J Pediatr Orthop* 1994;14:538-42.
197. Wattenmaker I, Kasser JR, McGravey A. Self-administered nitrous oxide for fracture reduction in children in an emergency room setting. *J Orthop Trauma* 1990;4:35-8.
198. Gamis AS, Knapp JF, Glenski JA. Nitrous oxide analgesia in a pediatric emergency department. *Ann Emerg Med* 1989;18:177-81.
199. Kalach N, Barbier C, el Kohen R, et al. [Tolerance of nitrous oxide-oxygen sedation for painful procedures in emergency pediatrics: Report of 600 cases] [article in French]. *Arch Pediatr* 2002;9:1213-5.
200. Michaud L, Gottrand F, Ganga-Zandzou PS, et al. Nitrous oxide sedation in pediatric patients undergoing gastrointestinal endoscopy. *J Pediatr Gastroenterol Nutr* 1999;28:310-4.
201. Baskett PJ. Analgesia for the dressing of burns in children: A method using neuroleptanalgesia and Entonox. *Postgrad Med J* 1972;48:138-42.
202. Veerkamp JS, van Amerongen WE, Hoogstraten J, Groen HJ. Dental treatment of fearful children, using nitrous oxide. Part 1: Treatment times. *ASDC J Dent Child* 1991;58:453-7.
203. Veerkamp JS, Gruythuysen RJ, van Amerongen WE, Hoogstraten J. Dental treatment of fearful children using nitrous oxide. Part 3: Anxiety during sequential visits. *ASDC J Dent Child* 1993;60:175-82.
204. Veerkamp JS, Gruythuysen RJ, van Amerongen WE, Hoogstraten J. Dental treatment of fearful children using nitrous oxide. Part 2: The parent's point of view. *ASDC J Dent Child* 1992;59:115-9.
205. Veerkamp JS, Gruythuysen RJ, Hoogstraten J, van Amerongen WE. Dental treatment of fearful children using nitrous oxide. Part 4: Anxiety after two years. *ASDC J Dent Child* 1993;60:372-6.
206. Hout MI, Limb R, Livingston RL. Clinical effects of nitrous oxide conscious sedation in children. *Pediatr Dent* 2004;26:29-36.
207. Shapira J, Holan G, Guelmann M, Cahan S. Evaluation of the effect of nitrous oxide and hydroxyzine in controlling the behavior of the pediatric dental patient. *Pediatr Dent* 1992;14:167-70.
208. Primosch RE, Buzzi IM, Jerrell G. Effect of nitrous oxide-oxygen inhalation with scavenging on behavioral and physiological parameters during routine pediatric dental treatment. *Pediatr Dent* 1999;21:417-20.

209. McCann W, Wilson S, Larsen P, Stehle B. The effects of nitrous oxide on behavior and physiological parameters during conscious sedation with a moderate dose of chloral hydrate and hydroxyzine. *Pediatr Dent* 1996;18:35-41.
210. Wilson S, Matusak A, Casamassimo PS, Larsen P. The effects of nitrous oxide on pediatric dental patients sedated with chloral hydrate and hydroxyzine. *Pediatr Dent* 1998;20:253-8.
211. Litman RS, Kottra JA, Berkowitz RJ, Ward DS. Breathing patterns and levels of consciousness in children during administration of nitrous oxide after oral midazolam premedication. *J Oral Maxillofac Surg* 1997;55:1372-7.
212. Litman RS, Kottra JA, Verga KA, Berkowitz RJ, Ward DS. Chloral hydrate sedation: The additive sedative and respiratory depressant effects of nitrous oxide. *Anesth Analg* 1998;86:724-8.
213. American Academy of Pediatric Dentistry. Guideline on appropriate use of nitrous oxide for pediatric dental patients. *Pediatr Dent* 2005;27(suppl):107-9.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.