

# Guideline on Use of Anesthesia Personnel in the Administration of Office-based Deep Sedation/General Anesthesia to the Pediatric Dental Patient

## Originating Committee

Clinical Affairs Committee – Sedation and General Anesthesia Subcommittee

## Review Council

Council on Clinical Affairs

## Adopted

2001

## Revised

2005, 2007

## Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that there exists a patient population for whom routine dental care using nonpharmacologic behavior guidance techniques is not a viable approach.<sup>1</sup> The AAPD intends this guideline to assist the dental practitioner who elects to use anesthesia personnel for the administration of deep sedation/general anesthesia for pediatric dental patients in a dental office or other facility outside of an accredited hospital or surgicenter. This document discusses personnel, facilities, documentation, and quality assurance mechanisms necessary to provide optimal and responsible patient care.

## Methods

The revision of this guideline is based upon a review of current dental and medical literature pertaining to deep sedation/general anesthesia of dental patients, including the 2006 guideline on pediatric sedation co-authored by the American Academy of Pediatrics and the AAPD.<sup>2</sup> A MEDLINE search was performed using the terms “office-based anesthesia”, “pediatric sedation”, and “dental sedation”.

## Background

Pediatric dentists seek to provide oral health care to infants, children, adolescents, and those with special health care needs in a manner that promotes excellence in quality of care and concurrently induces a positive patient attitude toward dental treatment. Behavior guidance techniques have allowed most pediatric dental patients to receive treatment in the office with minimal discomfort and without expressed fear. Minimal or moderate sedation has provided others with the ability to accept treatment. There exists, however, a subset of children and special needs patients that, because of extensive treatment needs, acute situational anxiety, uncooperative age-appropri-

ate behavior, immature cognitive functioning, disabilities, or medical conditions, requires deep sedation/general anesthesia to receive dental treatment in a safe and humane fashion. Access to hospital-based anesthesia services may be limited for a variety of reasons, including restriction of coverage of by third party payors. Many pediatric dentists and others who treat children have sought to provide for the administration of deep sedation/general anesthesia by properly trained individuals in their offices or other facilities outside of the traditional surgical setting.

Utilizing deep sedation/general anesthesia in the dental office offers benefits for the patient and the dental team. Access to care may be improved. Both the number of dentists utilizing these advanced behavior guidance techniques and the availability of surgical setting options for the operating dentist may be increased. The treatment may be scheduled more easily and efficiently. Facility charges and administrative procedures may be less than those associated with a surgical center. Complex or lengthy treatment can be provided comfortably while minimizing patient memory of the dental procedure. By decreasing patient movement, the quality of care may be improved. The dentist can use his/her customary delivery system with access to supplemental equipment, instrumentation, or supplies should the need arise.

The use of anesthesia personnel to administer deep sedation/general anesthesia in the pediatric dental population is an accepted treatment modality.<sup>2-6</sup> The AAPD supports the provision of deep sedation/general anesthesia when, in the judgment of the dental practitioner, clinical indications have been met and additional properly-trained and credentialed personnel and appropriate facilities are used.<sup>1</sup> In many cases, the patient may be treated in an appropriate outpatient facility (including the dental office) because the extensive medical resources of a hospital are not necessary. It is well documented that this is a safe and viable approach.<sup>2-6</sup>

This guideline does not supercede, nor is it to be used in deference to, federal, state, and local credentialing and licensure laws, regulations, and codes. It cannot and does not predict nor guarantee a specific patient outcome.

## Recommendations

### Personnel

Office-based deep sedation/general anesthesia techniques require at least 3 individuals. The anesthesia care provider's responsibilities are to administer drugs or direct their administration and to observe constantly the patient's vital signs, airway patency, and adequacy of ventilation.<sup>2</sup> In addition to the anesthesia care provider and operating dentist, staff shall be present to assist the operator and support resuscitative measures, if required.

It is the obligation of treating practitioners, when employing anesthesia personnel to administer deep sedation/general anesthesia, to verify their credentials and experience.

1. The anesthesia care provider must be a licensed dental and/or medical practitioner with appropriate state certification for deep sedation/general anesthesia.
2. The anesthesia care provider must have completed a 1- or 2-year dental anesthesia residency or its equivalent, as approved by the American Dental Association (ADA), and/or medical anesthesia residency, as approved by the American Medical Association (AMA).
3. The anesthesia care provider must be licensed by and in compliance with the laws of the state in which he/she practices. Laws vary from state to state and must be respected.
4. If state law permits a certified registered nurse anesthetist or anesthesia assistant to function under the supervision of a dentist, the dentist is required to have completed training in deep sedation/general anesthesia and be licensed or permitted, as appropriate to state law.

The dentist and anesthesia care provider must be compliant with the AAP/AAPD's Guideline on Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures<sup>2</sup> or other appropriate guideline(s) of the ADA, AMA, and their recognized specialties. The recommendations in this document may be exceeded at any time if the change involves improved safety and is supported by currently-accepted practice and/or is evidence-based.

The dentist and anesthesia personnel must work together to enhance patient safety. Effective communication is essential. The dentist introduces the concept of deep sedation/general anesthesia to the parent and provides appropriate preoperative instructions and informational materials. The dentist or his/her designee coordinates medical consultations when necessary. The anesthesia care provider explains potential risks and obtains informed consent for sedation/anesthesia. Office staff should understand their additional responsibilities and special considerations (eg, loss of protective reflexes) associated with office-based deep sedation/general anesthesia.

Advanced training in recognition and management of pediatric emergencies is critical in providing safe sedation and anesthetic care. Although it is appropriate for the most experienced professional (ie, the anesthesia provider) to assume responsibility in managing anesthesia-related emergencies, the operating dentist and clinical staff need to maintain current expertise in basic life support. An individual experienced in recovery care must be in attendance in the recovery facility until the patient, through continual monitoring, exhibits respiratory and cardiovascular stability and appropriate discharge criteria<sup>2</sup> have been met. In addition, the staff of the treating dentist must be well-versed in emergency protocols (including cardiopulmonary resuscitation) and have contact numbers for emergency medical services and ambulance services. Emergency preparedness must be updated and practiced on a regular basis.

### Facilities

There exists a continuum among the levels of sedation and general anesthesia. These levels are not easily differentiated and patients may drift among them. When anesthesia care providers are utilized for office-based administration of deep sedation or general anesthesia, the facilities in which the dentist practices must meet the guidelines and appropriate local, state, and federal codes for administration of the highest possible level of sedation/anesthesia. Facilities should comply with applicable laws, codes, and regulations pertaining to controlled drug storage, fire prevention, building construction and occupancy, accommodations for the disabled, occupational safety and health, and disposal of medical waste and hazardous waste.<sup>3</sup> The treatment room must accommodate the dentist and auxiliaries, the patient, the anesthesia care provider, the dental equipment, and all necessary anesthesia delivery equipment along with appropriate monitors and emergency equipment. Expedient access to the patient, anesthesia machine (if present), and monitoring equipment should be available at all times.

It is beyond the scope of this document to dictate equipment necessary for the provision of deep sedation/general anesthesia, but equipment must be appropriate for the technique used and consistent with the guidelines for anesthesia providers, in accordance with governmental rules and regulations. Because laws and codes vary from state to state, the Guideline on Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures<sup>2</sup> must be followed as the minimum requirements. For deep sedation, there shall be continuous monitoring of oxygen saturation and heart rate and intermittent recording of respiratory rate and blood pressure. When adequacy of ventilation is difficult to observe, use of a precordial stethoscope or capnograph is encouraged.<sup>2</sup> An electrocardiographic monitor should be readily available for patients undergoing deep sedation. In addition to the monitors previously mentioned, a temperature monitor and pediatric defibrillator are required for general anesthesia.<sup>2</sup> Emergency equipment must be readily accessible and should include suction, the necessary drugs (including

oxygen), and age-/size-appropriate equipment to resuscitate and rescue a nonbreathing and unconscious pediatric dental patient and provide continuous support while the patient is being transported to a medical facility.<sup>2,7</sup> The treatment facility should have medications, equipment, and protocols available to treat malignant hyperthermia when triggering agents are used.<sup>3</sup> Recovery facilities must be available and suitably equipped. Backup power sufficient to ensure patient safety should be available in case of an emergency.<sup>3</sup>

### Documentation

Prior to delivery of deep sedation/general anesthesia, appropriate documentation shall address rationale for sedation/general anesthesia, informed consent, instructions to parent, dietary precautions, preoperative health evaluation, and any prescriptions along with the instructions given for their use.<sup>2</sup> Because laws and codes vary from state to state, the Guideline on Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures<sup>2</sup> must be followed as minimum requirements for a time-based anesthesia record.

1. Vital signs: Pulse and respiratory rates, blood pressure, and oxygen saturation must be monitored and recorded at least every 5 minutes throughout the procedure and at specific intervals until the patient has met documented discharge criteria.
2. Drugs: Name, dose, route, site, time of administration, and patient effect of all drugs, including local anesthesia, must be documented. When anesthetic gases are administered, inspired concentration and duration of inhalation agents and oxygen shall be documented.
3. Recovery: The condition of the patient, that discharge criteria have been met, time of discharge, and into whose care the discharge occurred should be documented.

Various business/legal arrangements may exist between the treating dentist and the anesthesia provider. Regardless, because services were provided in the dental facility, the dental staff must maintain all patient records, including time-based anesthesia records, so that they may be readily available for emergency or other needs. The dentist must assure that the anesthesia provider also maintains patient records and that they are readily available.

### Risk management and quality assurance

Dentists who utilize in-office anesthesia care providers should take all necessary measures to minimize risk to patients. Knowledge, preparation, and communication between professionals are essential. The dental office must provide quality of care equal to that of the hospital-based facility. Prior to subjecting a patient to deep sedation/general anesthesia, the patient must undergo a preoperative health evaluation.<sup>2,8</sup> High-risk patients should be treated in a facility properly equipped to provide for their care.<sup>2,8</sup> The dentist and anesthesia care provider must communicate during treatment to share concerns

about the airway or other details of patient safety. Furthermore, they must work together to develop mechanisms of quality assurance.

Untoward and unexpected outcomes must be reviewed to monitor the quality of services provided. This will decrease risk, allow for open and frank discussions, and improve the quality of care for the pediatric dental patient.

### References

1. American Academy of Pediatric Dentistry. Guideline on behavior guidance for the pediatric dental patient. *Pediatr Dent* 2006;28(suppl):97-105.
2. American Academy of Pediatrics, American Academy of Pediatric Dentistry. Guideline for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures. *Pediatr Dent* 2006;28(suppl):115-32.
3. American Society of Anesthesiologists. Guidelines for office-based anesthesia. 2004. Available at <http://www.asahq.org/publicationsAndServices/standards/12.pdf>. Accessed May 5, 2007.
4. American Dental Association. Policy statement: The use of conscious sedation, deep sedation, and general anesthesia in dentistry. 2005. Available at: <http://www.ada.org/prof/resources/positions/statements/useof.asp>. Accessed May 6, 2007.
5. Nick D, Thompson L, Anderson D, Trapp L. The use of general anesthesia to facilitate dental treatment. *Gen Dent* 2003;51:464-8.
6. Wilson S. Pharmacologic behavior management for pediatric dental treatment. *Pediatr Clin North Am* 2000;47(5):1159-73.
7. American Society of Anesthesiologists. Guidelines for ambulatory anesthesia and surgery. 2003. Available at <http://www.asahq.org/publicationsAndServices/standards/04.pdf>. Accessed May 5, 2007.
8. American Dental Association. Guidelines for the use of conscious sedation, deep sedation, and general anesthesia for dentists. 2005. Available at: [http://www.ada.org/prof/resources/positions/statements/anesthesia\\_guidelines.pdf](http://www.ada.org/prof/resources/positions/statements/anesthesia_guidelines.pdf). Accessed May 6, 2007.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.