# Scientific Article



# Mixed Dentition Cavitated Caries Incidence and Dietary Intake Frequencies

Oitip Chankanka, DDS, MSc, PhD<sup>1</sup> • Teresa A. Marshall, PhD<sup>2</sup> • Steven M. Levy, DDS, MPH<sup>3</sup> • Joseph E. Cavanaugh, PhD<sup>4</sup> • John J. Warren, DDS, MS <sup>5</sup> • Barbara Broffitt, MS<sup>6</sup> • Justine L. Kolker, DDS, MS, PhD<sup>7</sup>

**Abstract:** *Purpose:* This study examined risk factors for children having new cavitated caries between 5 and 9 years old. **Methods:** Subjects were Iowa Fluoride Study cohort children (mostly Caucasian and of relatively high socioeconomic status) with both primary and mixed dentition caries exams and at least 2 diet diaries recorded between 5 and 8 years old (n=198). Using surface-specific transitions, combined counts of new cavitated caries ( $d_{2,3}f$  and/or  $D_{2,3}F$ ) were determined from 4 primary second molars, 8 permanent incisors, and 4 permanent molars. Food and beverage intake frequencies were abstracted. Other factors were assessed using periodic questionnaires. Logistic regression identified predictors of new cavitated caries. **Results:** Thirty-seven percent had new cavitated caries. The mean new cavitated caries count for all children was 1.17 surfaces ( $\pm 2.28$  SD). In multivariable logistic regression, the following were significantly associated (P<.10) with having new cavitated caries: noncavitated caries experience at 5 years old (odds ratio [OR]=2.67, P=.03); cavitated caries experience at 5 years old (OR=3.39, P=.004); greater processed starch at snack frequency (OR=3.87, P=.07); being older (OR=1.68, P=.04); and less frequent tooth-brushing (P=.001). **Conclusion:** Results suggested that increased tooth-brushing frequency and reduced consumption of processed starches as snacks may reduce caries incidence in younger school-aged children. (Pediatr Dent 2011; 33:233-40) Received November 1, 2009 I Last Revision March 2, 2010 I Accepted August 11, 2010

KEYWORDS: DENTAL CARIES, INCIDENCE, CARIOGENIC DIET, MIXED DENTITION, CHILD

Dental caries remains a major public health problem for U.S. children. The National Health and Nutrition Examination Survey 1999-2002 reported a national prevalence of 49% of children with dental caries experience in the primary teeth and 20% of children with caries experience in the permanent teeth<sup>1</sup> for 6- to 11-year-old children. Early school-age children are at a stage of transition from the primary to the mixed dentition. Moreover, they also begin elementary school, which generally changes beverage and food intake patterns.

Most studies have assessed risk factors for dental caries in the mixed dentition cross-sectionally. Since the dental caries process takes time to develop into clinically detectable lesions, risk factors for dental caries should be assessed by examining factors that occur before and during the time of clinical caries detection. There are a limited number of nonexperimental studies, however, that have assessed caries incidence/increment in the transition period from primary to mixed dentition.

Associations between previous cavitated caries and new cavitated caries in primary school-age children have been

investigated.<sup>2-9</sup> Consistently, findings show a strong and significant association between previous caries experience and new caries. Since previous dental caries experience is not a modifiable factor, studies should examine modifiable risk factors for dental caries beyond previous caries experience. Few studies<sup>3,4,9,10</sup> have included modifiable factors, such as dietary intakes, fluoride usage, and tooth-brushing frequency. Two of these studies<sup>3,10</sup> found significant associations between consumption of sweets/sweetened drinks and new cavitated caries, while the other 2<sup>4,9</sup> found no significant association between dietary consumption (sweets, sugars, or snacks) and new cavitated caries.

The objective of the analyses described in this paper was to examine modifiable and nonmodifiable risk factors for children having new cavitated caries from 5- to 9-years-old. The frequencies of dietary intake of specific food and beverage categories and other caries-related factors were compared between 5- to 9-year-olds with and without new cavitated caries.

## Methods

The data used in this analysis are part of the ongoing, longitudinal Iowa Fluoride Study (IFS). Subjects were recruited immediately postpartum from 8 Iowa hospitals and followed thereafter by mail and periodic assessments/exams. The children were mostly Caucasian and relatively high socioeconomic status (SES).<sup>11</sup> Details of the recruitment of IFS subjects have been published elsewhere.<sup>12</sup> This study was approved by the Human Subjects Committee at the

<sup>&</sup>lt;sup>1</sup>Dr. Chankanka is faculty. Department of Preventive Dentistry. Faculty of Dentistry. Prince of Songkla University. Songkhla province, Thailand: <sup>2</sup>Dr. Marshall is an associate professor, Drs. <sup>3</sup>Levy and <sup>5</sup>Warren are professors, and <sup>6</sup>Ms. Broffitt is Research Statistician, all in the Department of Preventive and Community Dentistry, and <sup>7</sup>Dr. Kolker is an associate professor. Department of Operative Dentistry, all in the College of Dentistry, and <sup>4</sup>Dr. Cavanaugh is professor, Department of Biostatistics, College of Public Health, all at the University of Iowa. Correspond with Dr. Chankanka at oitip.c@psu.ac.th

University of Iowa, Iowa City, Iowa. Parents of the participating subjects provided informed consent and children provided assent for the exams.

Subjects. Subjects with both primary dentition (-5 years old) and mixed dentition (-9 years old) caries exams and at least 2 diet diaries, including at least 1 from 5- to 6.5-years-old and 1 from 7- to 8.5-years-old (to ensure at least 1 diary before starting school and at least 1 after) were included in these analyses (N=198).

**Dietary information.** The food and beverage intakes for 2 weekdays and 1 weekend day were recorded by parents using 3-day diaries that were sent every 1.5 to 6 months (1.5-months- to 8.5-years-old).<sup>13</sup> Detailed information of consumption times, types of items, and quantities were collected. Annual dietary diaries (ie, at 5-8 years old) were abstracted by trained registered dietitians or diet technicians. If the 5-, 6-, 7-, and/or 8-year-old diary was missing, then a substitution was made using the diaries 6 months succeeding or preceding the yearly diary.

Food and beverage intake frequencies were estimated from the diaries<sup>13</sup> and categorized so that consumption within a 30-minute interval counted as 1 eating event. More than 1 serving of the same beverage or food consumed within 30 minutes was considered 1 event. The eating events then were identified as either meals or snacks, based on the time of consumption and nature of the food. Only 3 meals per day were allowed: 1 during the morning; 1 at the middle of the day; and 1 during evening hours. Unlimited eating events, however, were allowed for snacks.

Foods and beverages were then classified into categories. Beverages were categorized by type of beverage and included: milk; 100% juices; juice drinks; powder-sugared beverages; regular (sugared) soda pop; diet soda pop; sports drinks; and water. Foods were categorized based on sugar and/or starch content and included: sugar-based desserts (jelly, pudding, etc); candy; added sugar (table sugar, honey, brown sugar, etc); baked starch with sugar (cookies, cake, etc); unsweetened cereals; presweetened cereals; unprocessed starches (boiled potato, bread, rice, etc); and processed starches (potato chips, etc). The intake frequency of each food and beverage category was counted and averaged for each child across the 3 days for each annual diary (at 5, 6, 7, or 8-years-old). Results at 5 and/or 6-years-old and at 7 and/or 8-years-old were averaged separately, and these 2 means were subsequently averaged to form an overall measure for each child. The dietary variables were presented in order by percentage of children with some intake (5-8 years old) at snacking occasions ineach of the beverage, sugar-based, and starchbased food groups.

Other related factors. Daily tooth-brushing frequency and composite water fluoride intakes were determined based on data from IFS questionnaires that were sent to parents at the same time as the diaries. Composite water fluoride levels were determined from all sources, including available public water documentation and an assay of multiple sources of water (ie, home/school, bottled/filtered/tap water) at each time point for each individual child.<sup>11,14</sup> Composite water fluoride levels and daily tooth-brushing frequency<sup>15</sup> from all available 5- to 8-year-old data for each child were averaged.

Three categories of SES were created based on family income and mother's education data from baseline questionnaires (completed at birth). The high SES category included children from families with incomes from \$30,000 to \$49,999 a year whose mothers had a graduate/professional degree, or from families with incomes of more than \$50,000/ year (regardless of the mother's educational level). Children from families with incomes less than \$30,000 a year whose mother had 2 years or less in college were categorized as low SES. All other children were placed in the middle SES category.

**Dental data.** Children were examined for dental caries at 5-years-old (primary dentition) and 9-years-old (mixed dentition) by the same trained and calibrated examiners. The examinations were conducted using a portable chair, halogen headlight and a DenLite mirror (Welch-Allyn Medical Product, Inc, Skaneatele Falls, NY).<sup>16</sup> The teeth were dried and examined primarily by visualization. Any questionable lesions, however, were confirmed by an explorer.<sup>16</sup> This study used the diagnostic criteria that were modified from the D<sub>1</sub>-D<sub>4</sub> system of Pitts and colleagues.<sup>17,18</sup> The examiners did not differentiate cavitated enamel (D<sub>2</sub>/d<sub>2</sub>) and dentine lesions (D<sub>3.4</sub>/d<sub>3.4</sub>), thus those lesions were categorized together as D<sub>2.2</sub>/d<sub>2.3</sub>.

Combined count of new cavitated caries  $(d_{2.3}f \text{ and/or } D_{2.3}F)$  included cavitated lesions and restored lesions from 4 primary second molars, 8 permanent incisors, and 4 permanent first molars. The counts of new cavitated caries were determined using surface-specific transitions during exams taken between 5 and 9-years-old. Having new cavitated caries was used as the dependent binary variable (none vs 1 or more new lesions and/or restorations) in these analyses.

Data analysis. Logistic regression analyses were used to explore the relationships between having or not having new cavitated caries from 5- to 9-years-old and other variables: dietary intakes; gender; SES; age at mixed dentition exam; cavitated caries experience at 5-years-old; noncavitated caries experience at 5-years-old; tooth-brushing frequency; and composite water fluoride level.

First, univariable logistic regression analyses were modeled separately for each dietary variable or other related factor. For dietary variables, only variables with P < .15 from the univariable logistic regression were selected for the next steps. The pairwise correlations between all pairs of continuous variables of selected dietary variables and other variables were determined. For those pairs with moderate to high correlations (Pearson's coefficients >0.3), only the variable with the stronger association with caries incidence would be included in an initial model for multivariable logistic regression analysis. All 2-way interactions were then assessed in bivariable models, except for those based on a product of 2 continuous variables. Dietary variables, other related variables, and interaction terms with P<.15 (and the corresponding main effects) were considered in the multivariable modeling and were used to create 2 separate models. The 2 models were initially based on:

- 1. dietary and other variables (tooth-brushing frequency, composite water fluoride level, gender, age at mixed dentition exam, and SES) excluding previous dental caries experience; and
- 2. all variables (including previous caries experience).

For each of these model formulations, a backward elimination procedure was performed to determine the final model. Variables that did not retain significance at P<.10 were omitted sequentially based on the associated P-value. All data were analyzed using Proc Logistic in SAS 9.1 (SAS Institute Inc, Cary, NC).

## Results

There were 198 children (55% girls) who met the inclusion criteria of having both exams and sufficient numbers of diet diaries. Mean ( $\pm$ SD) ages at the primary dentition and mixed dentition exams were 5.1 ( $\pm$  0.4) and 9.2 ( $\pm$  0.7), respectively. Most were in the middle or high SES categories (35% and 42%, respectively). The prevalences of noncavitated

caries at the primary and at the mixed dentition exams were approximately 19% and 39%, respectively. Cavitated caries prevalence at the primary and at the mixed dentition exams were approximately 21% and 49%, respectively. Sixty-three percent of the children had no new cavitated caries (zero surfaces of new cavitated caries). Among the 37% of children with some new cavitated caries, more than half (20%) had only 1 to 2 surfaces of new cavitated caries, approximately one fourth (9%) had 3 to 4 surfaces of new cavitated caries, and less than one fourth (8%) had 5 or more surfaces of new cavitated caries. The overall mean new cavitated caries for all children was 1.17 surfaces (±2.28 SD; range=0-15), with a mean only among those affected of 3.2 surfaces (±2.8 SD; range=1-15).

Descriptive analyses presenting medians and 25th and 75th percentiles of food and beverage intake frequencies at meals and at snacking occasions for individuals' intakes for each year (5-8 years old) and averaged from the 2 to 4 diaries (5-8 years old) are presented in Table 1. This table also

Variable		Median (25 <sup>th</sup> , 75 <sup>th</sup> percentile)							
		5-8 ys old (n=198)	5 ys old (n=171)	6 ys old (n=173)	7 ys old (n=164)	8 ys old (n=150)			
Beverage (occasions)									
Milk	Snack	0.2 (0.1, 0.4)	0.3 (0, 0.7)	0.3 (0, 0.7)	0 (0, 0.3)	0 (0, 0.3)			
	Meal	1.6 (1.2, 1.8)	1.7 (1.0, 2.0)	1.5 (1.0, 2.0)	1.3 (1.0, 2.0)	1.7 (1.0, 2.0)			
100% juice	Snack	0.1 (0, 0.3)	0 (0, 0.3)	0 (0, 0.3)	0(0, 0)	0(0,0)			
	Meal	0.3 (0.1, 0.6)	0.3 (0, 0.7)	0.3 (0, 0.7)	0 (0, 0.7)	0(0, 0.7)			
Regular soda pop	Snack	0.1 (0, 0.3)	0 (0, 0.3)	0 (0, 0.3)	0 (0, 0.3)	0 (0, 0,3)			
	Meal	0.3 (0.1, 0.4)	0.3 (0, 0.5)	0 (0, 0.3)	0.3(0, 0.3)	0 (0, 0, 5)			
Juice drinks	Snack	0.1 (0, 0.2)	0 (0, 0.3)	0 (0, 0)	0 (0, 0.2)	0 (0, 0)			
	Meal	0.2 (0, 0.3)	0 (0, 0.3)	0 (0, 0,3)	0 (0, 0.3)	0 (0, 0,3)			
Powder-sugared beverages	Snack	0 (0, 0.1)	0 (0, 0)	0 (0, 0)	0(0,0)	0(0, 0)			
	Meal	0 (0, 0.1)	0 (0, 0)	0 (0, 0)	0(0, 0)	0 (0, 0)			
Sport drinks	Snack	0 (0, 0)	0 (0, 0)	0 (0, 0)	0(0, 0)	0 (0, 0)			
-	Meal	0 (0, 0)	0 (0, 0)	0 (0, 0)	0(0, 0)	0 (0, 0)			
Diet soda pop	Snack	0(0, 0)	0 (0, 0)	0 (0, 0)	0(0, 0)	0 (0, 0)			
	Meal	0 (0, 0)	0 (0, 0)	0 (0, 0)	0(0, 0)	0(0, 0)			
Water	Snack	0.8 (0.3, 1.5)	0.7 (0, 1.3)	0.7 (0, 1.3)	0.7(0, 1.7)	0.7 (0, 1.7)			
	Meal	0.2 (0, 0.3)	0 (0, 0.3)	0 (0, 0.3)	0 (0, 0.3)	0 (0, 0.3)			
Food (occasions)									
Sugar-based desserts	Snack	0.3(0.1, 0.5)	03(0.05)	0.3(0.07)	03(007)	0 (0 0 2)			
ougar outed detterts	Meal	0.2(0.1, 0.3)	0.0(0, 0.3)	0.0(0, 0.7)	0.3(0, 0.7)	0(0, 0.3)			
Candy	Snack	0.3(0.1, 0.5)	0(0, 0.3)	0.3(0, 0.5)	0(0, 0.5)	0(0, 0.3)			
)	Meal	0.04(0.0.2)	0(0, 0.9)	0.0(0, 0.7)	0.0(0, 0.3)	0(0, 0.3)			
Added sugars	Snack	0(0, 0, 2)	0(0,0)	0(0, 0)	0(0, 0.5)	0(0,0)			
	Meal	0.3(0.2, 0.6)	03(0,07)	0.3(0,07)	03(0,07)	0.3(0,07)			
Baked starches with sugar	Snack	0.5(0.2, 0.8)	0.7(0.3,1.0)	0.7(0.3, 1.0)	0.3(0, 0.7)	0.5(0, 0.7)			
	Meal	0.5(0.3, 0.0)	0.3(0, 0.7)	0.3(0.3, 0.7)	0.5(0, 0.7)	0.4(0, 0.7)			
Processed starches	Snack	0.3(0.1, 0.4)	0(0, 0.3)	0.3(0.03)	0.0(0.03, 0.0)	0.0(0.0, 0.7)			
	Meal	0.3(0.2, 0.5)	0.3(0, 0.7)	0.3(0, 0.7)	03(0,0.5)	0.3(0.07)			
Unprocessed starches	Snack	0.2(0.1, 0.3)	0.0(0, 0.3)	0.0(0, 0.7)	0.5(0, 0.7)	0.3(0, 0.7)			
	Meal	25(23,28)	23(2030)	25(20,30)	27(20.30)	27(2330)			
Presweetened cereals	Snack	0(0,0)	0(0,0)	2.9(2.0, 9.0)	0(0,0)	(2.3, 3.0)			
	Meal	0.3(0.2, 0.5)	0.3(0,07)	0.3(0,07)	0.3(0,07)	0(0, 0)			
Unsweetened cereals	Snack	0 (0, 0)	0(0, 0)	0.0(0, 0.7)	0.5(0, 0.7)	0.5(0, 0.5)			
		~ (0, 0)	0 (0, 0)	$\circ$ $(\circ, \circ)$	$v_{(0,0)}$	0(0, 0)			

Table continues on next page

presents the percentages of children who had consumed from each specific food/beverage category at 5- to 8-years-old and at any of the 2 to 4 available time points (5-8 years old). Based on these percentages, milk was the most common

Table 1. Contin	uation							
37 - 11		Children with some intake (%) <sup>†</sup>						
Variable		5-8 ys old (n=198) %	5 ys old (n=171) %	6 ys old (n=173) %	7 ys old (n=164) %	8 ys old (n=150) %		
Beverage (occasion	ns)							
Milk	Snack	80	56	55	40	37		
	Meal	99	94	97	96	95		
100% juice	Snack	65	47	33	23	17		
	Meal	80	58	52	48	41		
Regular soda	Snack	65	30	34	32	30		
рор	Meal	80	55	47	57	49		
Juice drinks	Snack	57	26	23	25	23		
	Meal	69	32	36	37	36		
Powder-sugared	Snack	41	18	15	17	12		
beverages	Meal	39	18	17	13	12		
Sport drinks	Snack	8	2	2	1	2		
•	Meal	5	4	2	2	5		
Diet soda pop	Snack	19	6	8	8	6		
	Meal	23	5	8	9	11		
Water	Snack	93	71	73	74	73		
	Meal	62	39	38	40	39		
Food (occasions)								
Sugar-based	Snack	86	55	61	51	48 <sup>·</sup>		
desserts	Meal	78	42	47	39	42		
Candy	Snack	81	46	51	55	48		
,	Meal	50	22	23	27	19		
Added sugars	Snack	46	19	21	14	15		
Ū.	Meal	86	58	65	60	55		
Baked starches	Snack	96	80	77	69	69		
with sugar	Meal	95	64	75	79	77		
Processed	Snack	81	45	52	43	45		
starches	Meal	95	65	69	68	64		
Unprocessed	Snack	81	44	44	39	42		
starches	Meal	100	100	99	100	99		
Presweetened	Snack	24	8	9	9	7		
cereals	Meal	87	56	61	59	56		
Unsweetened	Snack	16	6	6	5	2		
cereals	Meal	55	29	22	23	23		

\* To compute these measures, mean dietary intakes from 5 and/or 6-years-old and 7 and/or 8-years-old were calculated, and these 2 means were subsequently averaged.

Note: There were 22% additional abstracted dietary diary responses at ages 5½, 6½, 7½, and 8½ that are not included in the year-specific data, but are included as substitutes when 5- to 8-year-olds were missing. † Percentage of children with some intake from diaries (among 2-4 annual time points per child).

drink for this age group (99%). Other common beverages were: 100% juice; regular soda pop; and juice drinks. Approximately one fourth of the children consumed diet soda pop. Sports drinks were not a common beverage for these

children (≤10%).

Medians, means, and standard deviations of other related factors at 5- to 8-years-old and averages of any available data from 5- to 8-years-old are presented in Table 2. On average, children brushed 1.5 times per day, and water fluoride levels fell into the optimal range of 0.7 to 1.2 ppm. More than three fourths of the children did not have noncavitated or cavitated caries at the primary dentition exam (81% and 79%, respectively). Mean surfaces of noncavitated caries and cavitated caries experience at approximately 5-years-old were  $0.5\pm1.6$  and  $1.2\pm4.3$ , respectively.

Table 3 summarizes the univariable logistic regression analyses assessing the associations between having new cavitated caries and the numbers of occasions for the dietary variables. These analyses were used to screen the dietary variables for inclusion in the multivariable logistic regression. Significant variables (P<.15) were snacktime intake of regular soda pop, unprocessed starches, and processed starches.

The univariable logistic regression analyses of factors other than dietary variables are presented in Table 4. Low SES was significantly associated with having new cavitated caries. Greater toothbrushing frequency was significantly associated with not having new cavitated caries, while gender, age at mixed dentition exam, and composite water fluoride level were not significantly associated with having or not having new cavitated caries. Having noncavitated caries experience at 5-years-old (odds ratio [OR]=4.00, 95% confidence interval [CI]=1.91, 8.39, P<.001) and having cavitated caries experience at 5years-old (OR=4.44, 95% CI=2.16, 9.12, P<.001) were significantly associated with having new cavitated caries from 5- to 9-years-old.

The multivariable logistic regression analyses are presented in Table 5. Three dietary variables were initially included: snacktime intake of regular

Table 2. DESCRIPTI	VE ANALYSIS	OF OTHER I	RELATED FA	CTORS							
Variable			Median				1	Mean±(SD)			
	5-8 ys old (n=198)	5 ys old (n=171)	6 ys old (n=173)	7 ys old (n=164)	8 ys old (N=150)	5-8 ys old (n=198)	5 ys old (n=171)	6 ys old (n=173)	7 ys old (n=164)	8 ys old (n=150)	
Daily tooth-brushing frequency	1.50	1.0	1.0	2.0	2.0	1.5±0.5	1.4±0.6	1.5±0.6	1.5±0.6	1.5±0.6	
Composite water fluoride level (ppm)	0.93	0.94	0.95	0.98	0.94	0.8±0.4	0.8±0.4	0.8±0.4	0.8±0.3	0.8±0.4	

soda pop, unprocessed starches, and processed starches. There was no pairwise collinearity concern with the 6 continuous variables (2 food variables, 1 beverage variable, toothbrushing frequency, age at mixed dentition exam, and composite water fluoride level).

Model 1 initially considered 8 variables, including 3 dietary variables and 5 other variables (daily tooth-brushing frequency, composite water fluoride level, gender, age at mixed dentition exam, and SES), and 2 interaction terms (between gender and composite water fluoride level and between low SES and age at mixed dentition dental exam). Greater snacktime intake frequency of processed starches (OR=1.50, P=.04) and low SES (OR=2.27, P=.04) were significantly associated with having new cavitated caries. Greater daily tooth-brushing frequency was significantly associated with not having new cavitated caries (OR=0.32, P<.001). A higher composite water fluoride level was significantly associated with not having new cavitated caries caries in girls; however, it was not significantly associated

Table 3. UNIVARIABLE LOGISTIC REGRESSION MODELS USING DIETARY VARIABLES TO PREDICT OCCURRENCE OF NEW CAVITATED CARIES FROM 5- TO 9-YEARS-OLD* (N=198)					
Variable		Odds ratio (95% confidence interval)	<i>P</i> -value		
Beverages (occasions)					
Milk	Snack	1.29 (0.64, 2.62)	48		
	Meal	0.87 (0.52, 1.46)	.60		
100% juice	Snack	0.90 (0.19, 4.40)	.90		
,	Meal	0.79 (0.35, 1.78)	.57		
Regular soda pop	Snack	5.15 (1.003, 26.43)	<.05†		
0 11	Meal	1.75 (0.52, 5.94)	.37		
Juice drinks	Snack	1.09 (0.28, 4.19)	.90		
	Meal	0.53 (0.16, 1.78)	.31		
Powder-sugared	Snack	2.04 (0.47, 8,89)	.35		
beverages	Meal	0.44 (0.04, 5.14)	.51		
Diet soda pop	Snack	0.48 (0.03, 7.03)	.60		
	Meal	1.92 (0.05, 70.26)	.72		
Water	Snack	1.03 (0.75, 1.41)	.18		
	Meal	1.30 (0.56, 3.04)	.34		
Food (occasions)					
Sugar-based desserts	Snack	1.33 (0.48, 3.72)	.59		
	Meal	0.38 (0.08, 1.76)	.22		
Candy	Snack	0.77 (0.31, 1.95)	.59		
	Meal	0.54 (0.07, 4.14)	.56		
Added sugar	Snack	4.42 (0.42, 47.10)	.22		
D-1-11	Meal	0.50 (0.19, 1.34)	.17		
baked starch	Snack	0.91 (0.38, 2.18)	.83		
Processed starches	Speek	0.79(0.51, 2.05) 2.85(0.83, 0.83)	.05 10t		
riocesseu statenes	Meal	0.79(0.25, 9.83)	.10		
Unprocessed starches	Snack	3.16 (0.76, 13.15)	12		
	Meal	1.16 (0.62, 2.18)	64		
Presweetened cereals	Snack	0.43 (0.01, 23.54)	68		
cereals	Meal	1.72 (0.56, 5.28)	.35		

\* Cavitated caries from exam at approximately 9-years-old that was not present at 5-years-old. <u>Note:</u> Each line of this table represents a separate regression model.

† P<.15 for initial consideration for multivariable logistic regression.

with not having new cavitated caries in boys. The *P*-value for the Hosmer and Lemeshow goodness-of-fit test was 0.72, indicating an adequate fit of the final model.

Model 2 initially included 10 variables (8 variables from model 1, noncavitated caries experience at 5-years-old (yes/ no) and cavitated caries experience at 5-years-old (yes/no), and 2 interaction terms (between gender and composite water fluoride level and between low SES and age at mixed dentition dental exam). Having noncavitated caries experience at 5-years-old (OR=2.67, P=.03), having cavitated caries experience at 5-years-old (OR=3.39, P=.004), greater snacktime intake frequency of processed starches (OR=3.87, P=.07), and being older (OR=1.68, P=.04) were significantly associated (P<.10) with having new cavitated caries from 5- to 9-years-old. Greater daily tooth-brushing frequency was significantly associated with not having new cavitated caries (OR=0.28, P=.001). A higher composite water fluoride level was significantly associated with not having new cavitated caries in girls (OR=0.88, P=.08); however, it was not significantly associated with cavitated caries incidence in boys. The P-value for the Hosmer and Lemeshow goodnessof-fit test was 0.93, indicating adequate fit of the final model.

Although the main analyses have focused on intake frequency data, additional analyses were conducted using quantity of intake data. These generally showed similar directions

Table 4. UNIVARIABLE LOGISTI USING DEMOGRAPHIC VARIABLES TO PREDIC CAVITATED CARIES FR (N=198)	UNIVARIABLE LOGISTIC REGRESSION MODELS USING DEMOGRAPHIC VARIABLES AND OTHER VARIABLES TO PREDICT OCCURRENCE OF NEW CAVITATED CARIES FROM 5- TO 9-YEARS-OLD* (N=198)					
Variable	Odds ratio (95% confidence interval)	<i>P</i> -value				
Gender • Boys • Girls	1.01 (0.59, 1.89)	.85				
Age at mixed dentition dental exam (year)	1.24 (0.83, 1.85)	.29				
Age interval between 1 <sup>st</sup> and 2 <sup>nd</sup> exam (year)	1.20 (0.82, 1.74)	.35				
Socioeconomic status • Low • Middle/high	2.07 (1.04, 4.13)	.04				
Daily tooth-brushing frequency from 5-8 years old	0.37 (0.20, 0.69)	.002				
Composite water fluoride level (0.1 ppm units) from 5-8 years old	0.91 (0.40, 2.04)	.81				
Non-cavitated caries experience at 5-years-old • Yes • No	4.00 (1.91, 8.39)	<.001				
Cavitated caries experience at 5-years-old • Yes • No	4.44 (2.16, 9.12)	<.001				

\* Cavitated caries from exam at approximately 9-years-old that was not present at 5-years-old. <u>Note:</u> Each line of this table represents a separate regression model.

and strengths of association to the analyses using frequency data. For example, in additional univariable analyses, both greater regular soda pop intake frequency and quantity at snacking occasions tended to increase the risk of having new cavitated caries (P<.05 and P=.12, respectively). Both greater regular pop intake frequency and quantity at meals tended to increase the risk of having new cavitated caries, but were not statistically significant (P=.37 and P=.43, respectively).

OCCURRENCE OF NEW OLD* (n=198)	CAVITATED	CARIES FROM 5 TO	9 YEARS
Models	Estimate	Odds ratio (95% confidence interval)	P-value
Model 1 (all variables excluding previ	ous caries exp	erience <sup>†</sup> initially inclu	ded)
Intercept			
Girls	2.77		
Boys	0.38		
starches	1.50	4.48 (1.14, 17.70)	.04
Daily tooth-brushing frequency from 5-8 years old	-1.15	0.32 (0.16, 0.63)	<.001
Socioeconomic status			
Low Middle/high (Reference category)	0.41	2.27 (1.08, 4.80)	.04
Composite water fluoride level (0.1 ppm units) from 5-8 years old			
Girls Boys	-0.16 0.06	0.86 (0.75, 0.98) 1.06 (0.93, 1.20)	.03 .39
Model 2 (all variables <sup>‡</sup> initially includ	ed)		
Intercept			
Girls	1.91		
Boys .	-3.79		
Snacktime intake of processed starches	1.35	3.87 (0.93, 16.16)	.07
Daily tooth-brushing frequency from 5-8 years old	-1.29	0.28 (0.13, 0.59)	.001
Non-cavitated caries experience at 5-years-old			
Yes No (Reference category)	0.49	2.67 (1.11, 6.42)	.03
Cavitated caries experience at 5-years-old			
Yes No (Reference category)	0.61	3.39 (1.48, 7.78)	.004
Age at mixed dentition exam (year)	0.52	1.68 (1.04, 2.73)	.04
(0.1 ppm units) from 5-8 years old	-0.13	0.88 (0.76, 1.01)	.08
Girls Boys	0.05	1.05 0.93, 1.20)	.44

\* Cavitated caries from exam at approximately 9-years-old that was not present at 5years-old. <u>Note:</u> The other variables initially considered for inclusion (not shown in the table) were removed in the backward elimination (*P*<0.10 to remain).

† All variables excluding caries experience: snacktime intake of regular soda pop; snacktime intake of processed starches; snacktime intake of unprocessed starches; tooth-brushing frequency; composite water fluoride level; gender; age at mixed dentition exam; and socio-economic status.

<sup>‡</sup> All variables: snacktime intake of regular soda pop; snacktime intake of processed starches; snacktime intake of unprocessed starches; tooth-brushing frequency; composite water fluoride level; gender; age at mixed dentition exam; socioeconomic status; previous noncavitated caries; and previous cavitated caries.

Although not our first emphasis, the relationship between SES and previous noncavitated caries and the relationship between processed starches intake frequency at snacking occasions and previous cavitated caries experience were explored. Thirty-three percent of children of low SES families had previous noncavitated caries experience, while only 15% of children of middle/high SES families had previous noncavitated caries experience. Additionally, 11% of children who

had reported no intake frequency of processed starches at snacking occasions had previous cavitated caries experience, while 24% of children with some intake of processed starches at snacktime had previous cavitated caries experience.

#### Discussion

This study sample was predominantly Caucasian, of relatively high SES, and had low-moderate caries risk. More than one third, however, had new caries in the 4-year period and nearly 10% had an average of more than 1 new surface per year. Even in a low-moderaterisk population, caries in school-age children is still a public health problem.

Infection by cariogenic bacteria and the demineralization process on tooth surfaces are part of the true dental caries disease processes, but noncavitated caries and cavitated caries are signs of the chronic disease that can be more reliably assessed clinically. Routine assessments of dynamic changes in bacteria and/or tooth demineralization that can be reversed or remineralized in a short time are not feasible clinically. Thus, the present study assessed signs of dental caries (clinical exam) and used them in determining the outcome variable of new cavitated caries.

From the descriptive analyses of the dietary variables, more than three fourths of these children consumed regular soda pop, and approximately one fourth consumed diet soda pop. Diet soda pop has been available in the United States for more than 50 years. The results, however, reveal that many more children consumed regular vs diet soda pop. In addition, consumption of presweetened cereal was more common among children than consumption of unsweetened cereals.

Several other studies found a strong association between previous caries experience of the primary dentition and new caries of the mixed dentition.<sup>2-9</sup> Our findings strongly support this conclusion. While presence of caries experience is a predictor of future caries, it is not possible to intervene and change previous caries experience. Therefore, studies should assess other risk factors besides previous caries experience.

The present analyses did not use a traditional 5% significance level but instead used a 10% significance level. The univariable results showed strong relationships between caries incidence and previous caries experience and weaker relationships between caries incidence and certain other explanatory factors, such as SES and processed starches intake at snacking occasions. In the multivariable logistic regression analyses, when a 0.05 significance level was used in the backward elimination

procedure, dietary variables were not retained in the final multivariable model. Because previous caries experience and the dental caries outcome is the same disease, it is possible that they share the same risk factors. Thus, we further explored the relationship between these factors and previous noncavitated and cavitated caries experiences. The results indicate associations between: SES and noncavitated caries experiences; and frequency of snacktime intake of processed starches and cavitated caries experience. Because of these relationships, the simultaneous inclusion of previous caries experience in a multivariable model containing the other explanatory factors reduced the contribution of the other factors. Therefore, the dietary variables which had weaker relationships with caries incidence would have been dropped out from the model with the traditional P-value of .05. Thus, we decided to use a larger P-value to allow those variables with weaker relationships to be considered in the multivariable logistic regression. These approaches allow us to examine other related factors, such as dietary and fluoriderelated factors, which are modifiable factors, in contrast with previous caries experience, which is not modifiable.

A study by Leroy et al.,<sup>9</sup> which included previous caries experience and other related factors in the multivariable model, reported similar findings in which other related factors that were significantly associated with new caries in the univariable analyses were not retained in the final model. Tagliaferro et al.,<sup>4</sup> discussed their study's findings that oral hygiene habits variable did not reach statistical significance in the final model because of the strong association between previous caries experience and caries incidence.

To show how explanatory variables affected each other in the model, we developed 2 separate models. Model 1 initially included all potential variables other than previous caries experience. In the backward elimination procedure, the results of using a level of significance of 0.05 or 0.10 were identical. In the final model, 3 modifiable factors, daily tooth-brushing frequency, composite water fluoride level, and frequency of snacktime intake of processed starches were significantly associated with having new cavitated caries. SES was also significant in the final model.

Model 2, which initially included all potential variables, including previous caries experience, showed that both previous caries experience variables (noncavitated and cavitated) and daily tooth-brushing frequency were significantly associated with new caries. Processed starch intake frequency at snacking occasions and composite water fluoride level are the 2 variables in the final model that had P-values between .05 and .10. These 2 variables would have been trimmed from the model with a .05 significance level. Including previous caries experience in the model strengthened the association between the age at mixed dentition exam variable and new caries, which was found statistically significant in the final model. Low SES consistently has been found to be related to caries outcome in published studies. In model 2, SES was not significantly associated with caries incidence, probably because SES was highly correlated with caries experience at 5-years-old (already in the model).

Processed starches are common snacks in developed countries. The association between caries incidence and processed starches is not surprising. No studies, however, have found this association in early school-aged children. Possibly there are no other studies that have detailed information on dietary intake at this level. Also, processed starch intake frequency at snacking occasions might be counted with other snack intakes in other studies. Thus, the effect of processed starch exposures on new caries might be part of the association between a combined dietary variable such as frequency of snack intake and new cavitated caries. Developmentally, children in this age group still need to have snacks. Thus, the defined findings of which snacks would increase the chances of having new caries can be very useful for making recommendations.

Daily tooth-brushing frequency is a strong preventive factor in this group of children. This confirmed the finding of other studies.<sup>3,9</sup> Some other studies<sup>4,10</sup> that assessed children's oral hygiene, however, did not find significant associations with caries incidence. In this study, more than 99% of children used fluoridated toothpaste. Thus, the daily toothbrushing frequency was the major factor concerning the effect of fluoride exposure from toothpaste. Daily tooth-brushing frequency is the modifiable factor that had the strongest relationship with new caries.

In model 2, there was a statistically significant interaction effect between gender and composite water fluoride level. Specifically, increased composite water fluoride level showed a significant protective effect in girls, but not in boys. Additional studies are needed to confirm and explain these findings.

Most other studies of caries incidence and diet collected dietary information at only one point (either at the beginning or the end of this period) and related it to the new caries. This study collected the dietary information longitudinally to evaluate the relationship between the dietary intake during the period between the 2 dental examinations and cavitated caries incidence, which is a major strength of the present study. There were limitations to this study, however, including a relatively small sample size, a predominance of subjects with high SES, and a low to moderate disease level in the cohort. Thus, results should be interpreted in light of these limitations. This study did not assess cariogenic bacteria. Thus, the results of this study did not control for differences in the cariogenic bacteria among individuals. Additionally, no analyses considered duration of exposures to foods and beverages.

For practicality, future research should focus more on modifiable factors. Moreover, it should assess risk factors at a detailed level that could lead to practical preventive measures.

# Conclusions

Based on this study's results, the following conclusions can be made:

 Snacktime intake of processed starches, toothbrushing, and socioeconomic status are variables associated with caries, consistent with other studies in preschool and school-age children.

- 2. In school-age children, many of those associations with modifiable factors are absent from multivariable models, which include previous caries experience. Studies concerning risk factors for dental caries should be focused to include and emphasize modifiable factors.
- 3. Results suggested that tooth-brushing with fluoridated toothpaste and dietary counseling could help decrease the risk of new cavitated caries in young school-aged children.

#### Acknowledgments

The Iowa Fluoride Study was supported by NIH grants M01-RR00059, R01-DE09551, and R01-DE12101, CDC grants TS-0652 and TS-1329, and Dr. Levy's Wright-Bush-Shreves Endowed Research Professorship. The contents of this paper are the responsibility of the authors and do not necessarily reflect the official views of the granting organizations.

#### References

- 1. Beltran-Aguilar ED, Barker LK, Canto MT, et al. Surveillance for dental caries, dental sealants, tooth retention, edentulism, and enamel fluorosis: United States, 1988-1994 and 1999-2002. MMWR Surveill Summ 2005; 54:1-43.
- Steiner M, Helfenstein U, Marthaler TM. Dental predictors of high caries increment in children. J Dent Res 1992;71:1926-33.
- 3. Vanobbergen J, Martens L, Lesaffre E, Bogaerts K, Declerck D. The value of a baseline caries risk assessment model in the primary dentition for the prediction of caries incidence in the permanent dentition. Caries Res 2001;35:442-50.
- Tagliaferro EP, Ambrosano GM, Meneghim Mde C, Pereira AC. Risk indicators and risk predictors of dental caries in schoolchildren. J Appl Oral Sci 2008;16: 408-13.
- 5. Zhang Q, van Palenstein Helderman WH. Caries experience variables as indicators in caries risk assessment in 6- to 7-year-old Chinese children. J Dent 2006;34: 676-81.
- 6. Vallejos-Sanchez AA, Medina-Solis CE, Casanova-Rosado JF, Maupome G, Minaya-Sanchez M, Perez-Olivares S. Caries increment in the permanent dentition of Mexican children in relation, to prior caries experience on permanent and primary dentitions. J Dent 2006;34:709-15.

- 7. Skeie MS, Raadal M, Strand GV, Espelid I. Caries in primary teeth at 5 and 10 years of age: A longitudinal study. Eur J Paediatr Dent 2004;5:194-202.
- 8. Peretz B, Ram D, Azo E, Efrat Y. Preschool caries as an indicator of future caries: A longitudinal study. Pediatr Dent 2003;25:114-8.
- 9. Leroy R, Bogaerts K, Lesaffre E, Declerck D. Multivariate survival analysis for the identification of factors associated with cavity formation in permanent first molars. Eur J Oral Sci 2005;113:145-52.
- 10. Mattila ML, Rautava P, Paunio P, et al. Caries experience and caries increments at 10 years of age. Caries Res 2001; 35:435-41.
- 11. Levy SM, Warren JJ, Broffitt B. Patterns of fluoride intake from 36 to 72 months of age. J Public Health Dent 2003;63:211-20.
- 12. Levy SM, Kiritsy MC, Slager SL, Warren JJ. Patterns of dietary fluoride supplement use during infancy. J Public Health Dent 1998;58:228-33.
- 13. Marshall TA, Broffitt B, Eichenberger-Gilmore J, Warren JJ, Cunningham MA, Levy SM. The roles of meal, snack, and daily total food and beverage exposures on caries experience in young children. J Public Health Dent 2005; 65:166-73.
- 14. Levy SM, Warren JJ, Davis CS, Kirchner HL, Kanellis MJ, Wefel JS. Patterns of fluoride intake from birth to 36 months. J Public Health Dent 2001;61:70-7.
- 15. Franzman MR, Levy SM, Warren JJ, Broffitt B. Toothbrushing and dentifrice use among children ages 6 to 60 months. Pediatr Dent 2004;26:87-92.
- 16. Warren JJ, Levy SM, Kanellis MJ. Dental caries in the primary dentition: Assessing prevalence of cavitated and noncavitated lesions. J Public Health Dent 2002;62: 109-14.
- 17. Pitts NB, Fyffe HE. The effect of varying diagnostic thresholds upon clinical caries data for a low prevalence group. J Dent Res 1988;67:592-6.
- Pitts NB. Diagnostic tools and measurements: Impact on appropriate care. Community Dent Oral Epidemiol 1997;25:24-35.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.