

Guideline on Perinatal Oral Health Care

Originating Council

Council on Clinical Affairs

Review Council

Council on Clinical Affairs

Adopted

2009

Revised

2011

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that perinatal oral health, along with infant oral health, is one of the foundations upon which preventive education and dental care must be built to enhance the opportunity for a child to have a lifetime free from preventable oral disease. Recognizing that dentists, physicians, allied health professionals, and community organizations must be involved as partners to achieve this goal, the AAPD proposes recommendations for perinatal oral health care, including caries risk assessment, anticipatory guidance, preventive strategies, and therapeutic interventions, to be followed by the stakeholders in perinatal and pediatric oral health.

Method

This document is an update of the previous guideline adopted in 2009. The update is based on a review of the current dental and medical literature related to perinatal oral health care. An electronic search was conducted using PubMed® with the following parameters: Terms: "early childhood caries", "perinatal", "perinatal oral health", and "early childhood caries prevention"; Fields: all; Limits: within the last 10 years, humans, English, and clinical trials. Papers for review were chosen from the resultant list of 113 articles and from references within selected articles. When data did not appear sufficient or were inconclusive, recommendations were based upon expert and/or consensus opinion by experienced researchers and clinicians.

Background

The perinatal period is defined as the period around the time of birth, beginning with the completion of the 20-28th week of gestation and ending 1-4 weeks after birth.¹⁻³ Perinatal oral health plays a crucial role in the overall health and well-being of pregnant women.⁴ It is also essential for the health and well-being of their newborn children. Many women do not seek dental care during their pregnancy, and those that do often confront unwillingness of dentists to provide care.⁵⁻⁹ Many expectant mothers are unaware of the implications of poor oral health for themselves, their pregnancy, and/or their unborn child.^{7,10,11} Research continues to show links between peri-

odontal disease and adverse outcomes in pregnancy including preterm deliveries, low birth weight babies, and preeclampsia.¹²⁻²¹ Furthermore, mothers with poor oral health and high levels of cariogenic oral bacteria are at greater risk for infecting their children with the bacteria and increasing their children's caries risk at an early age.²² Dental caries in infants is a disease that generally is preventable. Determining those mothers at highest risk for transferring cariogenic bacteria to their children improves opportunities for preventive intervention.

The primary goal of perinatal oral health care, with regard to caries transmission, is to lower the numbers of cariogenic bacteria in an expectant mother's mouth so that mutans streptococci (MS) colonization of the infant can be delayed as long as possible.²³ Timely delivery of educational information and preventive therapies to these parents can reduce the incidence of early childhood caries (ECC), prevent the need for dental rehabilitation, and improve the oral health of their children.²⁴⁻²⁶ Physicians, nurses, and other health care professionals are far more likely to see expectant or new mothers and their infants than are dentists. Therefore, it is essential that these providers be aware of the infectious etiology and associated risk factors of dental caries and ECC, make appropriate decisions regarding timely and effective interventions for pregnant women, and facilitate the establishment of a dental home.²⁷⁻²⁹

Perinatal Oral Health

Dental caries

Dental caries is a common chronic infectious transmissible disease resulting from tooth-adherent specific bacteria, primarily MS, that metabolize sugars to produce acid which, over time, demineralizes tooth structure.³⁰ MS is considered to be the principal indicator group of bacterial organisms responsible for dental caries.^{31,32} MS colonization of an infant may occur from the time of birth.^{31,33-38} Significant colonization occurs after dental eruption as teeth provide non-shedding surfaces for adherence. Other surfaces also may harbor MS.³⁸⁻⁴⁰ For example, the furrows of the tongue appear to be an important ecological niche in harboring the bacteria in predentate infants.^{31,39}

Vertical transmission of MS from mother to infant is well documented.⁴¹⁻⁴⁴ Genotypes of MS in infants appear identical to those present in mothers in 17 reports, ranging from 24 to 100%.⁴³ The higher the levels of maternal salivary MS, the greater the risk of the infant being colonized.⁴⁵⁻⁴⁷ The success of transmission and resulting colonization of the mother's salivary MS may be related to several factors including the magnitude of the inoculum, frequency of inoculation, and a minimum infective dose.⁴⁸⁻⁵⁰ Along with maternal salivary levels of MS, the mother's oral hygiene, periodontal disease, snack frequency, and socioeconomic status also are associated with infant colonization.⁴⁰ Reports indicate that horizontal transmission (ie, transmission between members of a group such as siblings of a similar age or children in a daycare center) occurs.⁵¹⁻⁵⁴ A study of children with severe early childhood caries (S-ECC) revealed that non-maternal MS genotypes were identified in the majority (74%) of children.

Caries risk assessment can be performed to determine the patient's relative risk for dental caries. Its goal is to prevent disease by identifying and minimizing causative factors (eg, microbial burden, dietary habits, plaque accumulation) and optimizing protective factors (eg, fluoride exposure, oral hygiene, sealants).⁵⁵ Caries-risk assessment can aid in the identification of reliable predictors and allow health care professionals to identify and refer high-risk patients.⁵⁶ The early establishment of a dental home provides time-critical opportunities to implement preventive oral health practices, including caries risk assessment, and reduces the risk of preventable oral/dental conditions or disease for children.⁵⁷

Anticipatory guidance⁴⁵

Anticipatory guidance for the mother and/or other caregivers can help delay the onset and reduce the impact of MS colonization of the infant.^{24-26,58} Modification of the mother's oral hygiene and diet and the use of topical chlorhexidine and/or fluoride can have a significant effect on MS levels and, correspondingly, the infant's caries rate.⁵⁹⁻⁶²

Recommendations

Recommendations for perinatal oral health include the following:

Oral health education: The perinatal period is an opportune time to educate and perform dental treatment on expectant mothers.^{9,19,63,64} Pregnancy offers an opportunity to educate women regarding oral health by providing a "teachable moment" in self-care and future child-care.⁶⁵ Early intervention and counseling during the perinatal period from all health care providers (eg, physicians, dentists, nurses) are essential to ensure good oral health for the mother and infant.²⁴⁻²⁶

Oral hygiene: Toothbrushing with fluoridated toothpaste and flossing by the mother are important to help dislodge food and reduce bacterial plaque levels. Systematic literature reviews suggest an association between periodontal disease and an increased risk of adverse pregnancy outcomes, including preterm deliveries, low birth weight babies, and preeclampsia.^{14,15,17,18,66,67} Periodontal infections, which can be a reservoir for inflamma-

tory mediators, can pose a threat to the placenta and fetus which can increase the likelihood of preterm delivery.^{68,69} Mothers with severe periodontitis have high levels of prostaglandin in their gingival crevicular fluid and blood. In turn, these increased levels of prostaglandins may be associated with uterine contractions leading to preterm deliveries.^{69,70} While some research shows that scaling and root planning during pregnancy can reduce the likelihood of preterm deliveries and low birth weight babies,^{12,71-75} a recent randomized controlled trial did not support that treatment of periodontal disease during pregnancy prevents preterm birth, fetal growth restriction, or preeclampsia.⁷⁶ Regardless of the potential for improved oral health to improve pregnancy outcomes, the data on the relationship between maternal and child experience with dental caries is well established. Therefore, comprehensive dental services for pregnant women should be available so that, not only their own oral and general health is safeguarded, but also so that their children's caries risk is reduced.⁶⁵

The effects of pregnancy may negatively affect oral health behaviors among pregnant women. Nausea and vomiting may lead to avoidance of toothbrushing, resulting in an increased caries rate. For a pregnant woman experiencing frequent vomiting, rinsing with a cup of water containing a teaspoon of baking soda and waiting an hour before brushing can help minimize dental erosion.⁹ Using a fluoridated toothpaste, chewing sugarless or xylitol-containing gum, and eating small amounts of nutritious food throughout the day can help minimize their caries risk.⁹

Diet: Important components of the mother's diet need to be discussed fully. A healthy diet is necessary to provide adequate amounts of nutrients for the mother-to-be and unborn child. Food cravings may lead to the consumption of foods that increase the mother's caries risk. The caries potential of the mother's diet (ie, cariogenicity of certain foods, beverages, medicines), as well as its effect on her child, should be addressed. The frequency of consumption of cariogenic substances and resulting demineralization/remineralization process also are important discussion topics.

Fluoride: Using a fluoridated toothpaste and rinsing with an alcohol-free, over-the-counter mouth rinse containing 0.05% sodium fluoride once a day or 0.02% sodium fluoride rinse twice a day have been suggested to help reduce plaque levels and help promote enamel remineralization.²³

Professional oral health care: Routine professional dental care for the mother can help optimize her oral health. Every pregnant woman should have an oral evaluation, be counseled on proper oral hygiene, and be referred for preventive and therapeutic oral health care.^{9,19} It has been shown that maternal salivary levels of MS are related significantly to MS colonization in plaque as well as dental caries in their children.^{46,77,78} Removal of active caries with subsequent restoration of the remaining tooth structure is important to suppress maternal MS reservoirs and has the potential to minimize the transfer of MS to the infant, thereby decreasing the infant's risk of developing ECC.⁹ The safest time to perform dental treatment during

pregnancy is in the second trimester, or the 14th–20th weeks.^{9,19,65,79} The risk of pregnancy loss is lower in the second trimester compared to that in the first trimester, and organogenesis is complete.⁷ Even though the second trimester is usually optimal, dental treatment can be accomplished safely at any time during pregnancy.⁹

Treatment options may include diagnostic x-rays,^{9,80–84} dental prophylaxis, periodontal treatment, and restorations^{85–90} with the administration of local anesthetics containing epinephrine.^{9,91} Amalgam may be considered as a restorative material in pregnant women. There is no evidence that fetal exposure to mercury released from the mother's existing amalgam restorations causes any adverse effects.^{85,86,89,90} Since mercury vapor released during removal and placement of an amalgam restoration may be absorbed into the blood stream and cross the placental barrier, the use of rubber dam and high speed evacuation is recommended.⁹⁰ Antibiotics and analgesics for treating infection and controlling pain may be administered.⁹ Acute conditions, such as pain and swelling, should be treated as soon as possible.^{9,92} Delay in necessary treatment could result in significant risk to the mother and indirectly to the fetus.⁹ The consequences of not treating an active infection during pregnancy out-weigh the possible risks presented by most of the medications required for dental treatment.⁹³ Due to patient positioning, comfort is a consideration for treatment during the third trimester. In these cases, elective treatment sometimes is best deferred until after delivery.

Delay of colonization: Reducing maternal MS reservoirs, avoiding or delaying MS transmission, and implementing preventive practices for the child can help delay the colonization process.^{94–96} Maternal MS reservoirs can be suppressed by dietary counseling, reducing the frequency of simple carbohydrate intake, applying topical chlorhexidine and/or fluoride, removing and restoring active caries, and chewing xylitol-containing chewing gum.⁹ Evidence suggests that the use of xylitol chewing gum (at least 2 or 3 times per day by the mother) has a significant impact on mother-child transmission of MS and decreasing the child's caries rate.^{61,97–100} Avoidance or delay of MS transmission can be accomplished by educating the mother or caregiver on behaviors that directly pass saliva to the child (eg, sharing utensils or cups, cleaning a dropped pacifier by mouth).^{94,101,102} Routine preventive efforts should include toothbrushing, optimizing the child's fluoride exposure, and limiting the child's frequency of carbohydrate intake.

Additional recommendations

The AAPD recommends:

1. All primary health care professionals who serve pregnant women provide education on the etiology and prevention of ECC. Oral health counseling and referral for a comprehensive oral examination and treatment during pregnancy is especially important for the mother.
2. The curriculum of all medical, nursing, and allied health professional programs include education in perinatal oral health, including the infectious etiology of ECC, methods of oral health risk assessment, anticipatory guidance, and the need for early establishment of a dental home.
3. Parents/caregivers establish a dental home for infants by 12 months of age.
4. Legislators, policy makers and third party payors be educated about the benefits of perinatal intervention in order to support efforts that improve access to oral health care for pregnant women, including more frequent and comprehensive services.

References

1. Datasegment.com. Online dictionary. Available at: "<http://onlinedictionary.datasegment.com/word/perinatal>". Accessed November 27, 2010.
2. Merriam-Webster Online Dictionary. Available at: "<http://www.merriam-webster.com/dictionary/perinatal>". Accessed November 27, 2010.
3. The Free Encyclopedia Wikimedia Foundation Inc. Wikipedia. Available at: "<http://www.en.wikipedia.org/wiki/Perinatal>". Accessed November 27, 2010.
4. Brown A. Access for Oral Health Care During the Perinatal Period: A Policy Brief. National Maternal and Child Oral Health Resource Center. Georgetown University, Washington, DC; 2008. Available at: "<http://www.mchoralhealth.org/PDFs/PerinatalBrief.pdf>". Accessed November 27, 2010.
5. Gaffield ML, Gilbert BJ, Malvitz DM. Oral Health during pregnancy: An analysis of information collected by the pregnancy risk assessment monitoring system. *J Am Dent Assoc* 2001;132(7):1009-16.
6. Huebner CE, Milgrom P, Conrad D, Lee RS. Providing dental care to pregnant patients: A survey of Oregon general dentists. *J Am Dent Assoc* 2009;140(2):211-22.
7. Keirse MJ, Plutzer K. Women's attitudes to and perceptions of oral health and dental care during pregnancy. *J Perinat Med* 2010;38(1):3-8.
8. Kerpen SJ, Burakoff R. Improving access to oral health care for pregnant women. A private practice model. *NY State J* 2009;75(6):34-6.
9. New York State Department of Health. Oral health care during pregnancy and early childhood: Practice Guidelines. August, 2006. Available at: "<http://www.health.state.ny.us/publications/0824.pdf>". Accessed November 27, 2010.
10. Dimitrova MM. A study of pregnant women's knowledge of children's feeding practice as a risk factor for early childhood caries. *Folia Med (Plovdiv)* 2009;51(4):40-5.
11. Fadavi S, Sevandal MC, Koerber A, Punwani I. Survey of oral health knowledge and behavior of pregnant minority adolescents. *Pediatr Dent* 2009;31(5):405-8.
12. Dasanayake AP, Gennaro S, Hendricks-Muñoz KD, Chhun N. Maternal periodontal disease, pregnancy, and neonatal outcomes. *MCN Am J Matern Child Nurs* 2008;33(1):45-9.

13. Oettinger-Barak O, Barak S, Ohel G, et al. Severe pregnancy complication (preeclampsia) is associated with greater periodontal destruction. *J Periodontol* 2005;76(1):134-7.
14. Pitiphat W, Joshipura KJ, Gillman MW, Williams PL, Douglass CW, Rich-Edwards JW. Maternal periodontitis and adverse pregnancy outcomes. *Community Dent Oral Epidemiol* 2008;36(1):3-11.
15. Ruma M, Boggess K, Moss K, et al. Maternal periodontal disease, systemic inflammation, and risk for preeclampsia. *Am J Obstet Gynecol* 2008;198(4):389.e1-5.
16. Sacco G, Carmagnola D, Abati S, et al. Periodontal disease and preterm birth relationship: A review of the literature. *Minerva Stomatol* 2008;57(5):233-50.
17. Shetty M, Shetty PK, Ramesh A, Thomas B, Prabhu S, Rao A. Periodontal disease in pregnancy is a risk factor for preeclampsia. *Acta Obstet Gynecol Scand* 2010;89(5):718-21.
18. Shub A, Wong C, Jennings B, Swain JR, Newnham JP. Maternal periodontal disease and perinatal mortality. *Aust N Z J Obstet Gynaecol* 2009;49(2):130-6.
19. Silk H, Douglass AB, Douglass JM, Silk L. Oral health during pregnancy. *Am Fam Physician* 2008;77(8):1139-44.
20. Siqueira FM, Cota LO, Costa JE, Haddad JP, Lana AM, Costa FO. Maternal periodontitis as a potential risk variable for preeclampsia: A case-control study. *J Periodontol* 2008;79(2):207-15.
21. Xiong X, Buekens P, Fraser WD, Beck J, Offenbacher S. Periodontal disease and adverse pregnancy outcomes: A systematic review. *BJOG* 2006;113(2):135-43.
22. Ramos-Gomez FJ, Weintraub JA, Gansky SA, Hoover CI, Featherstone JD. Bacterial, behavioral and environmental factors associated with early childhood caries. *J Clin Pediatr Dent* 2002;26(2):165-73.
23. American Academy of Pediatrics. Policy on oral health risk assessment timing and establishment of the dental home. *Pediatrics* 2003;111(5Pt1):1113-6.
24. Lucey SM. Oral health promotion initiated during pregnancy successful in reducing early childhood caries. *Evid Based Dent* 2009;10(4):100-1.
25. Meyer K, Geurtsen W, Gunay H. An early oral health care program starting during pregnancy: Results of a prospective clinical long-term study. *Clin Oral Investig* 2010;14(3):257-64.
26. Plutzer K, Spencer AJ. Efficacy of an oral health promotion intervention in the prevention of early childhood caries. *Community Dent Oral Epidemiol* 2008;36(4):335-46.
27. Harrison R. Oral health promotion for high-risk children: Case studies from British Columbia. *J Can Dent Assoc* 2003;69(5):292-6.
28. Lewis CW, Grossman DC, Domoto PK, Deyo RA. The role of the pediatrician in the oral health of children: A national survey. *Pediatrics* 2000;106(6):E84.
29. Nowak AJ, Warren JJ. Infant oral health and oral habits. *Pediatr Clin North Am* 2000;47(5):1043-66.
30. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev* 1986;50(4):353-80.
31. Berkowitz RJ. Mutans streptococci: Acquisition and transmission. *Pediatr Dent* 2006;28(2):106-9; discussion 192-8.
32. Ge Y, Caufield PW, Fisch GS, Li Y. *Streptococcus mutans* and *Streptococcus sanguinis* colonization correlated with caries experience in children. *Caries Res* 2008;42(6):444-8.
33. Berkowitz RJ, Jordan HV, White G. The early establishment of *Streptococcus mutans* in the mouth of infants. *Arch Oral Biol* 1975;20(3):171-4.
34. Law V, Seow WK, Townsend G. Factors influencing oral colonization of mutans streptococci in young children. *Aust Dent J* 2007;52(2):93-100; quiz 159.
35. Loesche WJ. Microbial adhesion and plaque. In: *Dental Caries: A Treatable Infection*. 2nd ed. Grand Haven, Mich. Automated Diagnostic Documentation, Inc; 1993:81-116.
36. Stiles HM, Meyers R, Brunelle JA, Wittig AB. Occurrence of *Streptococcus mutans* and *Streptococcus sanguis* in the oral cavity and feces of young children. In: Stiles M, Loesch WJ, O'Brien T, eds. *Microbial Aspects of Dental Caries*. Washington, DC: Information Retrieval; 1976:187.
37. Wan AK, Seow WK, Purdie DM, Bird PS, Walsh LJ, Tudehope DI. A longitudinal study of *Streptococcus mutans* colonization in infants after tooth eruption. *J Dent Res* 2003;82(7):504-8.
38. Wan AK, Seow WK, Walsh LJ, Bird P, Tudehope DI, Purdie DM. Association of *Streptococcus mutans* infection and oral developmental nodules in pre-dentate infants. *J Dent Res* 2001;80(10):1945-8.
39. Tanner AC, Milgrom PM, Kent R Jr, et al. The microbiota of young children from tooth and tongue samples. *J Dent Res* 2002;81(1):53-7.
40. Wan AK, Seow WK, Purdie DM, Bird PS, Walsh LJ, Tudehope DI. Oral colonization of *Streptococcus mutans* in six-month-old predentate infants. *J Dent Res* 2001;80(12):2060-5.
41. Berkowitz R, Jones P. Mouth-to-mouth transmission of the bacterium *Streptococcus mutans* between mother and child. *Arch Oral Biol* 1985;30(4):377-9.
42. Davey AL, Rogers AH. Multiple types of the bacterium *Streptococcus mutans* in the human mouth and their intra-family transmission. *Arch Oral Biol* 1984;29(6):453-60.
43. Douglass JM, Li Y, Tinanoff N. Association of Mutans streptococci between caregivers and their children. *Pediatr Dent* 2008;29(5):375-87.
44. Li Y, Caufield PW. The fidelity of initial acquisition of Mutans streptococci by infants from their mothers. *J Dent Res* 1995;74(2):681-5.
45. American Academy of Pediatric Dentistry. Policy on early childhood caries (ECC): Classifications, consequences, and prevention strategies. *Pediatr Dent* 2011;33(special issue):47-9.

46. Kishi M, Abe A, Kishi K, Ohara-Nemoto Y, Kimura S, Yonemitsu M. Relationship of quantitative salivary levels of *Streptococcus mutans* and *Streptococcus sobrinus* in mothers to caries status and colonization of mutans streptococci in plaque in their 2.5-year-old children. *Community Dent Oral Epidemiol* 2009;37(3):241-9.
47. Parisotto TM, Steiner-Oliveira C, Silva CM, Rodrigues LK, Nobre-dos-Santos M. Early childhood caries and mutans streptococci: A systematic review. *Oral Health Prev Dent* 2010;8(1):59-70.
48. Acs G, Lodolini G, Kaminshy S, Cisneros GJ. Effect of nursing caries on body weight in pediatric populations. *Pediatr Dent* 1992;14(5):302-5.
49. Clarke M, Locker D, Berall G, Pencharz P, Kenny DJ, Judd P. Malnourishment in a population of young children with severe early childhood caries. *Pediatr Dent* 2006;28(3):254-9.
50. Ismail AI. Prevention of early childhood caries. *Comm Dent Oral Epidemiol* 1998;26(suppl):49-61.
51. Mattos-Graner RO, Li Y, Caufield PW, Duncan M, Smith JD. Genotypic diversity of Mutans streptococci in Brazilian nursery children suggests horizontal transmission. *J Clin Microbiol* 2001;39(6):2313-6.
52. Mitchell SC, Ruby JD, Moser S, et al. Maternal transmission of mutans streptococci in severe-early childhood caries. *Pediatr Dent* 2009;31(3):193-201.
53. Redmo Emanuelsson IM, Wang XM. Demonstration of identical strains of mutans streptococci within Chinese families by genotyping. *Eur J Oral Sci* 1998;106(3):778-94.
54. van Loveren C, Buijs JF, ten Cate JM. Similarity of bacteriocin activity profiles of mutans streptococci within the family when the children acquire strains after the age of 5. *Caries Res* 2000;34(6):481-5.
55. American Academy of Pediatric Dentistry. Guideline on periodicity of examination, preventive dental services, anticipatory guidance, and oral treatment for infants, children, and adolescents. *Pediatr Dent* 2010;32(special issue):93-100.
56. American Academy of Pediatric Dentistry. Guideline on caries-risk assessment and management for infants, children, and adolescents. *Pediatr Dent* 2011;33(special issue):110-7.
57. American Academy of Pediatric Dentistry. Policy on the dental home. *Pediatr Dent* 2010;32(special issue):25-6.
58. Nowak AJ, Casamassimo PS. Using anticipatory guidance to provide early dental intervention. *J Am Dent Assoc* 1995;126(8):1156-63.
59. Brambilla E, Felloni A, Gagliani M, Malerba A, García-Godoy F, Strohmer L. Caries prevention during pregnancy: Results of a 30-month study. *J Am Dent Assoc* 1998;129(7):871-7.
60. Ercan E, Dulgergil CT, Yildirim I, Dalli M. Prevention of maternal bacterial transmission on children's dental caries development; 4-year results of a pilot study in a rural-child population. *Arch Oral Biol* 2007;52(8):748-52.
61. Isokangas P, Söderling E, Pienihäkkinen K, Alanen P. Occurrence of dental decay in children after maternal consumption of xylitol chewing gum: A follow-up from 0 to 5 years of age. *J Dent Res* 2000;79(11):1885-9.
62. Köhler B, Andréen I, Jonsson B. The effects of caries-preventive measures in mothers on dental caries and the oral presence of the bacteria *Streptococcus mutans* and lactobacilli in their children. *Arch Oral Biol* 1984;29(11):879-83.
63. Boggess KA, Society for Maternal-Fetal Medicine Publications Committee. Maternal oral health in pregnancy. *Obstet Gynecol* 2008;111(4):976-86.
64. dela Cruz GG, Rozier RG, Slade G. Dental screening and referral of young children by pediatric primary care providers. *Pediatrics* 2004;114(5):e642-52.
65. Boggess KA, Edelstein BL. Oral health in women during preconception and pregnancy: Implications for birth outcomes and infant oral health. *Matern Child Health J* 2006;10(5 suppl):S169-74.
66. Khader YS, Ta'ani Q. Periodontal diseases and the risk of preterm birth and low birth weight: A meta-analysis. *J Periodontol* 2005;76(2):161-5.
67. Xiong X, Buekens P, Vastardis S, Yu SM. Periodontal disease and pregnancy outcomes: State-of-the-science. *Obstet Gynecol Surv* 2007;62(9):605-15.
68. Jeffcoat MK, Geurs NC, Reddy MS, Cliver SP, Goldenberg RL, Hauth JC. Periodontal infection and preterm birth: Results of a prospective survey. *J Am Dent Assoc* 2001;132(7):875-80.
69. Offenbacher S, Katz V, Fertik G, et al. Periodontal infections as a possible risk factor for preterm low birth weight. *J Periodontol* 1996;67(suppl 10):1103-13.
70. Offenbacher S, Jared HL, O'Reilly PG, et al. Potential pathogenic mechanisms of periodontitis associated pregnancy complications. *Ann Periodontol* 1998;3(1):233-50.
71. Dasanayake AP, Li Y, Wiener H, Ruby JD, Lee, MJ. Salivary *actinomyces naeslundii* genospecies 2 and *lactobacillus casei* levels predict pregnancy outcomes. *J Periodontol* 2005;76(2):171-7.
72. Jeffcoat MK, Hauth JC, Geurs NC, et al. Periodontal disease and preterm birth: Results of a pilot intervention study. *J Periodontol* 2003;74(8):1214-8.
73. López NJ, Smith PC, Gutiérrez J. Periodontal therapy may reduce the risk of preterm low birth weight in women with periodontal disease: A randomized controlled trial. *J Periodontol* 2002;73(8):911-24.
74. López NJ, Da Silva I, Ipinza J, Gutiérrez J. Periodontal therapy reduces the rate of preterm low birth weight in women with pregnancy-associated gingivitis. *J Periodontol* 2005;76(11 suppl):2144-53.
75. Scannapieco FA, Bush RB, Paju S. Periodontal disease as a risk factor for adverse pregnancy outcomes. A systemic review. *Ann Periodontol* 2003;8(1):70-8.
76. Niederman R. Periodontal treatment did not prevent complications of pregnancy. *Evid Based Dent* 2010;11(1):18-9.

77. Al Shukairy H, Alamoudi N, Farsi N, Al Mushayt A, Masoud I. A comparative study of *Streptococcus mutans* and lactobacilli in mothers and children with severe early childhood caries (SECC) versus a caries free group of children and their corresponding mothers. *J Clin Pediatr Dent* 2006;31(2):80-5.
78. Irigoyen Camacho ME, Sánchez Pérez L, García Pérez A, Zepeda Zepeda MA. Relationship between severe early childhood caries, mother's oral health and mutans streptococci in a low-income group: Changes from 1996 to 2007. *J Clin Pediatr Dent* 2009;33(3):241-6.
79. Gajendra S, Kumar JV. Oral health and pregnancy: A review. *NY State Dent J* 2004;70(1):40-4.
80. American College of Obstetrics and Gynecology. Committee Opinion, Number 299, September 2004 Guidelines for diagnostic imaging during pregnancy. *Obstet Gynecol* 2004;104(3):647-51.
81. American Dental Association, US Dept of Health and Human Services. The selection of patients for x-ray examinations: Dental radiographic examinations. Rockville, Md. Food and Drug Administration, 2004; HHS Publication Number 88-8273. Available at: "http://www.ada.org/sections/professionalResources/pdfs/topics_radiography_examinations.pdf". Accessed November 27, 2010.
82. National Council on Radiation Protection and Measurement. Report No. 54: Medical radiation exposure of pregnant and potentially pregnant women. Bethesda, Md; 1977.
83. Matteson SR, Joseph LP, Bottomley W, et al. The report of the panel to develop radiographic selection criteria for dental patients. *Gen Dent* 1991;39(4):264-70.
84. Toppenberg KS, Hill DA, Miller DP. Safety of radiographic imaging during pregnancy. *Am Fam Physician* 1999;59(7):1813-8, 1820.
85. Hujoel PP, Lydon-Rochelle M, Bollen AM, Woods JS, Geurtsen W, del Aguila MA. Mercury exposure from dental filling replacement during pregnancy and low birth weight risk. *Am J Epidemiol* 2005;161(8):734-40.
86. Life Sciences Research Office. Review and analysis of the literature on the potential adverse effects of dental amalgam. Bethesda, Md. December 9, 2004. Available at: "http://www.lbro.org/amalgam/frames_amalgam_report.html". November 27, 2010.
87. March of Dimes. During your pregnancy: Things to avoid: Mercury. 2008. Available at: "http://www.marchofdimes.com/pnhec/159_15759.asp". Accessed November 27, 2010.
88. Olea N, Pulgar R, Perez P, et al. Estrogenicity of resin-based composites and sealants used in dentistry. *Environ Health Perspect* 1996;104(3):298-305.
89. US Food and Drug Administration. White paper: FDA update/review of potential adverse health risks associated with exposure to mercury in dental amalgam. Available at: "<http://www.fda.gov/MedicalDevices/ProductsandMedicalProcedures/DentalProducts/DentalAmalgam/ucm171117.htm>". November 27, 2010.
90. Whittle KW, Whittle JG, Sarll DW. Amalgam fillings during pregnancy. *Br Dent J* 1998;185(10):500.
91. Rosen MA. Management of anesthesia for the pregnant surgical patient. *Anesthesiol* 1999;91(4):1159-63.
92. Chiodo GT, Rosenstein DI. Dental treatment during pregnancy: A preventive approach. *J Am Dent Assoc* 1985;110(3):365-8.
93. Moore PA. Selecting drugs for the dental patient. *J Am Dent Assoc* 1998;129(9):1281-6.
94. Berkowitz RJ. Causes, treatments and prevention of early childhood caries: A microbiologic perspective. *J Can Dent Assoc* 2003;69(5):304-7.
95. Kohler B, Andreen I. Influence of caries-preventive measures in mothers on cariogenic bacteria and caries experience in their children. *Arch Oral Biol* 1994;39(10):907-11.
96. Li Y, Dasanayake AP, Caufield PW, Elliot RR, Butts JT III. Characterization of maternal mutans streptococci transmission in an African American population. *Dent Clin North Am* 2003;47(1):87-101.
97. Gomez SS, Weber AA. Effectiveness of a caries preventive program in pregnant women and new mothers on their offspring. *Int J Paediatr Dent* 2001;11(2):117-22.
98. Söderling E, Isokongas P, Pienihäkkinen K, Tenovou J. Influence of maternal xylitol consumption on acquisition of mutans streptococci by infants. *J Dent Res* 2000;79(3):882-7.
99. Söderling E, Isokongas P, Pienihäkkinen K, Tenovou J, Alanen P. Influence of maternal xylitol consumption on mother-child transmission of Mutans streptococci: 6-year follow-up. *Caries Res* 2001;35(3):173-7.
100. Thorild I, Lindau B, Twetman S. Caries in 4-year-old children after maternal chewing of gums containing combinations of xylitol, sorbitol, chlorhexidine, and fluoride. *Eur Arch Paediatr Dent* 2006;7(4):241-5.
101. Berkowitz RJ. Acquisition and transmission of Mutans streptococci. *J Calif Dent Assoc* 2003;31(2):135-8.
102. Caufield PW, Wannemuehler YM, Hansen JB. Familial clustering of *Streptococcus mutans* cryptic plasmid strain in a dental clinic population. *Infect Immun* 1982;38(2):907-11.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.