Guideline on Xylitol Use in Caries Prevention

Originating Committee

Council on Clinical Affairs

Adopted 2011

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes the benefits of caries preventive strategies involving sugar substitutes, particularly xylitol, on the oral health of infants, children, adolescents, and persons with special health care needs. This guideline is intended to assist oral health care professionals make informed decisions about the use of xylitolbased products in caries prevention.

Methods

This guideline is based upon a review of current dental and medical literature related to the use of xylitol in caries prevention. An electronic search was conducted using PubMed[®] with the following parameters: Terms: "xylitol AND dental caries", "caries prevention", "plaque reduction", "maternal *Streptococcus mutans* transmission", and "*Streptococcus mutans* long term suppression with xylitol"; Fields: all; Limits: within the last 10 years, humans, English, birth through 18. Two hundred forty articles matched these criteria. Fifty-one papers were chosen for review from this list and from the references within selected articles. When data did not appear sufficient or were inconclusive, recommendations were based upon expert and/or consensus opinion by experienced researchers and clinicians.

Background

Xylitol is a naturally occurring 5-carbon sugar polyol currently approved for use in foods, pharmaceuticals, and oral health products in more than 35 countries. It is found naturally in various trees, fruits, and vegetables and is an intermediate product of the glucose metabolic pathway in man and animals.¹ Xylitol was approved by the Food and Drug Administration as a dietary food additive in 1963 and has been used widely in the general market since the mid 1970s. European countries such as Finland have national programs promoting the use of xylitol chewing gum among children in an effort to reduce dental caries. The AAPD supports the use of xylitol in caries prevention.²

Xylitol has properties that reduce levels of mutans streptococci (MS) in the plaque and saliva. Xylitol disrupts the energy production processes of MS leading to a futile energy consumption cycle and cell death.³ Further, consumers of clinically effective levels of xylitol show MS strains with reduced adhesion to the teeth and other reduced virulence properties such as less acid production.⁴⁻⁸ There are numerous clinical studies evaluating the effectiveness of xylitol.⁹⁻³⁷ Several studies of children who have consumed xylitol for 3 weeks or more have reported both short- and long-term reduction in salivary and plaque MS levels.⁹⁻¹⁵ A few studies, however, have not shown a long-term reduction in salivary and plaque MS levels.³⁹⁻⁴² The mechanical action of chewing a gum containing xylitol along with subsequent increased volume of saliva may assist with caries reduction.³⁸ Evaluation for this guideline was done with the consideration that several of the published studies used "no chewing" groups instead of placebo controls.

Numerous clinical studies have demonstrated a decrease in caries rates, increment, and/or onset among children who were exposed to daily xylitol use for 12 to 40 months.¹⁶⁻²⁵ Long-term benefits on caries rates, increment, and/or onset also have been observed up to 5 years after the cessation of xylitol intervention.^{26,27} Xylitol works most effectively on teeth that are erupting.²⁷ There is also evidence that maternal consumption of xylitol may reduce the acquisition of MS and dental caries by their children.²⁸⁻³³

Recommendations

Clinicians may consider recommending xylitol use to moderate or high caries-risk patients. Those recommending xylitol should be familiar with the product labeling and recommend age-appropriate products. They should routinely reassess (not less than once every 6 months) a patient for changes in cariesrisk status and adjust recommendations accordingly.

Dosage

There is accumulating evidence that total daily doses of 3 to 8 grams of xylitol are required for a clinical effect with the currently available delivery methods of syrup, chewing gum, and lozenges.^{40,42} Dosing frequency should be a minimum of 2 times a day⁴², not to exceed 8 grams per day. Although tables of clinically effective xylitol containing products have recently been published, the products are continually changing.^{34,40,42}

Modality

Chewing gum has been the predominant modality for xylitol delivery in clinical studies.³⁵ Studies^{24,36} that have utilized xylitol-containing mints and hard candies have shown them to be as effective as xylitol-containing chewing gum. The American Academy of Pediatrics (AAP) does not recommend use of chewing gum, mints, or hard candy by children less than 4 years of age due to the risk of choking. A randomized trial of xylitol syrup (8 g/day) reduced early childhood caries by 50-70 percent in children 15 to 25 months of age.²⁵ Another study showed that gum or lozenges consumed by children at 5 grams total dose per day at about age 10 resulted in 35-60 percent reductions of tooth decay, with no differences between the delivery methods.²⁴ Xylitol containing gummy bears,¹⁴ other confections,³⁷ and even milk⁴³ have been studied as delivery vehicles, but they are neither well established scientifically nor available commercially at present. A pacifier with a pouch containing slow release xylitol in tablet form, not yet available in the US, has shown high salivary xylitol concentrations and may be a potential delivery vehicle for infants.44 Currently, xylitol-containing chewing gum, mints, energy bars and foods, nasal sprays, and oral hygiene products (eg, mouth rinse, gels, wipes, floss) are commercially available through retail or online venues. However, they may not contain the necessary therapeutic level, xylitol as the only sweetener, or adequate labeling.45

Studies⁴⁶⁻⁴⁸ using toothpaste formulations with 10% xylitol (dose of 0.1 g/brushing) have shown reduction in MS levels and caries in children. The toothpastes that were studied are not for sale in the US. Furthermore, the xylitol-containing toothpastes that currently are sold in the US have never been tested and their formulas differ from those tested.

Current evidence supports the following recommendations for children at moderate or high caries risk:

Age	Xylitol Product	Dosage
< 4 years old	Xylitol syrup*	3 – 8 grams/day in divided doses
≥ 4 years old	Age appropriate products such as chewing gum*, mints, lozenges, snack foods such as gummy bears.	3 – 8 grams/day in divided doses

* AAP does not recommend chewing gum use in children less than 4 years of age due to the risk of choking.⁴⁹

Side effects

Parents need to control the amount of xylitol and other polyols that their child consumes. Xylitol is safe for children when consumed in therapeutic doses for dental caries prevention. Common side effects that may occur with the use of xylitol are gas and osmotic diarrhea. These symptoms usually occur at higher dosages⁵⁰⁻⁵¹ and will subside once xylitol consumption is stopped.⁵¹ To minimize gas and diarrhea, xylitol should be introduced slowly, over a week or more, to acclimate the body to the polyol, especially in young children.

References

- 1. Bär A. Caries prevention with xylitol. A review of the scientific evidence. World Rev Nutr Diet 1988;55:183-209.
- 2. American Academy of Pediatric Dentistry. Policy on the use of xylitol in caries prevention. Pediatr Dent 2010;32 (special issue):36-8.
- 3. Trahan L, NéronS, Bareil M. Intracellular xylitolphosphate hydrolysis and efflux of xylitol in *Streptococcus sobrinus*. Oral Microbiol Immunol 1991;6(1):41-50.
- 4. Trahan L, Söderling E, Dréan MF, Chevrier MC, Isokangas P. Effect of xylitol consumption on the plaque-saliva distribution of mutans streptococci and the occurrence and long-term survival of xylitol resistant strains. J Dent Res 1992;71(11):1785-91.
- 5. Söderling E, Trahan L, Tammiala-Salonen T, Häkkinen L. Effects of xylitol, xylitol-sorbitol, and placebo chewing gums on the plaque of habitual xylitol consumers. Eur J Oral Sci 1997;105(2):170-7.
- 6. Roberts MC, Riedy CA, Coldwell SE, et al. How xylitolcontaining products affect cariogenic bacteria. J Am Dent Assoc 2002;133(4):435-41.
- 7. Tanzer JM, Thompson A, Wen ZT, Burne RA. *Streptoco-ccus mutans*: Fructose transport, xylitol resistance, and virulence. J Dent Res 2006;85(4):369-73.
- 8. Trahan L. Xylitol: A review of its action on mutans streptococci and dental plaque: its clinical significance. Int Dent J 1995;45(1 Suppl 1):77-92.
- 9. Loesche WJ, Grossman NS, Earnest R, Corpron R. The effect of chewing xylitol gum on the plaque and saliva levels of *Streptococcus mutans*. J Am Dent Assoc 1984; 108(4):587-92.
- 10. Autio JT. Effect of xylitol chewing gum on salivary *Streptococcus mutans* in preschool children. ASDC J Dent Child 2002;69(1):81-6.
- 11. Thaweboon S, Thaweboon B, Soo-Ampon S. The effect of xylitol chewing gum on mutans streptococci in saliva and dental plaque. Southeast Asian J Trop Med Public Health 2004; 35(4):1024-7.
- 12. Mäkinen KK, Isotupa KP, Mäkinen PL, et al. Six-month polyol chewing-gum programme in kindergarten-age children: A feasibility study focusing on mutans streptococci and dental plaque. Int Dent J 2005;55(2):81-8.
- 13. Holgerson PL, , Sjöström I, Stecksén-Blicks C, Twetman S. Dental plaque formation and salivary mutans streptococci in school children after use of xylitol-containing chewing gum. Int J Paediatr Dent 2007;17(2):79-85.
- 14. Ly KA, Riedy CA, Milgrom P, Rothen M, Roberts MC, Zhou L. Xylitol gummy bears snacks: A school-based randomized clinical trial. BMC Oral Health 2008;8 (Jul):20.
- 15. Mäkinen KK, Alanen P, Isokangas P, et al. Thirty-nine month xylitol chewing-gum programme in initially 8year-old school children: A feasibility study focusing on mutans streptococci and lactobacilli. Int Dent J 2008;58 (1):41-50.

- Scheinin AK, Pienihäkkien K, Tiekso J, et al. Collaborative WHO xylitol field studies in Hungary. VII. Twoyear caries incidence in 976 institutionalized children. Acta Odontol Scand 1985;43(6):381-7.
- 17. Isokangas P. Xylitol chewing gum in caries prevention: A longitudinal study on Finnish school children. Proc Finn Dent Soc 1987;3(suppl 1):1-117.
- Kandelman D, Bar A, Hefti A. Collaborative WHO xylitol field study in French Polynesia, part I: baseline prevalence and 32 month caries increment. Caries Res 1988;22(1): 55-62.
- 19. Isokangas P, Alanen P, Tiekso J, Mäkinen KK. Xylitol chewing gum in caries prevention: A field study in children. J Am Dent Assoc 1988;117(2):315-20.
- 20. Kandelman D, Gagnon G. Clinical results after 12 months from a study of the incidence and progression of dental caries in relation to consumption of chewing-gum containing xylitol in school preventive programs. J Dent Res 1987;66(8):1407-11.
- 21. Mäkinen KK, Bennett CA, Hujoel PP, et al. Xylitol chewing gums and caries rates: A 40-month cohort study. J Dent Res 1995;74:12:1904-13.
- 22. Mäkinen KK, Mäkinen PL, Pape HR Jr, et al. Conclusion and review of the Michigan Xylitol Programme (1986-1995) for the prevention of dental caries. Int Dent J 1996;46(1):22-34.
- 23. Mäkinen KK, Hujoel PP, Bennett CA, et al. A descriptive report of the effects of a 16-month xylitol chewing-gum programme subsequent to a 40-month sucrose gum programme. Caries Res 1998;32(2):107-12.
- 24. Alanen P, Isokangas P, Gutman K. Xylitol candies in caries prevention: Results of a field study in Estonian children. Community Dent Oral Epidemiol 2000;28(3):218-24.
- 25. Milgrom P, Ly KA, Tut OK, et al. Xylitol pediatric topical oral syrup to prevent dental caries. Arch Pediatr Adolesc Med 2009;163(7):601-7.
- 26. Isokangas P, Mäkinen KK, Tiekso J, Alanen P. Long-term effect of xylitol chewing gum in the prevention of dental caries: A follow-up 5 years after termination of a prevention program. Caries Res 1993;27(6):495-8.
- 27. Hujoel PP, Mäkinen KK, Bennett CA, et al. The optimum time to initiate habitual gum-chewing for obtaining long-term caries prevention. J Dent Res 1999;78(3):797-803.
- Söderling E, Isokangas P, Pienihäkkinen K, Tenovuo J. Influence of maternal xylitol consumption on acquisition of mutans streptococci by infants. J Dent Res 2000;79 (3):882-7.
- 29. Thorild I, Lindau B, Twetman S. Effect of maternal use of chewing gums containing xylitol, chlorhexidine or fluoride on mutans streptococci colonization in the mother's infant children. Oral Health Prev Dent 2003;1(1):53-7.
- 30. Thorild I, Lindau B, Twetman S. Salivary mutans streptococci and dental caries in three-year-old children after maternal exposure to chewing gums containing combi-

nations of xylitol, sorbitol, chlorhexidine and fluoride. Acta Odontol Scand 2004;62(5):245-50.

- 31. Nakai Y, Shinga-Ishihara C, Kaji M, et al. Xylitol gum and maternal transmission of mutans streptococci. J Dent Res 2010;89(1):56-60.
- 32. Isokangas P, Söderling E, Pienihäkkien K, Alanen P. Occurrence of dental decay in children after maternal consumption of xylitol chewing gum, a follow-up from 0 to 5 years of age. J Dent Res 2000;79(11):1885-9.
- 33. Thorild I, Lindau B, Twetman S. Caries in 4-year-old children after maternal chewing of gums containing combinations of xylitol, sorbitol, chlorhexidine and fluoride. Eur Arch Paediatr Dent 2006;7(4):241-5.
- 34. Milgrom P, Ly KA, Rothen M. Xylitol and its vehicles for public health needs. Adv Dent Res 2009;21(1):44-7.
- 35. Ly KA, Milgrom P, Rothen M. The potential of dentalprotective chewing gum in oral health interventions. J Am Dent Assoc 2008;139(5):553-63.
- 36. Honkala E, Honkala S, Shyama M, Al-Mutawa SA. Field trial on caries prevention with xylitol candies among disabled school children. Caries Res 2006;40(6):508-13.
- Lam M, Riedy CA, Coldwell SE, Milgrom P, Craig R. Children's acceptance of xylitol-based foods. Community Dent Oral Epidemiol 2000;28(2):97-101.
- Machiulskiene V, Nyvad B, Baelum V. Caries preventive effect of sugar-substituted chewing gum. Community Dent Oral Epidemiol 2001;29(4):278–88.
- Fontana M, Catt D, Sissons C, et al. Xylitol: Effects on the acquisition of cariogenic species in infants. Pediatr Dent 2009;31(3):257-66.
- 40. Milgrom, P, Zero DT, Tanzer JM. An examination of the advances in science and technology of prevention of tooth decay in young children since the Surgeon General's Report on Oral Health. Acad Pediatr 2009;9(6):404-9.
- Mäkinen KK. An end to crossover designs for studies on the effect of sugar substitutes on caries? Caries Res 2009; 43(5):331-3.
- 42. Deshpande A, Jadad AR. The impact of polyol-containing chewing gums on dental caries: A systematic review of original randomized controlled trials and observational studies. J Am Dent Assoc 2008;139(12):1602-14.
- 43. Castillo JL, Milgrom P, Coldwell SE, Castillo R, Lazo R. Children's acceptance of milk with xylitol or sorbitol for dental caries prevention. BMC Oral Health 2005;5(1):6.
- 44. Taipale T, Pienihäkkien K, Alanen P, Jodela J, Söderling E. Dissolution of xylitol from a food supplement administered with a novel slow-release pacifier: Preliminary results. Eur Arch Paediatr Dent 2007;8(2):123-5.
- 45. Ly KA, Milgrom P, Rothen M. Xylitol, sweeteners, and dental caries. Pediatr Denti 2006;28(2):154-63.
- 46. Sintes JL, Escalante C, Stewart B, et al. Enhanced anticaries efficacy of a 0.243% sodium fluoride/10% xylitol/ silica dentifrice: 3 year clinical results. Am J Dent 1995;8 (5):231-5.

- 47. Sintes JL, Elias-Boneta A, Stewart B, Volpe AR, Lovett J. Anticaries efficacy of sodium monofluorophosphate dentifrice containing xylitol in a dicalcium phosphate dehydrate base. A 30-month caries clinical study in Costa Rica. Am J Dent 2002;15(4):215-9.
- Jannesson L, Renvert S, Kjellsdotter P, Gaffar A, Nabi N, Birkhed D. Effect of a triclosan-containing toothpaste supplemented with 10% xylitol on mutans streptococci in saliva and dental plaque. A 6-month clinical study. Caries Res 2002;36(1):36-9.
- 49. American Academy of Pediatrics. Health issues: Choking prevention. What can I do to keep my child from choking?

In: Caring for Your Baby and Young Child: Birth to Age 5. American Academy of Pediatrics, 2009. Available at: "http://www.healthychildren.org/English/health-issues/ injuries-emergencies/Pages/Choking-Prevention.aspx". Accessed June 10, 2011.

- 50. Förster H, Quadbeck R, Gottstein U. Metabolic tolerance to high doses of oral xylitol in human volunteers not previously adapted to xylitol. Int J Vitam Nutr Res Suppl 1982;22:67-88.
- 51. Akerblom HK, Koivukangas T, Puukka R, Mononen M. The tolerance of increasing amounts of dietary xylitol in children. Int J Vitam Nutr Res Suppl 1982;22:53-66.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.