

# Policy on Early Childhood Caries (ECC): Unique Challenges and Treatment Options

## Originating Council

Council on Clinical Affairs

## Review Council

Council on Clinical Affairs

## Adopted

2000

## Revised

2003, 2007, 2008, 2011

## Purpose

The American Academy of Pediatric Dentistry (AAPD), to promote appropriate, quality oral health care for infants and children with early childhood caries (ECC), must educate the health community and society about the unique challenges and treatment options of this disease, including the need for advanced preventive, restorative, and behavioral guidance techniques.

## Methods

The proceedings of the Conference on Early Childhood Caries held in Bethesda, Maryland, in October 1997<sup>1</sup> were reviewed. The update of this policy used electronic and hand searches of English written articles in the dental and medical literature within the last 10 years using the search terms "infant oral health", "infant oral health care", and "early childhood caries". When data did not appear sufficient or were inconclusive, recommendations were based upon expert and/or consensus opinion by experienced researchers and clinicians.

## Background

ECC is highly prevalent and increasing in poor and near poor US preschool children.<sup>2</sup> In the US and most other countries, this disease is largely untreated in children under age 3.<sup>3</sup> Those children with caries experience have been shown to have high numbers of teeth affected. Consequences of ECC include a higher risk of new carious lesions,<sup>4,5</sup> increased treatment costs and time,<sup>6</sup> risk for delayed physical growth and development,<sup>7-9</sup> loss of school days and increased days with restricted activity,<sup>10,11</sup> diminished ability to learn,<sup>12</sup> diminished oral health-related quality of life,<sup>13</sup> and hospitalizations and emergency room visits.<sup>14-16</sup>

Because of the aggressive nature of ECC, areas of demineralization and hypoplasia can rapidly develop cavitation. If untreated, the disease process can rapidly involve the dental pulpal tissue leading to dental infection and possibly life-

threatening fascial space involvement. Such infections may result in a medical emergency requiring hospitalization, antibiotics, and extraction of the offending tooth.<sup>17</sup>

Prevention of ECC begins during the pre-natal and perinatal periods.<sup>18</sup> Women should be advised to optimize nutrition during their pregnancy and the infant's first year, when enamel is undergoing maturation. Enamel defects are common in children with low birthweight or systemic illness or undernutrition during the perinatal period.<sup>19,20</sup>

Although enamel hypoplasia is a risk factor because the teeth are not as well formed, the etiology of ECC is bacterial; with mutans streptococci (MS) is the group of microorganisms most studied regarding the pathogenesis of ECC.<sup>21</sup> Children at high caries risk are colonized early by MS<sup>22</sup> that is transmitted most frequently from caregiver to child through salivary contact. The bacteria also can be transmitted between other members of a family or other children.<sup>23</sup> In association with the microbial etiology, high frequency sugar consumption is a caries risk factor. Caries-conducive dietary practices, including prolonged and/or frequent bottle or training cup feeding with sugar-containing drinks and frequent in between meal consumption of sugar-containing snacks or drinks (eg, juice, formula, soda), increase the risk of caries.<sup>24</sup>

Those children at risk for ECC should have care provided by a practitioner who has the training, experience, and expertise to manage both the child and the disease process. The use of anticariogenic agents, especially twice daily brushing with fluoridated toothpaste and the frequent application of fluoride varnish, may reduce the risk of development and progression of caries. A 'smear' of fluoridated toothpaste for children less than 2 years of age may decrease risk of fluorosis. A 'peasized' amount of toothpaste is appropriate for children aged 2 through 5 years.<sup>17</sup> When determining the risk-benefit of fluoride, the key issue is mild fluorosis versus preventing devastating dental disease. Interim therapeutic restorations (ITR), using materials such as glass ionomers that release fluoride, are efficacious in both preventive and therapeutic approaches.<sup>25,26</sup>

Stainless steel crowns often are indicated to restore teeth with large carious lesions and extensive white spot lesions and, at this early age, are less likely than other restorations to require retreatment.<sup>27,28</sup> Low levels of compliance with follow-up care and a high rate of children requiring additional treatment also can influence a practitioner's decisions for a more definitive restorative management of ECC.<sup>29</sup>

The extent of the disease process as well as the patient's developmental level and comprehension skills affect the practitioner's behavior guidance approaches. To perform treatment safely, effectively, and efficiently, the practitioner caring for a child with ECC often must employ advanced behavior guidance techniques. These may include protective stabilization and/or sedation or general anesthesia. The success of restorations may be influenced by the child's level of cooperation during treatment, and general anesthesia may provide better conditions to perform restorative procedures. General anesthesia, under certain circumstances, may offer a cost-saving alternative to sedation for children with ECC.<sup>30</sup>

### Policy statement

The AAPD recognizes the unique and virulent nature of ECC. Non-dental health care providers who identify ECC should either provide therapy or refer the patient to a licensed dentist for treatment and establishment of a dental home.<sup>31</sup> Immediate intervention is medically necessary to prevent further destruction, as well as more widespread health problems. Because children who experience ECC are at greater risk for subsequent caries development, preventive and therapeutic measures such as optimizing home care, ITR, more frequent professional visits with regiminated applications of topical fluoride, and full crown coverage often are necessary. The dentist must assess the patient's developmental level and comprehension skills, as well as the extent of the disease process, to determine the need for advanced behavior guidance techniques such as protective stabilization, sedation, or general anesthesia.

### References

1. Proceedings of the Conference on Early Childhood Caries, Bethesda, Md; October 1997. *Community Dent Oral Epidemiol* 1998;26(suppl):1-119.
2. Dye BA, Tan S, Smith V, et al. Trends in oral health status: United States, 1988-1994 and 1999-2004. National Center for Health Statistics. *Vital Health Stat* 11(248). Hyattsville, Md; 2007.
3. Tinanoff N, Reisine S. Update on early childhood caries since the Surgeon General's Report. *Academic Pediatr* 2009;9(6):396-403.
4. Al-Shalan TA, Erickson PR, Hardie NA. Primary incisor decay before age 4 as a risk factor for future dental caries. *Pediatr Dent* 1997;19(1):37-41.
5. Peretz B, Ram D, Azo E, Efrat Y. Preschool caries as an indicator of future caries: A longitudinal study. *Pediatr Dent* 2003;25(2):114-8.
6. Kanellis MJ, Damiano PC, Monamy ET. Medicaid costs associated with the hospitalization of young children for restorative dental treatment under general anesthesia. *J Public Health Dent* 2000;60(1):28-32.
7. Acs G, Lodolini G, Kaminshy S, Cisneros GJ. Effect of nursing caries on body weight in pediatric populations. *Pediatr Dent* 1992;14(5):302-5.
8. Ayhan H, Suskan E, Yildirim S. The effect of nursing or rampant caries on height, body weight, and head circumference. *J Clin Pediatr Dent* 1996;20(3):209-12.
9. Sheller B, Churchill SS, Williams BJ, Davidson B. Body mass index of children with severe early childhood caries. *Pediatr Dent* 2009;31(3):216-21.
10. Resine ST. Dental health and public policy: The social impact of disease. *Am J Public Health* 1985;75(1):27-30.
11. Gift HC, Reisine ST, Larach DC. The social impact of dental problems and visits. *Am J Public Health* 1992;82(12):1663-8.
12. Blumenshine SL, Vann WF, Gizlice Z, Lee JY. Children's school performance: Impact of general and oral health. *J Public Health Dent* 2008;68(2):82-7.
13. Filstrup SL, Briskie D, daFonseca M, Lawrence L, Wandera A, Inglehart MR. The effects on early childhood caries (ECC) and restorative treatment on children's oral health-related quality of life (OHRQOL). *Pediatr Dent* 2003;25(5):431-40.
14. Ladrillo TE, Hobdell MH, Caviness C. Increasing prevalence of emergency department visits for pediatric dental care 1997-2001. *J Am Dent Assoc* 2006;137(3):379-85.
15. Oliva MG, Kenny DJ, Ratnapalan S. Nontraumatic dental complaints in a pediatric emergency department. *Pediatr Emerg Care* 2008;24(11):757-60.
16. Griffin SO, Gooch BF, Beltran E, Sutherland JN, Barsley R. Dental services, costs, and factors associated with hospitalization for Medicaid-eligible children, Louisiana 1996-97. *J Public Health Dent* 2000;60(3):21-7.
17. Sheller B, Williams BJ, Lombardi SM. Diagnosis and treatment of dental caries-related emergencies in a children's hospital. *Pediatr Dent* 1997;19(8):470-5.
18. Ismail AI. Prevention of early childhood caries. *Community Dent Oral Epidemiol* 1998;26(suppl):49-61.
19. Davies GN. Early childhood caries: A synopsis. *Community Dent Oral Epidemiol* 1998;26(suppl):106-16.
20. Seow WK. Biological mechanisms of early childhood caries. *Community Dent Oral Epidemiol* 1998;26(suppl):8-27.
21. Kanasi E, Johansson J, Lu SC, et al. Microbial risk markers for childhood caries in pediatrician's offices. *J Dent Res* 2010;89(4):378-83.
22. Karn T, O'Sullivan DA, Tinanoff N. Mutans streptococci levels in 8-15 month-old children. *J Public Health Dent* 1998;58(3):248-9.
23. Berkowitz RJ. Mutans streptococci: Acquisition and transmission. *Pediatr Dent* 2006;28(2):106-9.

24. Tinanoff NT, Palmer C. Dietary determinants of dental caries in preschool children and dietary recommendations for preschool children. *J Pub Health Dent* 2000;60(3):197-206.
25. van't Hof MA, Frencken JE, van Palenstein Helderma WH, Holmgren CJ. The atraumatic restorative treatment (ART) approach for managing dental caries: A meta-analysis. *Int Dent J* 2006;56(6):345-51.
26. American Academy of Pediatric Dentistry. Guideline on Restorative Dentistry. *Pediatr Dent* 2010;32(special issue):87-93.
27. Randall RC, Vrijhoef MM, Wilson NH. Efficacy of preformed metal crowns vs amalgam restorations in primary molars: A systematic review. *J Am Dent Assoc* 2000;131(3):337-43.
28. Eidelman E, Faibis S, Peretz B. A comparison of restorations for children with early childhood caries treated under general anesthesia or conscious sedation. *Pediatr Dent* 2000;22(1):33-7.
29. Almeida AG, Roseman MM, Sheff M, Huntington N, Hughes CV. Future caries susceptibility in children with early childhood caries following treatment under general anesthesia. *Pediatr Dent* 2000;22(4):302-6.
30. Lee JY, Vann WF, Roberts MW. A cost analysis of treating pediatric dental patients using general anesthesia vs conscious sedation. *Pediatr Dent* 2000;22(1):27-32.
31. American Academy of Pediatric Dentistry. Policy on a dental home. *Pediatr Dent* 2011;33(special issue):24-5.

Copyright of Pediatric Dentistry is the property of American Society of Dentistry for Children and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.