Dental Erosion in Cerebral Palsy Patients

Ginna Kércia Matos Gonçalves, DDSc, MS Fabíola Grammatico Carmagnani, DDSc, MS Maria Salete Nahás Pires Corrêa, DDSc, PhD Danilo A. Duarte, DDSc, PhD Maria Teresa Botti Rodrigues Santos, DDSc, PhD

ABSTRACT

Purpose: The objective of this study was to evaluate the presence and severity of dental erosion in cerebral palsy (CP) patients.

Methods: A group of 48 noninstitutionalized individuals with CP, 2 to 18 years old (8.8 ± 3.9 years), has been diagnosed with the disorder. Dental erosion was evaluated according to O'Brien.¹⁶ Results were compared to a control group composed by 26 patients with no neurological damage aged 4 to 18 years (11.3 ± 3.5 years).

Results: Presence of deep dental erosion was significantly higher in CP individuals when compared to the control group. Concerning motor disorder, no statistical difference has been observed between the types of CP. Most of the affected teeth observed in the study group were the upper and lower molars and upper incisors.

Conclusions: Cerebral palsy patients have a higher risk of dental erosion development. (J Dent Child 2008;75:117-20)

Received June, 2006; Revision Accepted November 2, 2006.

Keywords: cerebral palsy, erosion, special health care needs

Dental erosion has been described as "the physical result of a pathologic, chronic localized loss of dental hard tissue that is chemically etched away from the tooth surface by acid and/or chelation without bacterial involvement."¹⁻⁴ Teeth affected by erosion become rounded and lose their surface characteristics, and any existing restorations become raised above the adjacent eroded occlusal surfaces. Once dentin is exposed, the "cupping" of "lesions" on the occlusal surface can be seen.⁵

Dental erosion can be the result of various systemic conditions, which often makes the etiology difficult to identify.⁶ Those conditions include: upper gastrointestinal disorders with an acid diet (43%), upper gastrointestinal disorders (25%), an acid diet (24%), eating disorders (6%), and unknown causes (2%). Unfortunately, the cause of dental erosion often goes undiagnosed, or the presence of other factors such as abrasion and attrition make diagnosis more difficult to determine.⁷ The underlying etiology of dental erosion is a source of acid, which may be intrinsic or extrinsic, acting on susceptible tooth. Extrinsic causes of dental erosion may arise from several sources, including occupational, medications, or through lifestyle practices.⁸

The propulsion of gastric contents into the mouth, such as in gastroesophageal reflux, is the most common source of intrinsic acids in the mouth. Since the stomach's acidity may be below pH 1, dental erosion has been observed in disorders associated with chronic vomiting, persistent regurgitation, or gastroesophageal reflux or with protracted rumination.⁹

This may be more significant in the etiology of tooth surface loss in children with CP than the parafunctional activity, which has been classically regarded as the cause. CP patients are known to have a high incidence of feeding difficulties with food loss due to excessive drooling, coughing, choking, and poor motor control; and difficulties with swallowing, vomiting, recurrent chest infections, and irritability.^{10,11}

Oral motor skills are often dysfunctional in CP patients, and lingual dysfunction, specifically tongue thrust, is consistently observed. Other patterns of oral dysfunction include: hyperactive and hypoactive gag reflexes, oral hypersensitivity, and prolonged and exaggerated bite reflexes. Inadequate function of cheek and lip musculature may prevent

Dr. Gonçalves has a master degree in Pediatric Dentistry; Dr Carmagnani is orthodontist and has a master degree in Pediatric Dentistry ; Drs. Corrêa and Santos are professors, all in the Division of Dentistry for Persons with Disabilities; and Dr. Duarte is professor, Department of Pediatric Dentistry, all at the School of Dentistry, University of Cruzeiro do Sul of São Paulo, São Paulo, Brazil.

Correspond with Dr. Gonçalves at ginnagois@ yahoo.com.br

formation of an adequate oral lip seal, leading to food/liquid loss and inhibiting distal propulsion of an organized bolus.¹²⁻¹⁵

The objective of this study was to evaluate the presence and severity of dental erosion in cerebral palsy patients.

METHODS

Forty-eight noninstitutionalized patients (29 males and 19 females), aged 2 to 18 yrs (8.8±3.9) with a medical diagnosis of CP, who were being treated at the Disabled Individuals Division of the Cruzeiro do Sul University (UNICSUL), São Paulo, Brazil, were selected at random. (The characteristics of the CP patients evaluated in this study are shown in Table 3.) The Ethics Committee of the UNICSUL

approved the study protocol and, before the data collection, a written informed consent was obtained from parents/care givers responsible for each child participating in the study.

From the patients' health chart, the authors obtained the following general information: name, birth date, gender, and clinical diagnosis regarding motor disorder and clinical patterns.

The control group consisted of 26 patients (14 males and 12 females), 4 to 18 years old (11.3 \pm 4.9) from similar socioeconomic backgrounds and with no neurological damage (Table 1). Dental erosion was evaluated according to O' Brien;¹⁶ methodology using codes listed on Table 2.

Table 1. Control Group's Characteristics								
			Dental erosion (%)					
	N (%)	Age (mean±SD)	Presence	Absence				
Male	12 (46)	10.5±4.4	2 (8)	10 (38)				
Female	14 (54)	12.3±4.8	0	1 (54)				
Total	26 (100)	11.3±4.9	26 (100)					

Table 2. D	able 2. Diagnosis criteria for Dental Erosion Index				
Depth Code	CRITERIA				
0	Normal				
1	Enamel only; loss of surface characterization				
2	Enamel and dentin; loss of enamel, exposing dentin				
3	Enamel into pulp; loss of enamel and dentin, resulting in a pulp exposure				
9	Assessment cannot be made				

 Table 3. Motor Disorder and Clinical Patterns of Involvement for

 Cerebral Palsy Group

Motor disorder	CLINICAL PATTERN	Female %	Male %	Age mean±SD	Total %
Spastic	Quadriplegia	8 (44)	13 (43)	8±4.1	21 (44)
	Diplegia	5 (28)	8 (27)	9.5±4.3	13 (27)
	Hemiplegia	2 (11)	2 (7)	6.7±3.8	4 (8)
	Double hemiplegia	2 (11)	0 (0)	11±0	2 (4)
Athetoid		1 (6)	5 (17)	10.2±1.8	6 (13)
Ataxic		0 (0)	2 (7)	11±2.8	2 (4)
Total		18 (38)	30 (63)	8.8±3.9	48 (100)

Both CP and control groups were examined at the dental clinic on a dental chair under appropriate light-reflector illumination with the teeth air dried and the help of a mirror. Intraexaminer reliability was established through the reexamination of 20 children at 2 different visits (K=0.93).

Degree of tooth wear was evaluated and Chi-square analysis was used to compare the prevalence and depth of dental erosion in the CP group compared to a control group. The prevalence of erosion was also examined within the CP group by motor disorder. Alpha levels were set at P<.05.

RESULTS

Twenty-six patients (54%) from the CP group and 2 patients (8%) from the control group showed dental erosion. Thus significantly higher dental erosion was observed in CP patients when compared to the control group (P<.0001). Concerning gender, the authors observed dental erosion in the CP group in 11 females (58%) and 15 males (52%). In the control group, the authors observed the presence of dental erosion in only two of the boys (12%) and none of the girls. The authors did not observe a significant association between males and females in the CP group (P=.451) as well as in the control group (P=.280). Concerning motor disorder, no statistical difference (P=.282) was observed between spastic and dystonic types (athetoid and ataxic).

The findings related to depth of erosion showed that 17 (65%) patients from the CP group patients presented with code 1 and 9 patients (35%) presented with code 2. Chisquare test shows a statistical difference regarding depth of erosion between code 1 and 2 (P=.027) for the CP group.

In the control group, one patient (4%) presented with erosion depth code 1 and one presented with code 2 (4%). The CP group presented with significantly higher erosion depth code 1 (P=.001) and 2 (P=.005), however, when compared to the control group (Figure 1).

Most of the affected teeth observed in the CP group were upper molars (54%), lower molars (58%), and upper incisors (54%).



Figure 1. CP Group Control Group

DISCUSSION

The incidence of dental erosion in patients who have not shown any alteration of the neuromotor system is widely studied. Although this prevalence in CP patients is not yet known, studies have shown that gastroesophageal reflux occurs frequently in CP.²⁰⁻²¹ Gastric dysfunction, which is a problem in neurologically impaired children, is considered one of the principal risk factors associated with dental erosion.¹¹ Acidic fruit juices were shown to provoke reflux, and they may have, therefore, extrinsic and intrinsic effects.²⁰

It is also acknowledged that the etiology of dental erosion is multifactorial. Extrinsic factors, such as diet, play a significant role. However, intrinsic factors like gastroesophageal reflux disease (GERD) and other medical conditions, as oralmotor disabilities, are also responsible for dental erosion.¹⁸ The inability to produce the necessary motor skills for ingestion^{10,19} leads to a longer permanence of food and beverages in the oral cavity. This makes it difficult to identify the true etiological factors through basic questionnaires, as one must consider the "silent refluxes." In this study, the result was extremely significant for dental erosion (P<.0001) in CP individuals. The maxillary teeth show a pattern of higher incidence and greater severity than the mandibular teeth.

The mandibular arch has a higher incidence of dental erosion on the premolars and molars, with minimal or no erosion of the lower anterior teeth.²¹ This could be attributed to the fact that regurgitated gastric acid predominantly contacted the lingual surfaces of the upper anterior teeth. Due to the salivary glands' proximity, the tongue and the buffering ability of saliva may have protected the lower anterior teeth.²¹ Premolars could not be evaluated because the majority of patients were in the mixed dentition, and premolars were not present.

Lack of early guidance and of prevention of dental deterioration caused by chemical agents can aggravate its prognosis. The early diagnosis and adequate management of dental erosion may prevent further dental tissue loss and consequently the diminished vertical dimension of occlusion and complex restorative treatments. Since dental erosion is irreversible, the goal should be the immediate referral of the patient to a gastroenterologist, who can more accurately confirm the diagnosis and begin treating the disorder. Dental practitioners may be able to help minimize erosion in their patients through the clinical applications of fluoridated solutions, aiming at reducing erosion progression in enamel and dentin.²²

CONCLUSIONS

Based on this study's results, the following conclusions can be made: this study demonstrates that dental erosion is common in cerebral palsy patients, and these results suggest that dentists should be alert to early signs of dental erosion in cerebral palsy patients and provide appropriate preventive therapy and referral for diagnosis and treatment of gastroesophageal reflux disease to avoid irreversible damage to the dentition.

REFERENCES

- 1. Ten Cate JM, Imfeld T. Dental erosion: Summary. Eur J Oral Sci 1996;104:241-4.
- 2. Ecles JD. Tooth surface loss from abrasion, attrition, and erosion. Dent Update 1982;9:373-81.
- 3. Watson IB, Tulloch EN. Clinical assessment of cases of tooth surface loss. Br Dent J 1985;159:144-8.
- Meurman JH, Ten Cate JM. Pathogenesis and modifying factors of dental erosion. Eur J Oral Sci 1996; 104:199-206.
- 5. Deery C, Wagner ML, Longbottom C, et al. The prevalence of dental erosion in a United States and a United Kingdom sample of adolescents. Pediatr Dent 2000;22:505-10.
- 6. Bouquet JE, Seime RJ. Nervous bulimia: Dental perspectives. Pract Periodontics Aesthet Dent 1997;9:655-63.
- 7. Cardoso AC, Canabarro S, Myers SL. Dental erosion: Diagnosis based noninvasive treatment. Pract Periodontics Aesthet Dent 2000;12:223-8.
- 8. Pontefract H, Hughes J, Kemp K, et al: The erosive effects of some mouthrinses on enamel: A study in situ. J Clin Periodontol 2001;28:319-24.
- 9. Scheutzel P. Etiology of dental erosion—intrinsic factors [review]. Eur J Oral Sci. 1996;104:178-90.
- Krick J, Van Duyn MA. The relationship between oral-motor involvement and growth: A pilot study in a pediatric population with cerebral palsy. J Am Diet Assoc 1984;84:555.
- 11. Jarvinen VK, Rytomaa II, Heinonen OP. Risk factors in dental erosion. J Dent Res 1991;70:942-7.
- Mueller HF. Handling the young cerebral palsied children at home. London: William Heinemann Medical Books, Ltd; 1974:111-30.
- 13. Helfrich-Miller KR, Rector KL, Straka JA. Dysphagia: Its treatment in the profoundly retarded patient with cerebral palsy. Arch Phys Med Rehabil 1986;67:520-5.

- 14. Griggs CA, Jones PM, Lee RE. Video fluoroscopic investigation of feeding disorders of children with multiple handicap. Dev Med Child Neurol 1989;31:303-8.
- 15. Reid JA, King PL, Kilpatrick NM. Desensitization of the gag reflex in an adult with cerebral palsy: a case report. Spec Care Dentist 2000;20:56-60.
- 16. O'Brien M. Children's Dental Heath in the United Kingdom 1993. London: Office of Population Censuses and Survey, Majesty's Stationery Office; 1994.
- 17. Koman LA, Smith BP, Shilt JS. Cerebral palsy. Lancet 2004;363:1619-31.
- Nunn JH, Gordon PH, Morris AJ, et al. Dental erosion —changing prevalence? A review of British national children' surveys. Int J Paediatr Dent 2003;13:98-105.
- 19. Gisel EG, Birnbaum R, Schwartz S. Feeding impairments in children: Diagnosis and effective intervention. Int J Orofacial Myology 1998;24:27-33.

- 20. Shaw L, Weatherill S, Smith A. Tooth wear in children: An investigation of etiological factors in children with cerebral palsy and gastroesophageal reflux. J Dent Child 1998;65:484-6.
- 21. Su JM, Tsamtsouris A, Laskou M. Gastroesophageal reflux in children with cerebral palsy and its relationship to erosion of primary and permanent teeth. J Mass Dent Soc 2003;52:20-4.
- 22. Ganss C, Klimek U, Schaffer T. Effectiveness of two fluoridation measures on erosion progression in human enamel and dentine in vitro. Caries Res 2001;35:325-30.