Association of Denture Use with Sleep-disordered Breathing Among Older Adults

Yohannes W. Endeshaw, MD, MPH; Sheri Katz, DDS; Joseph G. Ouslander, MD; Donald L. Bliwise, PhD

Abstract

Objective: This study investigates the relationship between sleep-disordered breathing and denture use. Methods: This was a cross-sectional study of community-dwelling older adults. Information about denture use was obtained using a questionnaire. Ambulatory sleep recording in subjects' homes was performed using Embletta PDS (Medcare, Iceland). Chi-square tests and logistic regression analysis were used for statistical analysis. Results: A total of 58 subjects completed the study. The mean apnea hypopnea (AHI) index was 15.1±16.1. Twenty-two subjects (38%) used dentures and most removed them before sleep. There was significant association between denture use and AH≥15 per hour of sleep (odds ratio [OR]=6.29; 95% confidence interval [CI]=1.71, 23.22; P=.006). Conclusions: This preliminary study found an association between sleep-disordered breathing and denture use, which may represent a proxy for a relationship between sleep-disordered breathing and edentulism. Given the common occurrence of both conditions among older adults, the observed relationship warrants a more detailed investigation of the mechanisms whereby loss of teeth leads to upper airway closure during sleep. [J Public Health Dent 2004;64(3):181-83]

Key Words: sleep apnea syndrome, dentures, edentulous mouth.

Edentuiism (loss of all natural permanent teeth) is a common problem affecting close to 25 percent of older adults in the United States (1). The anatomical changes of the face associated with edentulism include decrease in the vertical dimension of occlusion (2), change in the position of the mandible (3) and the hyoid bone (4), and possible impaired function of the oropharyngeal muscles (5,6). Loss of posterior teeth may also decrease the vertical dimension of occlusion (7).

Sleep-disordered breathing is also a prevalent condition among older adults with reported prevalence between 20 percent to 50 percent (8,9), based on specific definitions used. Obstructive sleep-disordered breathing is caused by partial or complete obstruction of the upper airway and reported risk factors include obesity, increased neck circumference, male sex, and anatomical abnormalities of the face (10-17). Although edentulism is reported to change the anatomy and also impair the function of the upper airway as described above, it has not been typically recognized as a risk factor for sleep-disordered breathing. This preliminary study was undertaken to examine the relationship between sleep-disordered breathing and denture use among communitydwelling older adults.

Methods

This was a cross sectional study of a convenience sample of communitydwelling older adults. Subjects were recruited, using a protocol approved by the Institutional Review Board of Emory University, from retirement living and senior housing facilities, and adult learning centers in Atlanta, GA. Individuals older than 64 years of age who were independent in their activities of daily living were included in the study. Individuals who were not capable of following instructions due to physical disability or cognitive impairment were excluded from the study. Subjects who volunteered to participate in the study gave written consent.

Medical history including denture use ("removable dental prosthesis") was obtained using a questionnaire. Questions included denture location (upper, lower, or both), type (partial or full), and removal during sleep. Body mass index (BMI) was calculated as: (weight in kilograms)/(height in meters)².

An overnight ambulatory sleep recording was performed for one night in the subjects' homes using Embletta PDS (Medcare, Iceland) sleep recording equipment to determine sleep-disordered breathing. The system records abdominal and chest movement, airflow through the nose and mouth, and pulse oximetry. Ambulatory recording systems similar to the Embletta PDS have been validated against polysomnography (18,19). The subjects were given both oral and written instructions on how to use the sleep recording equipment on the day of the sleep study. The sleep equipment was returned the next morning, and the data were downloaded and scored by the first author. The first author did not have any information about denture use by the subjects at the time the sleep studies were scored. At least five hours of technically adequate recordings were required for the record to be scored. The following conventional definitions were used to score the sleep recordings: (1) apnea

Send correspondence and reprint requests to Dr. Endeshaw, Division of Geriatrics and Gerontology, Emory University, 1841 Clifton Road NE, 5th Floor, Atlanta, GA 30329. E-mail: yendesh@emory.edu. Dr. Katz is in private practice in Decatur, GA. Drs. Ouslander and Bliwise are both affiliated with Emory University. Dr. Endeshaw received salary support from Hartford /AFAR Academic Fellowship Program, pilot research grant from the Hartford Southeast Center of Excellence in Geriatric Medicine, and K-12 Emory Mentored Clinical Research Program from Emory University. Preliminary results of this research were presented in poster form at the 2003 annual meeting of the American Geriatrics Society in Baltimore, MD. Manuscript received: 9/11/03; returned to authors for revision: 10/30/03; final version accepted for publication: 12/3/03. was defined as the cessation of airflow through the mouth and nose for at least 10 seconds; (2) hypopnea was defined as the reduction in airflow by 50 percent or more associated with a decrease in the pulse oximetry reading by at least 3 percent (20). Apnea hypopnea index (AHI) was defined as the number of apneas and/or hypopneas per hour of sleep. Total sleep time was estimated based on the clock time recorded by the subject for initiation of sleep and wake up time. The result of the sleep study was reported to the subjects, and with the subjects' permission, a copy of the report was sent to their physician.

T-tests for independent samples and chi-square tests were used to compare the demographic characteristics between those who completed and those who did not complete the study. A chi square test was used for comparing AHI categories between those who did and did not use dentures. Logistic regression analysis was performed with AHI ≥15 as the dependent variable and denture use, and known risk factors for sleep-disordered breathing—including age, sex, and body mass index—as covariates included in the model.

Results

Seventy-two subjects were enrolled and 58 subjects (81%) completed the study. There were 5 inadequate initial ambulatory sleep recordings (9%) requiring repeated evaluation.

Of the 14 subjects who did not complete the study, three moved to other locations, two were hospitalized, one had exacerbation of an illness, and eight declined further participation. The characteristics of subjects who completed and did not complete the study is shown in Table 1. There were no statistically significant differences in sex, age, marital status, educational status, and denture use between those who completed and did not complete the study. Of the 58 subjects who completed the study, 22 (38%) used dentures while 36 (62%) did not. Of the 22, 9 used both upper and lower, 10 used upper or lower, and 3 subjects did not

,	*		1 ,
Characteristics	Completed Study (N=58)	Did Not Complete Study (n=14)	P-value
Sex: female	44 (76%)	13 (93%)	.27*
Age mean (SD)	77.7 (6.7%)	78.43 (3.78%)	.75 †
Years of education			
<12	2%	7%	
12	23%	36%	.11*
13–15	28%	28%	
≥16	47%	29%	
Living facility			
Home	50%	7%	.003*
Independent	50%	93%	.41*
Denture use while sleeping	38%	50%	

 TABLE 1

 Characteristics of Subjects Who Completed and Did Not Complete Study

*Chi-square test.

†Independent sampe T-test.

TABLE 2 Denture Use by Apnea Hypopnea Index (AHI) Category						
Denture Use	AHI 0-4 (%)	AHI 514 (%)	AHI ≥15 (%)	Total (%)		
Yes	6 (27)	3 (14)	13 (59)	22 (100)		
No	8 (22)	22 (61)	6 (17)	36 (100)		
Total	14 (24)	25 (43)	19 (33)	58 (100)		

specify denture location. Seven subjects used full dentures, 13 subjects used partial dentures, and 2 subjects did not specify the type of dentures they used. Eighteen of the 22 subjects who used dentures (82%) removed them before going to bed.

The mean standard deviation (SD) AHI for the 58 subjects who completed the study was 15.1 ± 16.1 . The number of subjects with AHI ≥ 15 was significantly higher in the group who used dentures (Pearson's chi-square= 11.157; degree of freedom=1; *P*=.001). Table 2 shows denture use by AHI category.

By logistic regression analysis, denture use was significantly associated with AHI \geq 15 per hour of sleep (OR=6.29; 95% CI=1.71, 23.22; *P*=.006) even after controlling for other risk factors including age, sex, and body mass index. There was no significant association between AHI \geq 15 and age or sex. Body mass index was marginally associated with AHI \geq 15 (OR= 1.15; 95% CI=0.97, 1.37; *P*=.103).

There was no statistically significant difference in AHI between those who used partial versus full dentures (Fisher's exact test P=.32), or those who used either upper or lower versus both upper and lower dentures (Fisher's exact test P=.34). But due to the limited sample size, our study did not have enough statistical power to detect a difference in AHI between these groups. Only four subjects (18%) reported not removing their dentures before going to bed, thus preventing an evaluation of this practice.

Discussion

This study found a significant association between sleep-disordered breathing and denture use. As 82 percent of the study subjects removed their dentures before they went to sleep, we believe the association between sleep-disordered breathing and denture use may serve as a proxy for the relationship between sleep-disordered breathing and teeth loss, rather than a specific cause for sleep-disordered breathing. This is consistent with a previous note, where subjects were found to have worsening of the AHI and decrease in their antero-posterior oropharyngeal wall distance when examined without their dentures (21). Other studies have also noted associations between denture

use and sleep-disordered breathing (8,22,23). Because only four of the 22 subjects (28%) reported not removing their dentures before going to sleep, there was insufficient statistical power to detect any significant difference in this regard.

The mechanism underlying the relationship between teeth loss and sleep-disordered breathing likely involves changes in the upper airway anatomy and function. Reduction in the retropharyngeal space associated with impaired function of the genioglossus and other upper airway dilating muscles results in increase in the upper airway resistance (24), predisposing the subjects to apnea, hypopnea, or the upper airway resistance syndrome.

Of note was that there was no difference in AHI between those who used full and those who used partial dentures. Impaired oral function in subjects who lack their posterior teeth have been reported previously (7), which may suggest the importance of the posterior occlusion in maintaining the vertical dimension of occlusion. Unfortunately, because information on the types of partial dentures used by the subjects was not collected and the vertical dimension of occlusion was not measured, we must be circumspect regarding the mechanisms that may underlie the observed relationships. Future studies will need to evaluate such additional information.

The use of dentures may be associated with chronic inflammatory changes leading to papillary hyperplasia and other complications (24). To minimize these complications, subjects who use dentures are advised to remove their dentures before going to bed. As subjects who remove their dentures at night may be at increased risk of sleep-disordered breathing, clinical investigations are needed to determine if wearing dentures during sleep improves the upper airway patency and reduces sleep-disordered breathing. If indeed this is the case, the advice to remove dentures while going to bed may have to be modified (e.g., avoiding dentures for several hours during the daytime rather than at night), depending upon the individual's risk for sleep-disordered breathing and the risk of cardiovascular complications associated with this sleep

disorder (25-28). Given this apparent association between teeth loss and sleep-disordered breathing, it would also be prudent for denture wearers to be asked routinely about snoring and other related sleep problems associated with sleep-disordered breathing during clinic visits.

References

- Center for Disease Control and Prevention. Total tooth loss among persons aged greater than or equal to 65 years—selected states, 1995-1997. MMWR Morbid Mortal Wkly Rep 1999;48:206-11.
- 2. Unger J. Comparison of vertical morphologic measurements on dentulous and edentulous patients. J Prosthet Dent 1990;64:232-4.
- Lambadakis J, Karkazis H. Changes in the mandibular rest position after removal of remaining teeth and insertion of complete dentures. J Prosthet Dent 1992; 68:74-7.
- Tallgren A, Lang BR, Walker GF, Ash MM Jr. Changes in jaw relations, hyoid position and head posture in complete denture wearers. J Prosthet Dent 1983;50: 148-56.
- Otsuka R, Ono T, Ishiwatz Y, Kuroda T. Respiratory related genioglossus electromyographic activity in response to head position and changes in body position. Angle Orthod 2000;70:63-9.
- Lome A, Johnston W. Tongue and jaw muscle activity in response to mandibular rotations in a sample of normal and anterior open-bite subjects. Am J Orthod 1979;76:565-76.
- Tallgreen A, Mizutani H, Tryde G. A two-year kinesiographic study of mandibular movement patters in denture wearers. J Prosthet Dent 1989;62:594-600.
- Ancoli-Israel S, Kripke DF, Klauber MR, Mason WJ, Fell R, Kaplan O. Sleep disordered breathing among communitydwelling elderly. Sleep 1991;14:486-95.
- Young T, Shahar E, Nieto F, Redline S, Newman A, Gottlieb D, et al. Predictors of sleep disordered breathing in community-dwelling adults. The Sleep Heart Study. Arch Intern Med 2002;162:893-900.
- Bliwise D, Feldman D, Bliwise N, Carskadon M, Kraemer H, North C, et al. Risk factors for sleep disordered breathing in heterogeneous geriatric populations. J Am Geriat Soc 1987;35:132-41.
- Tishler P, Larkin E, Schluchter M, Redline S. Incidence of sleep-disordered breathing in an urban adult population: the relative importance of risk factors in the development of sleep-disordered breathing. JAMA 2003;289:223-37.
- Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleepdisordered breathing among middleaged adults. N Engl J Med 1993;328:1230-5
- Redline S, Kump K, Tishler PV, Browner I, Ferrette V. Gender differences in sleep disordered breathing in a community-

based sample. Am J Respir Crit Care Med 1994;149:722-6.

- 14. Liistro G, Rombaux P, Belge C, Dury M, Aubert G, Rodenstein DO. High Mallampati score and nasal obstruction are associated risk factors for obstructive sleep apnoea. Eur Respir J 2003;21:248-52.
- Jamieson A, Guilleminault C, Partinen M, Quera-Salva MA. Obstructive sleep apneic patients have craniomandibular abnormalities. Sleep 1986:469-77.
- Bixler EO, Vgontzas AN, Ten Have T, Tyson K, Kales A. Effects of age on sleep apnea in men. Am J Respir Crit Care Med 1998;157:144-8.
- Paoli J, Lauwers F, Lacassagne L, Tiberge M, Dodart L, Boutault F. Craniofacial differences according to the body mass index of patients with obstructive sleep apnoea syndrome: cephalometric study in 85 patients. Br J Oral Maxillofac Surg 2001;39:40-5.
- Emsellem HA, Corson WA, Rappaport BA, Hackett S, Smith LG, Hausfeld JN. Verification of sleep apnea using a portable sleep apnea screening device. South Med J 1990;83:748-52.
- Redline S, Tosteson T, Boucher MA, Millman RP. Measurement of sleep-related breathing disturbances in epidemiologic studies. Assessment of the validity and reproducibility of a portable monitoring device. Chest 1991;100:1281-6.
- Report of an American Academy of Sleep Medicine Task Force. Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. Sleep 1999;22:667-89.
- Bucca C, Carossa S, Pivetti S, Gai V, Rolla G, Preti G. Edentulism and worsening of obstructive sleep apnea. Lancet 1999; 353:121-2.
- Knudson RC, Meyer JB. Fabrication of prosthesis to prevent sleep apnea in edentulous patients. J Prosthet Dent 1990;63:448-51.
- Reeves-Hoche MK. Absence of mandibular molars and other dental health issues as risk factors associated with obstructive sleep apnea: a case control study. Sleep Res 1996;25:342.
- Budiz-Jorgensen E. The edentulous patient. In: Owall B, Kayser A, Carlson G, eds. Prosthodontics: principles and management strategies. London: Mosby-Wolfe, 1996:65-79.
- Nieto J, Young T, Lind B, Shahar E, Samet J, Redline S, et al. Association of sleepdisordered breathing as a risk factor for hypertension. JAMA 2000;283:1829-36.
- Peppard PR, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. N Engl J Med 2000;342: 1378-84.
- 27. Peker Y, Hedner J, Kraiczi H, Loth S. Respiratory Disturbance Index. An independent predictor of mortality in coronary heart disease. Am J Resp Crit Care Med 2000;162:81-6.
- Mohsenin V. Sleep-related breathing disorders and the risk of stroke. Stroke 2001; 32:1271-8.