Racial and Ethnic Variation in the Provision of Dental Procedures

Christopher Okunseri, BDS, MSc, DDPHRCSE, FFDRCSI; Ruta Bajorunaite, PhD; Richards Matthew, BS; Anthony M. Iacopino, DMD, PhD

Abstract

Objectives: Most of the available information on racial/ethnic disparities in oral health is based on differences in sociodemographic variables related to dental disease burden, dental visits, and access to care. However, very little is known regarding racial/ethnic variation in the provision of dental procedures. This study examined trends in the provision of dental procedures and sought to determine whether there are racial/ethnic differences in the provision of dental procedures. Methods: This is a retrospective observational study of patients treated at a dental training institution. Data for all patients 18 to 60 years of age in axiUm (electronic database) for 2001 to 2003 were analyzed. Data include demographic information, poverty status, insurance coverage, dental procedure, and race/ethnicity. Separate logistic regression models (by dental procedure category and year) were fitted while considering race/ethnicity, insurance coverage, poverty status, marital status, and age as possible covariates. Results: The total number of dental procedures completed by providers increased by 14,000 between 2001 and 2003. African-Americans were significantly less likely to have restorative procedures [odds ratio (OR): 0.60, 95 percent confidence interval (CI): 0.42 to 0.86], (OR: 0.52, 95 percent CI: 0.38 to 0.73), (OR: 0.46, 95 percent CI: 0.36 to 0.58) in 2001, 2002, and 2003, respectively, than the White population. Significant differences in the use of other dental procedures (prosthodontics - removable) and oral surgery procedures by race/ethnicity were observed. Conclusions: Substantial racial/ethnic variation in the provision of dental procedures exists. This study presents findings beyond anecdotal information on racial/ethnic variation in the provision of dental procedures and requires further research to compile more detailed data.

Key Words: dental procedures, race, adults, oral health disparities, health services

Introduction

Racial and ethnic variations in the provision of appropriate dental and medical procedures are important component measures of health and oral health disparities. Medical and dental care services provided to patients based on their race/ethnic origin could lead to poor oral health outcomes. The Surgeon General's report Oral Health in America documents the burden and severity of dental disease and the existence of oral health disparities in the United States (1). The Institute of Medicine report *Unequal Treatment Confronting Racial and Ethnic Disparities in Health Care* identified significant variations in the use of medical procedures by race/ethnicity (2). Factors related to patient, provider, patient–doctor relations, and the health care system were identified as possible causes of health disparities in the use of health procedures (2).

Peterson et al. showed that Blacks with acute myocardial infarction received substantially fewer cardiac catheterization, coronary angioplasty, coronary bypass surgery, and subsequent cardiac revascularization procedures than Whites (3). Ellison et al. examined racial variation in the receipt of bowel surveillance after receiving curative colorectal cancer surgery treatment and found that elderly Blacks were less likely than non-Hispanic Whites to receive posttreatment bowel surveillance after adjusting for sociodemographic, hospital, and clinical characteristics (4). Disparities in health outcomes are associated with health disparities use and process, with racial/ethnic minority groups reporting lower health care satisfaction and greater discrimination (5, 6).

Results from a multivariate analysis show that racial/ethnic variation in the use of dental service exists, with African-Americans being much less likely to receive dental prophylaxis, restorative treatment procedures, and fixed prosthodontic procedures (7). In a study that examined the influence of patient race/ethnicity on dentist decisions to extract or retain a decayed tooth, results show that racial variation

Send correspondence and reprint requests to Christopher Okunseri BDS, MSc, DDPHRCSE, FFDRSCI, Marquette University School of Dentistry, Department of Clinical Services, Room 356, P.O. Box 1881, Milwaukee, Wisconsin. 53201-1881. Tel.: 414-288-6524; Fax: 414-288-3586; e-mail: christopher.okunseri@marquette.edu. Christopher Okunseri and Richards Matthew are with the Department of Clinical Services, Marquette University School of Dentistry, Milwaukee, Wisconsin. Ruta Bajorunaite is with the Department of Mathematics, Statistics and Computer Science, Marquette University. Anthony M. Iacopino is with the Department of General Dental Sciences, Marquette University School of Dentistry. A poster presentation presented in part as an abstract at the 7th Joint Meeting of the Association of State and Territorial Dental Directors and the American Association of Public Health Dentistry in Little Rock, Arkansas, May 2006. Manuscript received: 7/13/06; accepted for publication: 10/20/06.

occurred irrespective of dentist sociodemographic characteristic or whether they have had postgraduate training (8). However, in terms of practice setting, racial variation was identified in company and private practice settings but not in public and military practice settings (8). Another study, using a veteran population, showed substantial racial/ ethnic variation in dentist decisions to prescribe root canal treatment versus tooth extraction (9).

Public perception is that dental teaching institutions (just like medical teaching institutions) are probably excellent places to get appropriate health care at a fairly reasonable cost. This study investigates trends in the provision of dental procedures and whether race/ethnicity is associated with the provision of dental procedures in a dental teaching facility. To the best of our knowledge, no study has examined the provision of a battery of dental treatment procedures together in a dental teaching institution.

Methods

Α retrospective observational study of patients treated at a dental training institution (Marquette University School of Dentistry, Milwaukee, WI) was conducted. Data for all patients 18 to 60 years of age in the axiUm (electronic database) for 2001 to 2003 were analyzed. Data included demographic information, poverty status, insurance coverage, marital status, code of the dental procedure performed, and race/ ethnicity (African-American, White, Hispanic, and others). Race/ethnicity is entered into axiUm based on patient's self-reported preference. The "unknown" category in the insurance column represents people who mostly pay directly out of pocket for their dental care service. The database consists of a small number of repeat patients as well as those who only attended the dental clinic once in a given year.

Dental treatment procedure codes were grouped according to the American Dental Association categories of preventive, diagnostic, restorative, endodontic, periodontics, prosthodontic (removable), surgical, prosthodontic (fixed), orthodontic, and others. Descriptive statistics were used to describe sample characteristics, with Chisquared tests to determine significant differences between groups. Separate logistic regression was used to model the probability of certain dental procedures being performed while considering race/ethnicity, insurance coverage, poverty status, marital status, and age as possible covariates. Race/ethnicity represents the main focus, and it was therefore included in each final model. A backward elimination model selection procedure was employed to identify statistical significant covariates to be added into the model. A statistical significance (alpha) level of 0.05 was used throughout. SAS version 9.1 (SAS Institute, Inc., Cary, NC) was used to perform all statistical analysis. The study was approved by the Marquette University Institutional Review Board.

Results

Table 1 shows the frequencies of study population characteristics. A total of 5,111 patients visited the dental institution for treatment, and 29,975 dental treatment procedures were completed for patients 18 to 60 years of age during the 3-year period. Significant differences (P <0.05) exist in all three years of study in sex, age, poverty status, race/ ethnicity, marital status, and insurance coverage. More than 50 percent of all patients were females, and over 40 percent of those treated at the clinic were below poverty level. In each year, the minimum number of African-Americans was 19 percent, and at least 42 percent of the patients had public insurance.

Table 2 reveals that in 2001, 2002, and 2003 a significant difference (P < 0.0001) exists in poverty status, insurance coverage, and age by race/ethnicity. A higher percent of White population (25 to 32 percent) were above poverty status compared with African-Americans (11 to 17 percent) and Hispanics (9 to 12 percent). In terms of insurance coverage, a higher percent of African-Americans (range: 60 to 63 percent) had public insurance compared with Hispanics (range: 42 to 54 percent) and Whites (range: 30 to 32 percent).

Table 3 shows the bivariate analysis between race/ethnicity and dental treatment procedures for each year. Significant differences (P <0.05) exist in the use of diagnostic/ preventive treatment procedures in 2003, but not in 2002 and 2001. Significant differences (P < 0.05) exist in the use of restorative and surgery treatment procedures in all three years and endodontic treatment procedures for 2002 and 2003. In 2001 and 2002, the differences in the use of periodontics and prosthodontic (removable) procedures were significant (P < 0.05). In 2003, all dental treatment procedure categories were found to be significantly different (P < 0.05).

Table 4 shows the final logistic regression models. African-Americans were significantly (P <0.05) less likely to have: restorative procedure (OR: 0.60, 95 percent CI: 0.42 to 0.86), (OR: 0.52, 95 percent CI: 0.38 to 0.73), (OR: 0.46 95 percent CI: 0.36 to 0.58) and twice more likely to have oral surgery procedures (OR: 1.57, 95 percent CI: 1.07 to 2.32), (OR: 2.19, 95 percent CI: 1.56 to 3.06), (OR: 2.52, 95 percent CI: 2.03 to 3.12) in 2001, 2002, and 2003, respectively. Diagnostic-preventive treatment procedures were more likely to be received by those who pay out of pocket in all three years. But in 2002 and 2003, only patients 48 years plus and those who identified themselves as single in 2003 were more likely to have diagnostic-preventive treatment procedures.

Odds ratio for restorative and surgery treatment procedures for the Hispanic group were very similar to those seen in the African-American population. African-Americans were less likely to have diagnosticpreventive procedures in 2001 and 2002 and more likely to have them in 2003. Insurance coverage and age were significant factors (P < 0.05)

| Frequencies o | Frequencies of Study Population Characteristics | | | | | | | |
|--|---|-------|--------|--------|--|--|--|--|
| | 2001 | 2002 | 2003 | Р | | | | |
| Total number of patients | 882 | 1.281 | 2.948 | | | | | |
| Total number of dental procedures done | 4,903 | 6,953 | 18,119 | | | | | |
| * | | % | | | | | | |
| Sex | | | | 0.03 | | | | |
| Female | 54 | 57 | 56 | | | | | |
| Male | 45 | 42 | 42 | | | | | |
| Unknown | 0.5 | 1 | 2 | | | | | |
| Age (years) | | | | < 0.01 | | | | |
| 18 to 33 | 22 | 27 | 38 | | | | | |
| 33 to 48 | 37 | 37 | 36 | | | | | |
| 48 and older | 41 | 36 | 26 | | | | | |
| Poverty status | | | | < 0.01 | | | | |
| Above | 24 | 20 | 14 | | | | | |
| Below | 49 | 49 | 41 | | | | | |
| Unknown | 27 | 31 | 45 | | | | | |
| Race/Ethnicity | | | | < 0.01 | | | | |
| White | 57 | 50 | 42 | | | | | |
| African-American | 23 | 19 | 20 | | | | | |
| Hispanic | 5 | 6 | 5 | | | | | |
| Others | 15 | 25 | 33 | | | | | |
| Marital status | | | | < 0.01 | | | | |
| Single | 50 | 49 | 49 | | | | | |
| Married | 28 | 25 | 23 | | | | | |
| Divorced, widowed | 16 | 14 | 11 | | | | | |
| Unknown | 6 | 12 | 17 | | | | | |
| Insurance | | | | 0.03 | | | | |
| Public | 42 | 45 | 44 | | | | | |
| Private | 9 | 8 | 7 | | | | | |
| Out-of-pocket | 30 | 33 | 32 | | | | | |
| Unknown | 19 | 14 | 17 | | | | | |

Table 1requencies of Study Population Characteristics

for use of restorative procedures in all three years and poverty status was significant in only 2002 and 2003.

African-Americans were least likely to receive endodontic treatment procedures in 2002 (OR: 0.46, 95 percent CI: 0.26 to 0.83) and in 2003 (OR: 0.43, 95 percent C.I: 0.29 to 0.66), and similar trends were seen for Hispanics. Patients with private insurance (OR: 1.71, 95 percent CI: 1.26 to 2.31), those paying out of pocket (OR: 1.71, 95 percent CI: 1.26 to 2.31), and females (OR: 1.46, 95 percent CI: 1.11 to 1.91) were significantly more likely to have endodontic treatment procedures in 2003.

Discussion

While racial/ethnic variations in the use of procedures have been

documented in medicine, very little has been reported in the literature about racial/ethnic variation in the provision of dental procedures, even with the recent increased attention paid to racial/ethnic disparities in oral health care. Studies have confirmed the existence of racial and ethnic disparities by socioeconomic status, dental visits, and quality of health care. However, this study presents an opportunity to examine racial/ethnic variation in the provision of dental treatment procedures from a dental training institution.

Kressin et al. found substantial racial variation in dentist treatment use of root canal therapy and tooth extraction after adjusting for severity of tooth and gum related disease, age, sex, medical/psychiatric comorbidities, and prior use of preventive dental services in patients treated at the Department of Veterans Affairs Dental Clinic (9). Manski and Moeller identified a significant difference in the use of dental procedures, with Blacks and Hispanics being more likely to use oral surgical procedures than Whites (10). Results in our study were similar, with African-Americans being twice more likely to have oral surgery treatment procedures as compared with the White population.

African-Americans were also less likely to have diagnostic/preventive treatment procedures, restorative treatment procedures, and periodontics treatment procedures even after adjusting for insurance coverage. While patient insurance plan/ coverage is important in the final determination of the type of dental procedure or treatment a provider performs on a patient, our study shows that patients with private insurance, those paying out-ofpocket, and females were twice more likely to have endodontic treatment procedures. African-Americans were significantly (P < 0.05) less likely to have restorative care than the White population.

Although African-Americans and Hispanics have poorer oral health and are less likely to visit the dentist within a 12-month period compared with Whites, racial differences in the characteristics of practices attended independently contributed to patientspecific tooth loss and to the patientlevel racial disparity, even with detailed tooth-specific and patient factor-level characteristics taken into account (11). Cabral et al. reported that there was no racial variation identified in a public practice setting (7), but our study shows racial variation in the provision of dental procedures. Racial variation could be a result of patient cultural beliefs, but our research did not address this question.

Despite the teaching in dental schools that treatment planning should be based on good history, clinical assessment, investigations, correct diagnosis, and patient consent, dentist bias is suggested as a possible influence on the treatment decision when prescribing extraction

Racial and Ethnic Variation in Dental Procedures

| | White (%) | African- American (%) | Hispanic (%) | Other (%) | Unknown (%) | Р |
|---|-----------|--------------------------|--------------|------------|-------------|-------|
| Age (vears) | | | 2001 | | | <0.01 |
| 18 to 33 | 17 | 23 | 38 | 36 | 28 | <0.01 |
| 33 to 48 | 38 | 23 | 38 | 38 | 20 48 | |
| /8 and older | 50 /15 | 20 49 | 30 24 | 26 | 24 | |
| Poverty status | 1) | 1) | 21 | 20 | 21 | <0.01 |
| Above | 32 | 17 | 0 | 33 | 0 | <0.01 |
| Below | 42 | 68 | 76 | 41 | 28 | |
| Unknown | 26 | 15 | 16 | 26 | 20 72 | |
| Insurance | 20 | 1) | 10 | 20 | / 4 | |
| Public | 30 | 60 | 42 | 33 | 60 | <0.01 |
| Private | 11 | 9 | 7 | 10 | 5 | <0.01 |
| Out-of-pocket | 37 | 23 | 22 | 21 | 7 | |
| Unknown | 23 | 23 | 18 | 26 | 10 | |
| Ulikilowii | 25 | 2 | 2002 | 20 | 19 | |
| Age (vears) | | | 2002 | | | <0.01 |
| 18 to 33 | 20 | 27 | 40 | 35 | <i>/</i> 1 | <0.01 |
| $\frac{10 \text{ to } 55}{23 \text{ to } 48}$ | 20 | 27 | 17 | 35 | 40 | |
| 19 10 40 18 and older | J/ /12 | 33 40 | 17 | 30 | 40 | |
| Poverty status | 43 | 40 | 1/ | 30 | 19 | ~0.01 |
| Abovo | 20 | 15 | 11 | 24 | 0 | <0.01 |
| Below | 29 48 | 13 | 11 70 | 24 40 | 25 | |
| Linknown | 40 | /1 | 11 | 49 | 2) | |
| Unknown | 23 | 14 | 11 | <u>Z</u> / | /3 | |
| Dublic | 21 | 62 | 5 / | 22 | 62 | <0.01 |
| Public | 51 | 05 | 34 | 55 10 | 02 | <0.01 |
| Private Out of position | 11 | 4 | 4 | 10 |) 10 | |
| University | 45 | 10 | 55 | 30 10 | 19 | |
| UIIKIIOWII | 15 | 10 | 2002 | 19 | 15 | |
| Ago (Noors) | | | 2005 | | | <0.01 |
| 19 to 22 | 20 | 41 | 47 | 20 | 47 | <0.01 |
| 10 10 55 | 50 25 | 41 | 4/ | 30 20 | 4/ | |
| 55 10 40 48 and older | 25 | 54 25 | 59 14 | 39 22 | 50 17 | |
| 48 and older | 32 | 25 | 14 | 25 | 1/ | <0.01 |
| Above | 25 | 11 | 10 | 25 | 0 | <0.01 |
| ADOVE D-1 | 25 45 | | 12 | 25 (2 | 0 | |
| Below | 45 | 07 | 07 | 45 | 11 | |
| UIIKIIOWN | 20 | | 21 | 23 | 07 | |
| Insurance | 20 | (2 | 40 | 24 | 47 | -0.01 |
| PUDIIC | 52 | 03 | 49 | 54 | 4/ | <0.01 |
| Private | 11 | 5 | / | 8 | 4 | |
| Out-of-pocket | 42 | 22 | 36 | 45 | 22 | |
| Unknown | 15 | 10 | 8 | 14 | 2/ | |

 Table 2

 Patients' Characteristics by Race/Ethnicity for Each Year

versus a tooth-sparing procedure to a patient (12). Other factors include dentist beliefs about the extent/ prognosis of disease and oral hygiene as well as their school of graduation (8). While this study did not investigate dental student bias or beliefs in their provision of dental treatment procedures, some of our results are suggestive of these elements and the need for further research in this area.

Results from this study support previous findings from medicine and

dentistry on racial/ethnic variation in the use of medical and dental procedures. However, interpretation of our study results should be done with some degree of caution because of the possibility of coding errors in the electronic database used for the study. Another limitation of the study is the nonavailability of diagnostic codes to match dental treatment procedure codes for the patients and the fact that dental students in training were used as providers. Additionally, African-Americans and Hispanics have been shown to be episodic users of the dental care system. Thus, these patients may have presented with more severe disease, ruling out many restorative procedures.

Findings from this study are not generalizable to the population at large; nevertheless, we believe that the study findings may be generalized to other similar dental teaching institutions because, like many other dental schools located in the United States, our study has a racially/ ethnically diverse population and

| Treatment procedure | White (%) | African- American (%) | Hispanic (%) | Other (%) | Unknown (%) | Р |
|-----------------------|-----------|--------------------------|--------------|-----------|-------------|---------|
| | | | 2001 | | | |
| Diagnostic-preventive | 87 | 86 | 80 | 85 | 82 | 0.76 |
| Restorative | 46 | 33 | 36 | 23 | 34 | <0.70 |
| Endodontics | 10 | 33 7 | 9 | 15 | 7 | 0.38 |
| Periodontics | 15 | 0 | 4 | 5 | 7 | 0.00 |
| Prosthodontics | 5 | 10 | 2 | 0 | 3 | 0.01 |
| Implants | 3 | 0 | 0 | 3 | 1 | 0.01 |
| Fixed prosthodontics | 4 | 3 | 4 | 3 | 0 | 0.00 |
| Surgery | 20 | 34 | 22 | 23 | 31 | <0.01 |
| Orthodoptics | 1 | 0 | 0 | 25 | 2 | 0.15* |
| Other | 12 | 10 | 9 | 8 | 12 | 0.19 |
| ouler | 12 | 10 | 2002 | 0 | 12 | 0.00 |
| Diagnostic-preventive | 85 | 82 | 84 | 86 | 83 | 0.76 |
| Restorative | 46 | 33 | 30 | 46 | 27 | <0.01 |
| Endodontics | 10 | 6 | 7 | 13 | 4 | <0.01 |
| Periodontics | 12 | 13 | 12 | 24 | 4 | <0.01 |
| Prosthodontics | 3 | 10 | 3 | 2 | 2 | <0.01* |
| Implants | 2 | 0.5 | 0 | 5 | 0 | 0.01* |
| Fixed prosthodontics | - | 2 | 3 | 3 | 0 | 0.12* |
| Surgery | 19 | 38 | 32 | 27 | 34 | <0.12 |
| Orthodontics | 0.5 | 0 | 0 | 0 | 2 | 0.09* |
| Other | 16 | 20 | 21 | 11 | 20 | 0.23 |
| o unor | 10 | -0 | 2003 | | -0 | 0.20 |
| Diagnostic-preventive | 86 | 91 | 88 | 80 | 86 | < 0.01 |
| Restorative | 38 | 21 | 27 | 39 | 24 | <0.01 |
| Endodontics | 12 | 5 | 10 | 15 | 7 | < 0.01 |
| Periodontics | 17 | 9 | 12 | 22 | 6 | < 0.01 |
| Prosthodontics | 4 | 6 | 2 | 3 | 2 | < 0.01* |
| Implants | 3 | 0 | 1 | 3 | 0 | < 0.01* |
| Fixed prosthodontics | 3 | 1 | 2 | 3 | 1 | 0.02* |
| Surgery | 26 | 50 | 44 | 20 | 37 | < 0.01 |
| Orthodontics | 1 | 0 | 1 | 2 | 3 | < 0.01* |
| Other | 18 | 17 | 17 | 16 | 17 | 0.90 |

 Table 3

 Bivariate Analyses Between Race/Ethnicity and Dental Treatment Procedures

* Because of low counts in the contingency table, Fisher's exact test (two-sided) was used to test for association.

| Table 4 | |
|---|-----------|
| Multiple Regression Analysis with Dependent and Independent | Variables |

| | 2001 | | 2002 | | 2003 | |
|----------------------------|---------------------|-------------|-------------------------|--------|---------------------|--------|
| | Odds ratio (95% CI) | Р | Odds ratio (95% CI) | Р | Odds ratio (95% CI) | Р |
| | Diagnosis-pre | eventive tr | eatment procedures cate | gory | | |
| Race/Ethnicity: Reference | : White | 0.62 | 1 | 0.40 | | 0.01 |
| Hispanic | 1.13 (0.43, 2.99) | | 0.95 (0.48, 1.87) | | 1.32 (0.75, 2.31) | |
| African-American | 0.81 (0.51, 1.32) | | 0.68 (0.45, 1.04) | | 1.60 (1.14, 2.24) | |
| Other | 0.86 (0.34, 2.15) | | 1.28 (0.59, 2.76) | | 0.68 (0.41, 1.13) | |
| Unknown | 0.63 (0.33, 1.72) | | 0.93 (0.61, 1.43) | | 1.25 (0.93, 1.67) | |
| Type of insurance: Refere | ence: Public | < 0.01 | | < 0.01 | | < 0.01 |
| Out-of-pocket | 1.02 (0.60, 1.74) | | 1.02 (0.68, 1.54) | | 1.48 (1.09, 2.00) | |
| Private | 0.41 (0.22, 0.75) | | 0.36 (0.20, 0.62) | | 0.85 (0.55, 1.32) | |
| Unknown | 0.45 (0.27, 0.75) | | 0.23 (0.15, 0.35) | | 0.34 (0.25, 0.44) | |
| Age: Reference: 18 to 33 | | NIM | | < 0.01 | | < 0.01 |
| 33 to 48 | | | 1.47 (1.02, 2.13) | | 2.25 (1.72, 2.95) | |
| 48 and older | | | 2.41 (0.6, 3.62) | | 2.33 (1.71, 3.18) | |
| Marital status: Reference: | Married | NIM | | NIM | | 0.01 |
| Single | | | | | 1.57 (1.18, 2.08) | |
| Divorced, widowed | | | | | 0.95 (0.63, 1.45) | |
| Unknown | | | | | 1.53 (1.05, 2.23) | |

| | 2001 | | 2002 | | 2003 | |
|------------------------------|---------------------|--------------|--------------------------|--------|---------------------|--------|
| | Odds ratio (95% CI) | Р | Odds ratio (95% CI) | Р | Odds ratio (95% CI) | Р |
| | Restorat | ive treatm | ent procedures category | | | |
| Race/Ethnicity: Reference: | White | 0.02 | | < 0.01 | | < 0.01 |
| African-American | 0.60 (0.42, 0.86) | | 0.52 (0.38, 0.73) | | 0.46 (0.36, 0.58) | |
| Hispanic | 0.76 (0.39, 1.46) | | 0.51 (0.30, 0.87) | | 0.63 (0.42, 0.94) | |
| Other | 0.41 (0.19, 0.90) | | 1.16 (0.68, 1.99) | | 1.11 (0.74, 1.66) | |
| Unknown | 0.78 (0.47, 1.28) | | 0.65 (0.45, 0.95) | | 0.88 (0.69, 1.12) | |
| Insurance: Reference: Publi | c | < 0.01 | | < 0.01 | | < 0.01 |
| Private | 0.94 (0.57, 1.56) | | 1.73 (0.46, 1.18) | | 1.13 (0.83, 1.55) | |
| Out-of-pocket | 1.70 (1.21, 2.40) | | 1.08 (0.82, 1.43) | | 1.18 (0.97, 1.43) | |
| Unknown | 0.59 (0.39, 0.90) | | 0.42 (0.28, 0.62) | | 0.43 (0.33, 0.57) | |
| Age: Reference: 18 to 33 | | < 0.01 | | < 0.01 | | < 0.01 |
| 33 to 48 | 2.10 (1.40, 3.16) | | 1.49 (1.10, 2.03) | | 1.80 (1.47, 2.19) | |
| 48 and older | 2.34 (1.57, 3.50) | | 1.61 (1.32, 2.48) | | 2.18 (1.76, 2.70) | |
| Poverty status: Reference: I | Below | NIM | | < 0.01 | | < 0.01 |
| Above | | | 0.93 (0.67, 1.28) | | 0.97 (0.75-1.24) | |
| Unknown | | | 0.54 (0.40, 0.74) | | 0.62 (0.50, 0.76) | |
| | Endodon | itics treatm | nent procedures category | | | |
| Race/Ethnicity: Reference: | White | 0.40 | | < 0.01 | | < 0.01 |
| African-American | 0.66 (0.36, 1.23) | | 0.46 (0.26, 0.83) | | 0.43 (0.29, 0.66) | |
| Hispanic | 0.89 (0.31, 2.58) | | 0.54 (0.21, 1.39) | | 0.81 (0.45, 1.45) | |
| Other | 1.65 (0.66, 4.13) | | 1.12 (0.51, 2.45) | | 1.20 (0.69, 2.08) | |
| Unknown | 0.66 (0.28, 1.60) | | 0.32 (0.16-0.64) | | 0.80 (0.57, 1.12) | |
| Insurance: Reference: Publi | ic | NIM | | NIM | | < 0.01 |
| Private | | | | | 1.78 (1.14, 2.78) | |
| Out-of-pocket | | | | | 1.71 (1.26, 2.31) | |
| Unknown | | | | | 0.58 (0.35, 0.95) | |
| Sex: Reference: Male | | NIM | | NIM | | < 0.01 |
| Female | | | | | 1.46 (1.11, 1.91) | |
| Marital status: Reference: M | Iarried | NIM | | | | < 0.01 |
| Single | | | | | 0.70 (0.52, 0.95) | |
| Divorced, widowed | | | | | 1.05 (0.69-1.58) | |
| Unknown | | | | | 0.48 (0.2980) | |
| | Periodon | itics treatm | nent procedures category | | | |
| Race/Ethnicity: Reference: ' | White | 0.25 | | < 0.01 | | < 0.01 |
| African-American | 0.68 (0.39, 1.19) | | 1.44 (0.90, 2.30) | | 0.65 (0.47, 0.91) | |
| Hispanic | 0.34 (0.08, 1.49) | | 1.78 (0.82, 3.90) | | 0.95 (0.55, 1.66) | |
| Other | 0.38 (0.09, 1.67) | | 3.17 (1.62, 6.20) | | 1.68 (1.02, 2.76) | |
| Unknown | 0.62 (0.25, 1.55) | | 0.49 (0.24, 0.99) | | 0.76 (0.52, 1.11) | |
| Sex: Reference: Male | | 0.03 | | 0.01 | | NIM |
| Female | 0.63 (0.41, 0.96) | | 0.62 (0.43, 0.89) | | | |
| Insurance: Reference: Public | С | < 0.01 | | < 0.01 | | < 0.01 |
| Private | 3.24 (1.66, 6.30) | | 2.43 (1.34, 4.40) | | 2.68 (1.82, 3.95) | |
| Out-of-pocket | 2.52 (1.49, 4.25) | | 1.42 (0.93, 2.18) | | 1.93 (1.47, 2.55) | |
| Unknown | 0.62 (0.28, 1.39) | | 0.71 (0.37, 1.36) | | 0.47 (0.29, 0.78) | |
| Age: Reference: 18 to 33 | | < 0.01 | | < 0.01 | | < 0.01 |
| 33 to 48 | 3.47 (1.51, 8.01) | | 4.79 (2.37, 9.67) | | 3.15 (2.23, 4.44) | |
| 48 and older | 3.81 (1.67, 8.65) | | 6.81 (3.40, 13.63) | | 5.11 (3.61, 7.21) | |
| Poverty: Reference: Below | | NIM | | NIM | | < 0.01 |
| Above | | | | | 1.02 (0.74, 1.39) | |
| Unknown | | | | | 0.58 (0.42, 0.79) | |

Table 4 Continued

| | | | minucu | | | |
|----------------------------|---------------------|------------|------------------------|--------|---------------------|--------|
| | 2001 | | 2002 | | 2003 | |
| | Odds ratio (95% CI) | Р | Odds ratio (95% CI) | Р | Odds ratio (95% CI) | Р |
| | Surger | v treatmer | nt procedures category | | | |
| Race/Ethnicity: Reference: | White | 0.05 | 1 0 1 | < 0.01 | | < 0.01 |
| African-American | 1.57 (1.07, 2.32) | | 2.19 (1.56, 3.06) | | 2.52 (2.03, 3.12) | |
| Hispanic | 0.89 (0.42, 1.89) | | 1.72 (1.01, 2.92) | | 1.88 (1.30, 2.71) | |
| Other | 1.24 (0.56, 2.74) | | 1.61 (0.88, 2.92) | | 0.64 (0.40, 1.03) | |
| Unknown | 1.94 (1.11, 3.40) | | 1.95 (1.38, 2.74) | | 1.72 (1.40, 2.10) | |
| Insurance: Reference: Pub | lic | 0.02 | | < 0.01 | | < 0.01 |
| Private | 1.30 (0.73, 2.32) | | 0.54 (0.31, .94) | | 0.47 (0.33, 0.67) | |
| Out-of-pocket | 0.76 (0.51, 1.13) | | 0.69 (0.51, .93) | | 0.95 (0.79, 1.14) | |
| Unknown | 0.54 (0.33, 0.88) | | 0.30 (0.19, .49) | | 0.26 (0.20, 0.34) | |
| Poverty status: Reference: | Below | < 0.01 | | NIM | | NIM |
| Above | 0.46 (0.29, 0.72) | | | | | |
| Unknown | 0.44 (0.29, 0.67) | | | | | |
| Age: Reference: 18 to 33 | | NIM | | NIM | | < 0.01 |
| 33 to 48 | | | | | 0.81 (0.67, 0.97) | |
| 48 and older | | | | | 0.63 (0.51, 0.78) | |
| | Other | treatmen | t procedure category | | | |
| Race/Ethnicity: Reference: | White | 0.88 | | 0.25 | | 0.55 |
| African-American | 0.86 (0.51, 1.45) | | 1.06 (0.71, 1.58) | | 0.82 (0.63, 1.07) | |
| Hispanic | 0.74 (0.25, 2.13) | | 1.13 (0.62, 2.07) | | 0.80 (0.50, 1.28) | |
| Other | 0.63 (0.18, 2.11) | | 0.65 (0.29, 1.48) | | 0.78 (0.46, 1.32) | |
| Unknown | 1.08 (0.54, 2.14) | | 1.53 (0.99, 2.36) | | 0.94 (0.73, 1.20) | |
| Insurance: Reference: Pub | lic | NIM | | 0.01 | | < 0.01 |
| Private | | | 0.78 (0.42, 1.44) | | 1.07 (0.75, 1.55) | |
| Out-of-pocket | | | 0.79 (0.56, 1.12) | | 1.01 (0.81, 1.26) | |
| Unknown | | | 0.36 (0.21, 0.64) | | 0.25 (0.17, 0.38) | |
| Poverty status: Reference: | Below | NIM | | 0.02 | | NIM |
| Above | | | 0.76 (0.50, 0.16) | | | |
| Unknown | | | 0.58 (0.39, 0.85) | | | |
| Age: Reference: 18 to 33 | | NIM | | NIM | | < 0.01 |
| 33 to 48 | | | | | 0.70 (0.56, 0.87) | |
| 48 and older | | | | | 0.59 (0.46, 0.76) | |

Table 4 Continued

NIM, not included in the final model.

includes both urban and rural poor and all types of payers. This study documents racial/ethnic variations in the provision of dental procedures for patients seen at a dental teaching institution, there is also the need to investigate the impact on dental health outcomes of racial/ethnic variation in the provision of dental procedures. Racial/ethnic variation should be recognized as a serious problem within oral health disparities and, if not addressed, will impact the overall life quality for those groups most affected.

Conclusions

Substantial racial/ethnic variation in the use of dental treatment pro-

cedures exists for African-Americans and Hispanics. African-Americans were less likely to have restorative procedures and more likely to have surgery procedures than the White population. This study presents findings that support further investigation into racial/ethnic disparities in the provision of dental procedures in other dental settings.

Acknowledgments

We wish to thank Nick Pajewski for all his statistical advice and Patrick Lynch for his assistance with data collection.

References

1. US Department of Health and Human Services. Oral health in America. A report of the surgeon general. Rockville (MD): HHS, National Institutes of Health, National Institute of Dental and Craniofacial Research; 2000.

- Institute of Medicine. Unequal treatment: confronting racial and ethnic disparities in health care. Smedley BD, Stith AY, Nelson AR, editors. Washington, DC: National Academies Press; 2002
- Peterson ED, Wright SM, Daley J, Thibault GE. Racial variation in cardiac procedure use and survival following acute myocardial infarction in the department of veterans affairs. JAMA. 1994;271: 1175–80.
- Ellison GL, Warren JL, Knopf KB, Brown ML. Racial differences in the receipt of bowel surveillance following potentially curative colorectal cancer surgery. Health Serv Res. 2003;38:1885–903.
- Fiscella K, Franks P, Gold MR, Clancy CM. Inequality in quality addressing socioeconomic, racial, and ethnic disparities in health care. JAMA. 2000;283: 2579–84.

- The Henry J Kaiser Family Foundation. Race, ethnicity and family care: a survey of public perceptions and experiences. Menlo Park (CA): Henry J Kaiser Family Foundation; 1999.
- Gilbert GH, Shelton BJ, Duncan RP. Use of specific dental treatment procedures by dentate adults during a 24-month period. Community Dent Oral Epidemiol. 2002;30:260–76.
- 8. Cabral ED, Caldas AF Jr., Cabral HAM. Influence of the patient's race on the

dentist's decision to extract or retain a decayed tooth. Community Dent Oral Epidemiol. 2005;33:461–6.

- Kressin NR, Boehmer U, Berlowitz D, Christiansen C, Pitman A, Jones J. Racial variation in dental procedures: the case of root canal therapy versus tooth extraction. Med Care. 2003;41:1256–61.
- Manski RL, Moeller JF. Use of dental services an analysis of visits, procedures and provides, 1996. JADA. 2002;133: 167–75.
- Gregg GH, Shewchuk RM, Litaker MS. Effect of dental practice characteristics on racial disparities in patient-specific tooth loss. Med Care. 2006;44:414–20.
- Marcus SE, Drury TF, Brown LJ, Zion GR. Tooth retention and tooth loss in the permanent dentition of adults: United States, 1988–1991. J Dent Res. 1996;75: 684–95.

Professor & Chairperson – Dept of Dental Public Health Sciences Temple University School of Dentistry

Temple University School of Dentistry invites applicants for a full-time, tenure-track (Presidential) appointment as Professor and Chairperson of the newly established Department of Dental Public Health Sciences, available immediately. Applicant is expected to be nationally recognized as a leader in the dental profession, and must possess a dental degree, graduate education and experience relevant to the field of dental public health, outstanding written and verbal communication skills, and administrative experience in dental public health at a senior level. Salary will be commensurate with qualifications and experience in dental public health. Preference will be given to applicants with previous academic teaching and administrative experience at a senior level in a dental school setting, board certification in dental public health, and an established record of scholarship in dental public health as evidenced by peer-reviewed publications and/or extramural grant support. Dental licensure in the Common-wealth of Pennsylvania, or eligibility for the practice of dentistry for the limited purpose of teaching at the School of Dentistry, is required.

Responsibilities will include teaching, research, and overall administrative management of the School of Dentistry's Department of Dental Public Health Sciences. Graduates of non-US/Canadian dental schools who successfully pass Part I and II of the National Board Dental Examinations may be eligible for admission to a faculty DMD program at the School of Dentistry after one year of service. As soon as possible, applicant should submit a cover letter, a curriculum vitae signed and dated on each page, and names and addresses of at least three references to:

Dr. Thomas E. Rams Chair, Dental Public Health Sciences Search Committee Temple University School of Dentistry 3223 North Broad Street Philadelphia, PA 19140 of applications will begin immediately and continue unt

Review of applications will begin immediately and continue until the position is filled. Temple University is an equal opportunity/affirmative action employer. Minorities and women are encouraged to apply.